

## Letter To The Editor

## Noncompaction Requires Family Studies, Work-Up for Neuromuscular Disorders, and Close Monitoring

Josef Finsterer, MD, PhD.<sup>1</sup>, Claudia Stöllberger, MD.<sup>2</sup>

<sup>1</sup>Krankenanstalt Rudolfstiftung Messerli Institute Veterinary University of Vienna Vienna Austria.

<sup>2</sup>Medical Department Krankenanstalt Rudolfstiftung, Vienna Austria.

\*Corresponding Author  
Josef Finsterer, MD, PhD

**Keywords:** Diagnosing LVHT. hypertrabeculation / noncompaction (LVHT).

In a recent article Erdogan *et al.*, reported about a 55yo male with sudden onset anginal chest pain and mild systolic dysfunction in whom left ventricular hypertrabeculation / noncompaction (LVHT) was detected on echocardiography and anginal chest pain attributed to myocardial bridging of the posterior descending artery on multidetector computed tomography (Erdogan, E. *et al.*, 2013). We have the following comments and concerns.

Though LVHT is regarded as a congenital disorder in the majority of the cases, it may develop during postnatal life in rare cases as well (acquired LVHT) (Finsterer, J. *et al.*, 2008). Were any previous echocardiographic examinations available and reviewed, particularly intrauterine ultrasound to confirm the congenital nature of LVHT in this patient?

Diagnosing LVHT has significant therapeutic implications not only in terms of heart failure but also in terms of ventricular arrhythmias and cardio-embolic risk (Caliskan, K. *et al.*, 2012). Since LVHT patients carry an increased risk of ventricular arrhythmias with sudden cardiac death and of cardio-embolic events, it is essential to closely monitor these patients for arrhythmias and for the development of intertrabecular thrombus formation. In patients with a cardio-embolic event and LVHT, oral anticoagulation is indicated. Was the individual history positive for stroke /embolism, did the patient ever undergo cerebral MRI to look for previous embolic ischemias?

Since LVHT frequently occurs also in other family members, it is worthwhile to carry out family

studies in case LVHT is diagnosed. Interestingly, relatives of LVHT patients may also present with cardiomyopathy other than LVHT. Were echocardiographies carried out in first-degree relatives of the described patient?

Since the patient presented with two presumably congenital abnormalities and since LVHT is frequently associated with chromosomal abnormalities or mutations in genes encoding proteins of the contractile apparatus or the sarcolemma (Finsterer, J. 2009), it would be interesting to know if investigations for chromosomal aberrations were carried out. Was consanguinity excluded in the family of the patient?

Since LVHT is even more frequently associated with neuromuscular disorders there is a need to refer these patients to the myologist for neuromuscular evaluation. Were there any clinical indications, such as muscle weakness or wasting, muscle cramping, easy tiredness, joint contractures or gait disturbance, for neuromuscular disease in the presented patient? It is also essential to ask LVHT patients for sensory disturbances since it has been described even in association with hereditary neuropathy (Corrado, G. *et al.*, 2006).

So far, LVHT has been reported in association with mutations in the TAZ, DTNA, ZASP, lamin A/C, MYH7, MYH7B, ACTC1, TNNT2, TNNI3, MYBPC3, TPM1, dystrophin, DMPK, ZNF9, LAMP2, GAA, mtDNA genes, AMPD1, GBE1, RYR1, COL7A1, PMP22, MMACHC,  $\beta$ -globin, and DNAJC19 genes.

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easims/>

Article History

Received: 18.02.2019

Accepted: 27.02.2019

Published: 14.03.2019

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

Was he tested for mutations in any of these genes? Was the family history positive for hereditary disease?

Overall, this interesting case needs further individual and familial investigations to address these issues raised above and close monitoring to prevent fatal complications of LVHT.

#### REFERENCES

1. Erdogan, E., Akkaya, M., Bacaksiz, A., Tasal, A., & Sevgili, E. (2013). A very rare case of coexistence of ventricular noncompaction cardiomyopathy, myocardial bridging and atherosclerosis. *Postępy w Kardiologii Interwencyjnej= Advances in Interventional Cardiology*, 9(2), 176.
2. Finsterer, J., Stöllberger, C., & Schubert, B. (2008). Acquired left ventricular noncompaction as a cardiac manifestation of neuromuscular disorders. *Scandinavian Cardiovascular Journal*, 42(1), 25-30.
3. Caliskan, K., Ujvari, B., Bauernfeind, T., AMJ THEUNS, D. O. M. I. N. I. C., Van Domburg, R. T., Akca, F., ... & SZILI-TOROK, T. A. M. A. S. (2012). The prevalence of early repolarization in patients with noncompaction cardiomyopathy presenting with malignant ventricular arrhythmias. *Journal of cardiovascular electrophysiology*, 23(9), 938-944.
4. Finsterer, J. (2009). Cardiogenetics, neurogenetics, and pathogenetics of left ventricular hypertrabeculation/noncompaction. *Pediatric cardiology*, 30(5), 659.
5. Corrado, G., Checcarelli, N., Santarone, M., Stöllberger, C., & Finsterer, J. (2006). Left ventricular hypertrabeculation/noncompaction with PMP22 duplication-based Charcot-Marie-Tooth disease type 1A. *Cardiology*, 105(3), 142-145.