

Letter to Editor

Comprehensive Clinical and Genetic Work-Up in Carriers of the Variant C.1228G>A In *DNM1L* Is Warranted

Josef Finsterer, MD, PhD¹, Scorza Fulvio A, MD², Scorza Carla A, MD², Fiorini Ana C, MD³¹Krankenanstalt Rudolfstiftung, Messerli Institute, Vienna, Austria²Disciplina de Neurociência. Universidade Federal de São Paulo/Escola Paulista de Medicina (UNIFESP/EPM). São Paulo, Brasil³Programa de Estudos Pós-Graduado em Fonoaudiologia, Pontifícia Universidade Católica de São Paulo (PUC-SP), Brazil; Departamento de Fonoaudiologia, Escola Paulista de Medicina/Universidade Federal de São Paulo (EPM/UNIFESP), São Paulo, Brazil**Article History**

Received: 18.02.2020

Accepted: 20.03.2020

Published: 25.03.2020

Abstract:**Journal homepage:**<https://www.easpublisher.com/easjms>**Quick Response Code****Keywords:** DNM1L, cardiomyopathy, heart failure, mitochondrial disorder, multisystem.

Copyright @ 2020: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

LETTER TO THE EDITOR

In a recent article, Vandeleur *et al.*, reported about an 8 months-old female carrying the variant c.1228G>A in *DNM1L* which phenotypically manifested with microcephaly, failure to thrive, poor feeding, exsiccosis, cardiomyopathy (CMP), developmental motor delay, truncal hypotonia, exaggerated startle reaction, reduced visual tracking, myopathy, and lactic acidosis (Vandeleur, D. *et al.*, 2019). We have the following comments and concerns.

We do not agree that cardiac involvement in *DNM1L*-related disease “is undescribed” (Vandeleur, D. *et al.*, 2019). The *DNM1L*-gene encodes the dynamin-related protein-1 (DRP1). It is well known that DRP1 stimulates mitochondrial respiration, bioenergetics, and ROS-signalling in cardiomyocytes (Zhang, H. *et al.*, 2017). In mice with *Drp1* deficiency disassembly of the protein impairs mitophagy, leading to reduction of the mitochondrial membrane potential, aberrant calcium handling, reduced ATP synthesis,

myocardial inflammation, and lastly heart failure (Cahill, T. J. *et al.*, 2015).

There is a discrepancy concerning the cerebral myelination of the index patient. On the one hand the authors describe that myelination was normal for age (at age 6 months) on MRI and later in the case description they mention that there was hypomyelination (at age 8 months) at autopsy. We should know what is true and if the discrepancy results from evaluation at two different time points, is attributable to the disease course, or due to different types of investigation.

Missing is the specification of the CMP. It is described that the left ventricle was hypertrophic and dilated at the same time (Vandeleur, D. *et al.*, 2019). However, no information is provided if CMP was hypertrophic CMP with secondary dilatation, if CMP was primary dilative with secondary hypertrophy, if there was left ventricular hypertrabeculation (LVHT) / noncompaction, or if it was recurrent Takotsubo syndrome (TTS). Specifying the type of CMP is crucial as it may strongly determine the type of treatment and

the outcome. Arguments for the presence of LVHT are the subendocardial fibrosis (Stöllberger, C., & Finsterer, J. 2019), frequently found in LVHT, and the mitochondrial disorder (MID), as LVHT is most frequently associated with MIDs (Finsterer, J. 2009). Missing are the troponin and proBNP values, long-term ECG recordings, and the information about the course of heart failure. We should know if heart failure completely resolved after the first episode or not. Was pulmonary embolism excluded as a cause of clinical deterioration during blood transfusion?

Missing is the information if the parents were consanguineous or not and if the siblings and the parents were genetically tested for the culprit mutation.

Since the patient had myopathy we should know if respiratory insufficiency was truly only due to heart failure or if there was additional or exclusive muscular respiratory failure.

We should know if the increased startle reaction was attributable to seizure activity, thus, if the EEG showed epileptiform discharges or not.

Overall, this interesting case report has a number of shortcomings which need to be addressed before drawing final conclusions. We should know if the mutation was inherited or occurred spontaneously and evaluation of the phenotype needs to be reinforced.

REFERENCES

1. Cahill, T. J., Leo, V., Kelly, M., Stockenhuber, A., Kennedy, N. W., Bao, L., ... & Bellahcene, M. (2015). Resistance of dynamin-related protein 1 oligomers to disassembly impairs mitophagy, resulting in myocardial inflammation and heart failure. *Journal of Biological Chemistry*, 290(43), 25907-25919.
2. Finsterer, J. (2009). Cardiogenetics, neurogenetics, and pathogenetics of left ventricular hypertrabeculation/noncompaction. *Pediatric cardiology*, 30(5), 659.-681.
3. Stöllberger, C., & Finsterer, J. (2019). Understanding left ventricular hypertrabeculation/noncompaction: pathomorphologic findings and prognostic impact of neuromuscular comorbidities. *Expert review of cardiovascular therapy*, 17(2), 95-109.
4. Vandeleur, D., Chen, C. V., Huang, E. J., Connolly, A. J., Sanchez, H., & Moon-Grady, A. J. (2019). Novel and lethal case of cardiac involvement in DNMI1 mitochondrial encephalopathy. *American Journal of Medical Genetics Part A*, 179(12), 2486-2489.
5. Zhang, H., Wang, P., Bisetto, S., Yoon, Y., Chen, Q., Sheu, S.S., & Wang, W. (2017). A novel fission-independent role of dynamin-related protein 1 in cardiac mitochondrial respiration. *Cardiovasc Res*, 113, 160-170.