

Research Article

Environmental impact of some poisonous weeds in Damietta, Egypt

Mamdouh Mohamed S. Serag¹ Amina Zakreya AboEl-Naga¹ and Mohamed Adel Mohamed El-Gendey¹¹Botany and Microbiology Department, Faculty of Science, Damietta University, Egypt

*Corresponding Author

Mohamed Adel Mohamed El-Gendey

Abstract: Toxic plant contains a number of toxic alkaloids produced as secondary metabolites, which are a characteristic feature of plants, are especially important and can protect plants against a wide variety of microorganisms. The antibacterial effect of aqueous and organic solvents (ethanol and acetone) extracts of four poisonous weeds (*Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*) obtained from Damietta province of Egypt, were tested in vitro against growth of bacteria, *Escherichia coli* and *Helicobacter pylori*. Alkaloid can use in antibacterial activity. Bacterial drug resistant was important problem against antibiotic so researcher going to used alkaloid extract as antibacterial activity. The effect of alkaloids of four plant have inhibitors of growth of bacteria (*Escherichia coli* and *Helicobacter pylori*). Results for alkaloid content in the plants (*Datura stramonium* (all plant), *Datura stramonium* (Seed), *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*) have 150.5±6.1, 103.1±4.2, 30.7±2.4, 42.3±1.9 and 88.6±6.3 mg/g, respectively. The study confirms that all plants extracts used in this investigation possess *in vitro* antibacterial activity against the used organisms. The efficiency of the extracts varied with, solvent used in the extraction as well as, plant species and the part of plant used.

Keywords: Toxic plant, alkaloids, antibacterial activity.

INTRODUCTION

A wide variety of toxic plants produces compounds that may cause clinical symptoms in humans, some of which have caused severe poisonings. A few plants give rise to serious poisoning after ingestion even of a limited amount of plant material (Frohne *et al.*, 2004). Alkaloid extracts of the leaves of plants known to have toxic effects when ingested by livestock were screened for biological activity (McGaw *et al.*, 2005). Isolation of active compounds, in almost all cases, provided scientific validation for the use of the plants in traditional medicine. Although plants used medicinally are widely assumed to be safe, many are potentially toxic (Fennell *et al.*, 2004). On the basis that toxic plants have proven pharmacological activity. Resistance in microbes against antibiotics is a worldwide problem that is caused because of frequent exposure of antibiotics. There are millions of chemical compounds that are being synthesized, thousands of which have been confirmed for their antimicrobial potential (Tim *et al.*, 2014). Infectious diseases caused by these pathogenic microorganisms have become an important cause of death and mortality in immune-compromised patients in developing countries (Ara *et*

al., 2009). Despite the availability of a wide range of antibiotics, bacteria are constantly developing resistance to these antibiotics (AL-Bari *et al.*, 2006).

The present investigation aimed to study the effects of some extracts obtained from these four plants using different solvent (ethanol and acetone), on growth of some strains of *Escherichia coli* and *Helicobacter pylori*.

MATERIALS AND METHODS

Study Area

Damietta Province is located in the downstream part of the Damietta branch of the River Nile at 31° 25' 10" north to 31° 48' 54" east N-32° 00' longitude to the north east of the Nile Delta region of Egypt. The coast of Damietta Governate extends from El-Deeba village (about 20 km from Port-said) to Gamasa at west along the Mediterranean Sea for about 42 Km. This province is bounded by Lake Manzala at the east, Mediterranean sea from the north and El-Dakahlia Governate from the west and the south. The total average area of Damietta Province is about 1029

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easjals/>

Article History

Received: 12.09.2019

Accepted: 23.09.2019

Published: 10.10.2019

Copyright © 2019 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

DOI: 10.36349/easjals.2019.v02i10.004

cultured on brain heart infusion agar.

Preparation of plant extract

Antibacterial activities of different extracts from the four plant (*Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*) were used. The diluted extracts were prepared by using Dimethyl Sulfoxide (DMSO) to obtain the optimum dose concentrations of the different plant extract. DMSO was used as negative control. The antibacterial activity of each extracts were tested by Hole diffusion method (Al Somal *et al.*, 1994). Muller Hinton agar plates were inoculated by rubbing sterile cotton swabs after immerse 100 µl bacterial suspensions on plates (overnight cultures grown at 37°C on nutrient agar and adjusted to 0.5 McFarland in sterile saline) over the entire surface of the plate. After inoculation, 9 mm diameter wells were cut into the surface of the agar using a sterile cork borer. The different dose concentrations were added to the wells. Plates were incubated at 37°C for 24 h. Control wells contained DMSO. Bacterial growth inhibition was determined as the diameter of the inhibition zones around the discs. Zones of inhibition were measured by using ruler. The diameter of zones was recorded. Each assay was carried out in triplicate.

Soil sampling and analysis

The sediments were collected from the bottom of the studied stations and were carried to the laboratory in a plastic bag. Shortly after the collection, the samples were spread over large glass plate; air dried, thoroughly mixed, and passed through 2 mm sieve to remove gravel and debris and then packed in plastic bags ready for analysis.

Methods of Digestion Soil

Digestion procedures were applied to digest soil, involving nitric acid-hydrogen peroxide, nitric acid-sulfuric acid, and nitric acid-perchloric acid mixtures. Add 0.05 g of soil and heated in hot plate, after completed digested completed to 20 ml by deionized H₂O.

Apparatus

Flame Atomic Absorption Spectrometer (Thermo Electron Corporation-S series) with deuterium lamp background Correction was used.

Determination of heavy metals in soil

Final concentrations of the metals in the soil samples were calculated using the following formula (Uwah *et al.*, 2012).

$$\text{Concentration (mg/kg)} = \frac{\text{Concentration (mg/L)} \times V}{W}$$

Where V = Final volume (20 ml) of solution, and W = Initial weight (0.05 g) of sample measured.

Statistical Analysis

Data were analyzed using analysis of variance (SPSS, version 10.0, SPSS Inc., Chicago, Ill.). Means were separated with Duncan's multiple range tests (Kleinbaum *et al.*, 1998 ; Moussa and Amira, 2017).

RESULTS AND DISCUSSION

Total Alkaloid Content

Alkaloid of toxic plants is directly linked to the wide range of chemical compounds synthesized by the various biochemical pathways. These compounds are classified as secondary plant products, because they are not much related to the plant's survival. The results for alkaloid content in the different plant *Datura stramonium* (all plant), *Datura stramonium* (Seed), *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis* have total alkaloid contents 150.5±6.1, 103.1±4.2, 30.7±2.4, 42.3±1.9 and 88.6±6.3 mg/g, respectively as showed in Figure (1). The content of alkaloid can use as antimicrobial (Tim *et al.*, 2014; Kim *et al.*, 2004). Results showed that: (i) screening of presence of alkaloid confirmed the presence of alkaloid in the plant organs (ii) The alkaloid contents of the plant organs of the plant species were significantly different from one another and (iii) The total alkaloid content of *Datura stramonium* was the highest among the four plant species.




Fig. 2. Total alkaloid contents from four plants, *Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*.

Antibacterial activity of alkaloid extract of toxic plant

As shown in Tables (1, 2 and 3), the antibacterial activity showed positive response in all fractions of four plants extract (acetone and ethanol) as compared with antibiotic Amoxicillin, Clarithromycin and Trimethoprim. From this result we show that inhibition zone *E. coli* and *H. pylori* have good result can be improvement in future study. The antibacterial susceptibility tests showed that the alkaloid extract had potent antibacterial properties (Stavri *et al.*, 2007; Tim *et al.*, 2014).

Table 1. Antibacterial activity of Acetone extracted from four plants (*Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*).

Plant	Conc.%	Bacteria	Zone of inhibition (mm)
<i>Datura stramonium</i> (shoot, leaf and root)	13	<i>Escherichia coli</i>	8±0.05
	15	<i>Helicobacter pylori</i>	9±0.21
<i>Datura stramonium</i> (Seed)	22	<i>Escherichia coli</i>	7±0.11
	18	<i>Helicobacter pylori</i>	9±0.03
<i>Chenopodium murales</i>	25	<i>Escherichia coli</i>	8±0.02
	27	<i>Helicobacter pylori</i>	9±0.01
<i>Lotus corniculatus</i>	26	<i>Escherichia coli</i>	8±0.03
	24	<i>Helicobacter pylori</i>	12±0.09
<i>Amaranthus viridis</i>	21	<i>Escherichia coli</i>	9±0.07
	18	<i>Helicobacter pylori</i>	10±0.06

Each assay in these experiments was repeated three times and the results (mm of zone of inhibition) were expressed as average values ± standard deviation. Mean inhibition zone diameter (mm) after 24 h of incubation.

Table 2. Antibacterial activity of Ethanol extracted from four plants (*Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*).

Plant	Conc.%	Bacteria	Zone of inhibition (mm)
<i>Datura stramonium</i> (shoot, leaf and root)	15	<i>Escherichia coli</i>	7±0.07
	17	<i>Helicobacter pylori</i>	10±0.11
<i>Datura stramonium</i> (Seed)	20	<i>Escherichia coli</i>	5±0.09
	19	<i>Helicobacter pylori</i>	7±0.18
<i>Chenopodium murales</i>	23	<i>Escherichia coli</i>	6±0.22
	22	<i>Helicobacter pylori</i>	8±0.05
<i>Lotus corniculatus</i>	25	<i>Escherichia coli</i>	6±0.06
	22	<i>Helicobacter pylori</i>	11±0.11
<i>Amaranthus viridis</i>	20	<i>Escherichia coli</i>	8±0.07
	17	<i>Helicobacter pylori</i>	7±0.13

Each assay in these experiments was repeated three times and the results (mm of zone of inhibition) were expressed as average values ± standard deviation. Mean inhibition zone diameter (mm) after 24 h of incubation.

Table 3. Antibacterial activity of the antibiotic (Amoxicillin, Clarithromycin and Trimethoprim)

Antibiotic	Bacteria	Zone of inhibition (mm)
Amoxicillin	<i>Escherichia coli</i>	10±0.08
	<i>Helicobacter pylori</i>	13±0.05
Clarithromycin	<i>Escherichia coli</i>	8±0.11
	<i>Helicobacter pylori</i>	14±0.08
Trimethoprim	<i>Escherichia coli</i>	13±0.09
	<i>Helicobacter pylori</i>	8±0.21

Each assay in these experiments was repeated three times and the results (mm of zone of inhibition) were expressed as average values ± standard deviation. Mean inhibition zone diameter (mm) after 24 h of incubation.

Table 4. Plant species, life form, Life span, floristic category, and families associated with four plants (*Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*) in the Nile Delta coast.

Plant	Life form	Life span	Floristic category	Family
-------	-----------	-----------	--------------------	--------

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easjals/>

Article History

Received: 12.09.2019

Accepted: 23.09.2019

Published: 10.10.2019

Copyright © 2019 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

DOI: 10.36349/easjals.2019.v02i10.004

<i>Avena fatua</i>	Therophyte	Annuals	ME+MED	Poaceae
<i>Imperata cylindrica</i>	Graminoid	Perennials	+NEO	Poaceae
<i>Chenopodium murale</i>	Therophyte	Annuals	COSM	Amaranthaceae
<i>Tamarix nilotica</i>	Phanerophyte	Perennials	ME+SA-AR	Tamaricaceae
<i>Cynodon dactylon</i>	Geophytes	Perennials	ME+IT	Asclepiadaceae
<i>Juncus acutus</i>	Cryptophytes	Perennials	ME+ES+IT ME+IR-TR+ER-ES+SU	Juncaceae
<i>Malva parviflora</i>	Therophyte	Annuals	ME+ER	Malvaceae
<i>Rumex dentatus</i>	Therophyte	Annuals	ME+IT+ES	Polygonaceae
<i>Sonchus oleraceus</i>	Therophyte	Annuals	COSM	Asteraceae
<i>Cyperus rotundus</i>	Geophytes	Perennials	PAN	Cyperaceae
<i>Spergularia marina</i>	Therophyte	biennial	ER-R+ME+IR-TR	Caryophyllaceae
<i>Cynanchum acutum</i>	Phanerophyte	Perennials	ME+IR-TR+ER-SR	Asclepiadaceae
<i>Echinochloa stagnina</i>	Therophyte	Perennials	PAL	Poaceae
<i>Bassica indica</i>	Therophyte	Annuals	ME+ER	Chenopodiaceae
<i>Urospermum picroides</i>	Therophyte	Annuals	ME+ER	Asteraceae
<i>Beta vulgaris</i>	Hemicryptophytes	Annuals	ME+ER-SR+IR-TR	Amaranthaceae
<i>Urtica urens</i>	Therophyte	Annuals	ME+ER	Urticaceae
<i>Vicia sativa</i>	Therophyte	Annuals	Me+IR+-IR+SA-AR	Fabaceae
<i>Mesembryanthemum nodiflorum</i>	Therophyte	Annuals	ME+ER	Aizoaceae
<i>Solanum lycopersicum</i>	Hemicryptophytes	Perennials	IR-TR+ER-SR+ME	Poaceae
<i>Plantago major</i>	Hemicryptophytes	Perennials	COSM	Plantaginaceae
<i>Leptochloa fusca</i>	Raunkiaer	Perennials	PAL	Poaceae
<i>Solanum nigrum</i>	Therophyte	Annuals	ME+ER-SR+IR-TR	Solanaceae
<i>Rumex pectus</i>	Therophyte	Annuals	ME+SA-AR	Polygonaceae
<i>Ipomoea carnea jacq</i>	Chamaephytes	Perennials	ME+IT+ES	Convolvulaceae

Life form

The plants species were classified according to Raunkiaer (1937) classification into seven life form classes. there were represented in the present work as therophytes 56%, ge-helophytes 8% , chamaephytes 4%, hemicryptophytes 12%, cryptophytes 8%, and phanerophytes 12% (Table 4).

Plant life span

The 25-recorded can be classified based on life span into three groups: perennials 44% ,annuals 52% and biennials 4% . most of the recoded species are perennials and annuals with few biennials. These results are consistent with those of Boulos (1999 & 2005) as shown in (Table 4).

Table 5. Summary of heavy metal concentration of soils within 0–20 Cm depth in different sites

metal	site 1	site 2	site 3	site 4	site 5	site 6	site 7	site 8	site 9	site 0
Cu	16	9.6	20	51.6	27.6	27.2	12.4	13.6	12.4	12
Co	80	68.8	74.4	74	77.2	76.8	69.6	80	80.8	80.8
Ni	61.6	46.4	62.8	63.6	60.8	57.2	48	58	56.4	54.8
Zn	110.8	107.6	186.4	156.4	230	151.2	100	139.2	89.6	116
Mg	2664	2137.2	1231.6	848	1248.8	806	61.2	642.4	284.4	1159.6
Fe	351.6	214.8	686.4	732.4	772.8	654.8	297.6	462	256.4	116
Pb	184.8	179.6	176.4	190.4	196.8	185.6	216	218	229.6	220
Cd	23.6	22	19.2	24	26.8	24.4	23.6	28	29.2	26

The data of heavy metal concentration of soils within 0–20 Cm depth in different sites are shown in Table (5) which indicates that there is a variable changes in the concentration of the metals in the different sites.

CONCLUSION

Use of toxic plant alkaloid extract for antimicrobial compounds was important of humanity is necessitated by the inherent ability of pathogens to develop resistance against antibiotics. Potentially harmful side effects associated with use of new chemical entities synthesized artificially and the unsustainably high costs of drug development are slowly shifting the focus to plant derived

phytochemicals of medicinal significance (Scott *et al.*, 2009).

Extracts of alkaloids from (*Datura stramonium*, *Chenopodium murales*, *Lotus corniculatus* and *Amaranthus viridis*) were found to be inhibitors of growth in bacteria (*Escherichia coli* and *Helicobacter pylori*), the standard antibiotic (Amoxicillin, Clarithromycin and Trimethoprim) used in that study to compared with extract of alkaloid (four plant).

REFERENCES

1. Al, s. n., Coley, K.E., Molan, P.C & Hancock, B.M. (1994). Susceptibility of *Helicobacter pylori*

- to the antibacterial activity of manuka honey. *J R Soc Med* 87: 9-12.
2. AL-Bari, M.A., Sayeed, M.A., Rahman, M.S., & Mossadik, M.A. (2006). "Characterisation and antimicrobial activities of a phenolic acid derivative produced by *Streptomyces bangladesiensis* a novel species collected in Bangladesh." *Research Journal of Medicine and Medical Sciences* (1), 77–81, 2006.
 3. Ara, N., Nur, M.H., Amran, M.S., Wahid, M.I.I., & Ahmed, M. (2009). "In vitro antimicrobial and cytotoxic activities of leaves and flowers extracts from *Lippia alba*," *Pakistan Journal of Biological Sciences*, vol. 12, (1), pp. 87–90, 2009.
 4. Boulos, L. (1999). *Flora of Egypt. Volume I.* Al Hadara Publishing Cairo, Egypt.
 5. Boulos, L. (2005). *Flora of Egypt. Volume VI.* Al Hadara Publishing Cairo, Egypt.
 6. Fennell, C.W., Lindsey, K.L., McGaw, L.J., Sparg, S.G., Stafford, G.I, Elgorashi, E.E., & Van, S.J. (2004). Assessing African medicinal plants for efficacy and safety: pharmacological screening and toxicology. *Journal of ethnopharmacology*, 94(2-3), 205-217.
 7. Frohne, D., Pfänder, H.J. (2004). eds *Poisonous plants*, 2nd ed. New Mexico: Manson Publishing.
 8. Helal, R. M., & Amira, M.H. (2017). Selenium Affects Physiological Responses of *Phaseolus vulgaris* in Response to Salt Level. *INTERNATIONAL JOURNAL OF VEGETABLE SCIENCE*, 1-19.
 9. Kim, S.H., Shin, D.S., Oh, M.N., Chung, S.C., Lee, J.S & Oh, K.B. (2004). Inhibition of the bacterial surface protein anchoring transpeptidase sortase by isoquinoline alkaloids. *Biosci Biotechnol Biochem* 2004; 68,421–4.
 10. Kleinbaum ,D.G.,Kupper, L.L, & Muller, K.E. (1998). One-way analysis of variance. In: *Multivariate methods*, pp. 341-386. Boston: PWS-Kent Publishing Company.718.
 11. Mashaly, I. (2001). Contribution to the Ecology of the Deltaic Mediterranean Coast, Egypt. *Online J. Biol. Sci.*, 1(7), 628-635.
 12. McGaw, L.J., Eloff, J.N., & Meyer, J.M .(2005). Screening of 16 poisonous plants for antibacterial, anthelmintic and cytotoxic activity in vitro. *South African Journal of Botany*, 71(3-4), 302-306.
 13. Raunkiaer, C. (1937). *Plant life forms*. Clarendon, Oxford, pp;104.
 14. Shamsa, F., Monsef, H., Ghamooshi. R., Verdianrizi, M. (2008). Spectrophotometric determination of total alkaloids in some Iranian medicinal plants. *Thai J Pharm Sci.* 32;17–20.
 15. Sreevidya, N., & Mehrotra, S. (2003). Spectrophotometric method for estimation of alkaloids precipitable with Dragendorff's reagent in plant materials. *Journal of AOAC international*, 86(6), 1124-1127.
 16. Tim,C.T.P., Benjamart, C., & Andrew, J. L. (2014). Alkaloids: An overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *International Journal of Antimicrobial Agents* 44 (2014) 377–386.
 17. Trease, G.E., & Evans ,W.C.(2002). *Pharmacognosy*. 15th ed. Philadelphia: WB Saunders, Elsevier Science Limited; p. 336.
 18. Uwah, E.I., Buba, A., Gwaski, P.A. (2012). Heavy Metal Levels in Roots and Leaves of some Vegetables Grown in Gwoza, Borno State, Nigeria; *Book of Proceedings of the 35th Annual International Conference, Workshop & Exhibition of Chemical Society of Nigeria (CSN); (Heartland 2012) 17th-21st September, 2012; 1: 132 –134.*