

Original Research Article

Determination the Aetiological Pattern of Head Injury in the Region

Md. Jasim Uddin¹, Md. Shafi Mohaimen², Md. Iqbal Bahar Chowdhury³, Md. Nazir Hossain⁴, Rakibul Hasan Khan⁵, Md. Mashioor Rahman Rikabder⁶, Afroza Akter⁷, Malay Kumar Das⁸

¹Assistant Professor, Department Of Forensic Medicine, Ashiyan Medical College, Dhaka, Bangladesh

²Assistant Professor, Dept. of Forensic Medicine, Shaheed Tajuddin Ahmad Medical College, Gazipur, Bangladesh

³Assistant Professor, Department Of Forensic Medicine, Shaheed Monsur Ali Medical College, Dhaka, Bangladesh

⁴Assistant Professor, Department Of Forensic Medicine, Dhaka National Medical College, Dhaka, Bangladesh

⁵Lecturer, Department Of Forensic Medicine, Sheikh Hasina medical college, Tangail, Bangladesh

⁶Assistant Professor, Department Of Forensic Medicine, Jahurul Islam Medical College, Kishoreganj, Bangladesh

⁷General Practitioner, Hazi Md. Alfaj Uddin Diagnostic Centre, Kanchan, Rupgonj, Narayanganj, Bangladesh

⁸Assistant Professor, Department Of Forensic Medicine, Ad-Din Akij Medical College, Khulna, Bangladesh

Article History

Received: 29.09.2021

Accepted: 03.11.2021

Published: 08.11.2021

Journal homepage:

<https://www.easpublisher.com>

Quick Response Code



Abstract: *Introduction:* Earlier studies from the developing countries as well as from other terrains of Bangladesh showed various aetiologies of head injuries. However, there is no substantive government data's are available regarding the cause and manner of head injury. *Objective:* The current study was conducted with an objective to determine the aetiological pattern of head injury in the region. *Materials & Methods:* The study was conducted in Dept. of forensic medicine, Rajshahi Medical College Hospital, Rajshahi, Bangladesh from June-2019 to July-2021. Cases of fatal head injuries irrespective of their aetiology that were brought for medicolegal autopsies over a period of two years were included in this study. The data with regard to their age, sex, mode of injury in detail was collected and interpreted. *Results:* Out of total 59 cases of fatal head injury who succumb to death, 90% were males. Maximum numbers of cases (27.11%) were in age group of 21-30 followed by 31-40 year (22.03%). Roadside accident was the leading cause of head injury (65%) while only 15% were as a result of assault. Two wheeler riders comprises of 35.31% followed by pedestrians (24.3%). Maximum number of fatalities due to head injury was due to roadside accident and that too in a productive age group of 21- 40 years. Rider two wheelers and pedestrian were involved in majority. *Conclusion:* To minimize the morbidity and mortality resulting from head injury consequent to road accidents there is a need for better maintenance of roads, strict enforcement of traffic rules, compulsory wearing of crash helmets by two wheelers along with imparting of compulsory road safety education to school children from primary education level.

Keywords: Aetiology, Head Injury, Medico legal, Autopsy.

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Head injury is defined by the national advisory neurological diseases and stroke council as “a morbid state resulting from gross or subtle structural changes in the scalp, skull and/or the contents of the skull, produced by the mechanical forces” [1]. Head is one of the most accessible and vital part of the body that is why frequently involved in unintentional (accidents) and intentional (physical assault) trauma. Even a single blow to head can be fatal, taking as an example of recent time where tragic death of sports person by unintentional blow to head has shocked the whole world. Head injuries are the most dreadful trauma in terms of morbidity and mortality. They are often caused by road traffic accidents, assaults, falls, industrial and domestic accidents, sports accidents etc. [2]. Nine

percent of global mortality is due to injuries, which causes about five million deaths every year worldwide [3]. Vehicular accidents accounts for nearly fifty percent of injuries and at least 40% of deaths results from head injury, a percentage that is increasing day by day [4]. Surge in motorization, expansion in the road network and the rising population in the country contribute toward the growing numbers of road accidents. During the decade 2001 to 2011, the road network in Bangladesh have increased at a compound annual growth rate (CAGR) of 3.4%, the numbers of registered motor vehicles in the country have amplified at a compound annual growth rate of 9.9% while the country's population have increased at CAGR of 1.6%. The number of road accidents in the country increased at a CAGR of 2.1% during the same period and the

fatalities due to road accident increased by 5.8% [5]. Earlier studies from the developing countries as well as from other terrains of Bangladesh showed various aetiologies of head injuries, which includes trauma, vehicular accidents, pedestrian injuries, fall from height etc. However, there is no substantiative government datas are available regarding the cause and manner of head injury. Therefore, the study was conducted to determine the various aetiological patterns of fatal head injuries in the region.

MATERIAL & METHODS

This prospective study was planned and conducted in The study was conducted in Dept. of forensic medicine, Rajshahi Medical College Hospital, Rajshahi, Bangladesh from June-2019 to July-2021. All the cases of fatal head injuries irrespective of their aetiology that were brought for medico legal autopsies were included in this study. All the cases were admitted and primarily treated for head injury. The data with regard to their age, sex, mode of injury in detail was collected from inquest papers/ police proceedings and first information report of the police (FIR). A semi-structured questionnaire served as study tool. All the

questionnaires were manually checked and edited for completeness and were then coded for computer entry. After compilation of collected data, analysis was done using Statistical Package for Social Sciences (SPSS), version 20 (IBM, Chicago, USA). The results were expressed using appropriate statistical methods. When the data follow a normal distribution then it was presented in mean \pm SD, otherwise median \pm IQR (Inter Quartile Range) was applied.

RESULTS

Of the 59 cases of head injuries, which were brought for medico legal autopsy, 90% were males and only 10% were females. The age ranges from 5 years to 80 years with mean 37.53 and standard deviation of 17.616, median was found to be 33.5 with standard error of mean 2.274 (Table 1). Roadside accident was the most common cause of head injury (65%), followed by the injury sustained due to assault (15%). Head injury as a result from fall from height seen only in two cases while railway accident was the genesis in 5 cases. Miscellaneous cause such as fall from stairs or fall of heavy object over the head comprises of 11% of total cases (Figure 1).




Fig-1: Mode of injury.

Table 1: Age and sex distribution of study subjects.

Age in years	Male	Female	Total
<10 years	02 (3.77%)	-	02 (3.39%)
11-20 years	07 (13.2%)	-	07 (11.86%)
21-30 years	14 (26.41%)	02 (33.33%)	16 (27.11 %)
31-40 years	11 (20.75%)	02 (33.33%)	13 (22.03 %)
41-50 years	08 (15.09%)	01 (16.67%)	09 (15.25%)
51-60 years	05 (9.43%)	-	05 (8.47%)
61-70 years	04 (7.55%)	-	04 (6.77%)
>71 years	02 (3.77%)	01 (16.67%)	03 (5.08%)
Total	53 (100%)	06 (100%)	59 (100%)

Two wheelers were involved in majority of roadside accidents (37.83%) of which 35.13% were the rider. 24.3% of cases were pedestrians and 29.72% were occupants of four wheeler. None of the two wheeler occupants were wearing helmets. Similary, no seal belt was in use by the driver or passenger of four wheeler at the time of accident.

DISCUSSION

Males being more involved in outdoor activities therefore exhibit preponderance over females in terms of incidence of head injury. 90% males were involved in head injuries as compared to 10% females. Goyal *et al.*, also observed more involvement of males (87.7%) in head injuries in their study [8]. Male to

female ratio was observed to be 4.2:1 by Kumar *et al.*, [7] Crandon *et al.*; noticed in their study that out of 857 patients admitted with head injury, 629 (73.5%) were males [8]. In developing countries, the productive age group of 15-44 years is commonly affected in road traffic injuries [9]. 21-30 year age group was commonly affected in our study (28.33%) with 50% of total cases were young adults that is age group of 21-40 years. Our results were consistent with the observations of Kumar *et al.*, which conclude that 51.7% cases of head injury were in 21-40 year age group [7]. Study of another developing country also indicates that age groups with the greatest incidence of head injury were aged 16-20 years, 21-25 years, and 25-30 years [10]. Crandon *et al.*, stated that in head injury median (IQR) age for the entire group was 28 years with a range of less than one year to 98 years [8]. Frequent involvement of younger adults in head injury and that too due to an accident may be because of their risk taking behavior with tendency towards high speeding and ignorance towards rules and regulations. Roadside accident was found to be major contributor in total no. of tolls from head injury (61.67%) followed by injury due to assault (15%). The rising numbers of motor vehicles and poor enforcement of traffic safety rules are major contributing factor in the alarming toll of fatalities from road traffic crashes in developing countries [9]. *The ascending trend in injuries and death due to road traffic accidents (RTA) has become an important health issue in Bangladesh* [11]. India witnessed one road accident every minute in year 2011, which claimed one life every 3.7 minutes, one of the highest in the world [12]. *According to latest data of National Crime Records Bureau (NCRB) of Bangladesh, the total number of deaths every year due to road accidents has now passed the 1,35,000 mark* [13]. Congestions on the roads, poor illumination during night, faulty speed breakers, and ignorance towards the safety precautions while driving such as use of helmets or safety belts are main reasons behind accentuating incidents of deaths due to road accidents. In comparison to developed countries where lane driving is followed and speed limit is strictly adhered to, such regulations are not made or being followed strictly in developing countries. Kumar *et al.*; studied that head injury with skull fracture was found to be associated with road traffic accident in 92.8% of their total subjects [7]. Crandon *et al.*; in their prospective study over a period of three years observed that head injury in admitted patients is mainly due to road traffic accidents, falls and interpersonal violence. Intentional injuries accounted for 26.7% of those admitted [8]. Yattoo and Tabish in their two phase study; both prospective and retrospective in regard to profile of head injuries and traumatic brain injury deaths; concluded that maximum no. of cases (44.4%) of traumatic head injury were due to RTA [14]. Bhateja *et al.*, in their retrospective study of 298 patients with head injury found RTA as commoner (53%), followed by fall from height (28.9%) and assault (10.4%) [15]. In our observations, rider two-wheeler was involved in

35.13% of cases and pedestrians in 24.3%. None of the occupants of two wheelers were wearing helmets during the incident. *In Bangladesh, two wheelers were the offending vehicle in the majority of the fatal crashes (39.9%)* [16]. The result of this study is in agreement with previous study from rural Haryana [17-19]. Lee *et al.*, in their study over a period of 10 years observed that the most common causes of head injury were motorcycle accidents (56.3%) and pedestrian injury (29.47%) [10]. Kumar *et al.*, in their study observed that, 51.3% cases of RTA were two wheeler occupants. The next major group of victims was the pedestrian .i.e. 37.2% [7]. Not all the times the occupants of motor vehicles are at fault. Pedestrians also do not follow the rules such as not using the zebra crossing or subways when available. This study has several strengths. To our knowledge, assessment of aetiological pattern of fatal head injury in medicolegal autopsies has not been much investigated in this region. Very few similar studies are available in the literature. Findings of this study may assist in planning preventive aspects of road traffic accidents which is a very important aspect now days. The study has some limitations as well. First, some may argue that findings of this study may not be generalised. I agree because such aetiological pattern tends to vary from place to place. Second, sample size of this study is small. Data of sixty subjects was analysed in this study. Studies with bigger sample size are warranted.

CONCLUSION

Maximum number of fatalities due to head injury was subsequent to roadside accident and that too in a productive age group of 21-40 years. Rider two wheelers and pedestrian were involved in majority. To minimize the morbidity and mortality resulting from head injury consequent to road accidents there is a need for better maintenance of roads, strict enforcement of traffic rules, compulsory wearing of crash helmets by two wheelers and shoulder belt in cars along with imparting of compulsory road safety education right from primary education level. Injury surveillance and trauma registry should be done at every hospital so that preventive strategies can be formulated.

REFERENCES

1. Viji, K. (2011). Textbook of forensic medicine and toxicology. 5th ed. Kundli: Elsevier; 270-86.
2. Dhillon, S., Kapila, P., & Sekhon, H. S. (2007). Pattern of injuries present in road traffic accident in shimla hills. *Journal of Punjab Academy of Forensic Medicine and Toxicology*, 7(2), 7-10.
3. Gururaj, G. (2005). Injuries in India: A national perspective. Background Papers: Burden of Disease in India Equitable Development-Healthy Future. New Delhi: National Commission on Macroeconomics and Health, Ministry of Health & Family Welfare, Government of India, 325-347.
4. Kirkwood, R. (1995). Essentials of neuroimaging. 2nd ed. New York: Churchill Livingstone; 339-75.

5. Ruikar, M. (2013). National statistics of road traffic accidents in India. *Journal of Orthopedics, Traumatology and Rehabilitation*, 6(1), 1.
6. Goyal, M. K., Verma, R., Kochar, S. R., & Asawa, S. S. (2010). Correlation of CT scan with postmortem findings of acute head trauma cases at SMS Hospital.
7. Kumar, D., Bains, V., Sharma, B. R., & Harish, D. (2012). Descriptive study of head injury and its associated factors at tertiary hospital, Northern India. *J Community Med Health Educ*, 2(04), 10-4172.
8. Crandon, I. W., Harding-Goldson, H. E., McDonald, A., Fearon-Boothe, D., & Meeks-Aitken, N. (2007). The aetiology of head injury in admitted patients in Jamaica. *West Indian medical journal*, 56(3), 223-225.
9. Nantulya, V. M., & Reich, M. R. (2002). The neglected epidemic: road traffic injuries in developing countries. *Bmj*, 324(7346), 1139-1141.
10. Lee, S. T., Lui, T. N., Chang, C. N., Wang, D. J., Heimburger, R. F., & Fai, H. D. (1990). Features of head injury in a developing country--Taiwan (1977-1987). *The Journal of trauma*, 30(2), 194-199.
11. Gopalakrishnan, S. (2012). A public health perspective of road traffic accidents. *Journal of family medicine and primary care*, 1(2), 144.
12. Gupta, M. D. (2012). Every 3 minutes, an Indian dies in a road accident. *Hindustan Times. New Delhi*.
13. India has the highest number of road accidents in the world | Asia | DW.DE | 29.04.2010. [Cited 2014 Dec 8]. Available from: <http://www.dw.de/india-has-the-highest-number-of-road-accidents-in-the-world/a-5519345-1>
14. Yattoo, G. H., & Tabish, A. (2008). The profile of head injuries and traumatic brain injury deaths in Kashmir. *Journal of trauma management & outcomes*, 2(1), 1-9.
15. Bhateja, A., Shukla, D., Devi, B. I., & Kolluri, V. S. (2009). Coup and contrecoup head injuries: Predictors of outcome. *Indian Journal of Neurotrauma*, 6(02), 115-118.
16. Pruthi, N., Ashok, M., Shiva, K. V., Jhavar, K., Sampath, S., & Devi, B. I. (2012). Magnitude of pedestrian head injuries & fatalities in Bangalore, south India: A retrospective study from an apex neurotrauma center. *The Indian journal of medical research*, 136(6), 1039.
17. Kaur, B., Singh, A., Aggarwal, O. P., & Singh, H. (2012). Pattern of injuries due to fatal road traffic accidents in rural Haryana: an epidemiological survey. *Journal of Indian Academy of Forensic Medicine*, 34(3), 229-232.
18. Chaudhary, C., Singh, A., Pathak, R., Ahluwalia, S. K., Goel, R. K. D., & Mithra, P. (2013). Predictors of seatbelt and helmet usage among victims seeking care at emergency department in a tertiary care hospital in rural Northern India. *Nepal Journal of Medical Sciences*, 2(1), 57-61.
19. Singh, A., Bhardwaj, A., Pathak, R., & Ahluwalia, S. K. (2011). An epidemiological study of road traffic accident cases at a tertiary care hospital in rural Haryana. *Indian Journal of Community Health*, 23(2), 53-55.

Cite This Article: Md. Jasim Uddin (2021). Determination the Aetiological Pattern of Head Injury in the Region. *East African Scholars J Med Surg*, 3(11), 206-209.