

Role of Phytoremediation in Removing Air Pollutants: A Review

Hasnain Raza^{1*}, Tehmina Bibi¹, Haseena Bibi¹, Muhammad Junaid Asim¹, Huda Bilal², Awais Rasheed¹, Muhammad Bilal Shoukat¹, Attiq Ur Rehman¹

¹Department of Soil and Environmental Sciences, MNS-University of Agriculture, Multan, Pakistan

²Institute of Plant Protection, MNS-University of Agriculture, Multan, Pakistan

*Corresponding author: Hasnain Raza

| Received: 16.08.2021 | Accepted: 25.09.2021 | Published: 01.10.2021 |

Abstract: Air pollution has become a global issue in recent years due to increasing health and socioeconomic risks. It has negative effects on human health in both indoor and outdoor environments. There is a growing severity and impact of these threats, especially in developing countries such as Pakistan that have no adequate systems of alert, management, and protection. The main problem facing the scientific community now is to reduce air pollutant emissions properly. Phytoremediation seems a promising prospect: an environmentally sustainable, low-cost, plant-based approach to maintenance, soil stabilization, and aesthetical pleasure. The present review discusses Pakistan's indigenous plants, which have the potential to mitigate specific air pollutants.

Keywords: Air Pollutants, Phytoremediation, Native Plants, Human Health, Emissions.

INTRODUCTION

The accumulation of toxic or poisonous substances in the earth's atmosphere, which has negative consequences for human health and the ecosystem, is referred to as air pollution. The planet is on the verge of a global climate crisis, with air pollution being the primary cause. It is a critical global problem, the most serious environmental threat to human health, and the cause of 4.2 million deaths every year (WHO, 2021). Due to a lack of pollution controls and air quality regulations, Pakistan has some of the world's most polluted cities (Colbeck *et al.*, 2009). Particulate matter (PMs), Nitrogen oxide (NO₂), Sulphur dioxide (SO₂) and Ground-level ozone (O₃) are all major air pollutants. The first step in reducing air pollution is to eradicate or minimize anthropogenic-caused emissions (Ahmad and Aziz, 2013). Due to the serious consequences of air pollution, Pakistan's lack of progress in implementing various technological steps to avoid pollution is cause for concern. The second step is

to clean up any toxins that have already been released into the environment.

Various air pollution reduction techniques, policies, and models have been proposed (Macpherson *et al.*, 2017). Biological remediation, also known as bioremediation, can be used to reduce air pollution. It is the process of species assimilating, degrading, or transforming harmful substances into less harmful or non-toxic forms. Phytoremediation is the process of using plants to remove toxins from the air, soils, and water (Raza *et al.*, 2021). The key benefits of phytoremediation technology are that it is an aesthetically appealing and solar energy-driven cleanup technology and that it can treat a range of environmental pollutants at the same time. It is a cost-effective technique since the cost of phytoremediation is 60-80% less than that of traditional physio-chemical or mechanical systems (Singh and Verma, 2007).

Quick Response Code



Journal homepage:

<http://crosscurrentpublisher.com>

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

Citation: Hasnain Raza *et al* (2021). Role of Phytoremediation in Removing Air Pollutants: A Review. *Cross Current Int J Agri Vet Sci*, 3(6), 54-59.

S.r.	Pollutant	Sources	Diseases	References
1	Particulate matters (PMs)	Industrial activities, Forest fires, Volcano eruptions, Agricultural activities, Resuspension of soil, Combustion of fossil fuel, Vehicle emissions, and energy production.	Diseases in nervous system and respiratory system. Heart, bladder and lung cancer and Cardiovascular morbidity.	(Kelly and Fussell, 2015)
2	Nitrous oxide (NO ₂)	Industrial manufacturing, Vehicular traffic, Fossil fuel combustion, Industrial activities, and Domestic heating.	Respiratory disease, Lung illness, Asthma, and Breathing disorder.	(Peel <i>et al.</i> , 2013)
3	Sulfur dioxides (SO ₂)	Burning of fossil fuels, oil, coal and diesel. Sources that contain sulfur from burning of material are power plants, metals processing and smelting facilities, and vehicles.	Asthma, Chronic bronchitis, Lung diseases, and Emphysema	(Manisalidis <i>et al.</i> , 2020)
4	Ozone (O ₃)	This happens when pollutants emitted by cars, power plants, industrial boilers, refineries, chemical plants, and other sources chemically react in the presence of sunlight. Ozone is most likely to reach unhealthy levels on hot sunny days in urban environments, but can still reach high levels during colder months.	Asthma, emphysema, and chronic bronchitis and climate change	(Zhang <i>et al.</i> , 2019)
5	Heavy metals (HMs)	Industrial activities, Agricultural activities, and Combustion of fossil fuel.	Carcinogens, lung, liver, kidney and brain damage, skin irritation, reduced fetal growth, damage to blood vessels, cancer, and death.	(Lentini <i>et al.</i> , 2017)

Phytoremediation is a process in which plants absorb contaminants from the air and then degrade or detoxify them via a variety of mechanisms. This has been shown to be a successful plant-based, environmentally friendly, and long-term method of reducing air pollutants in both indoor and outdoor settings (Weyens *et al.*, 2015). Phytoremediation stands out among existing technologies because of its self-maintaining, cost-effective procedures, as well as increased ethical and societal acceptance (Doty *et al.*, 2007). Numerous studies have been conducted on plants' ability to absorb certain toxins under various environmental circumstances. Many studies further explore their broader applications, such as the use of

taller plants to remove airborne pollutants, which is particularly useful in outdoor settings.

Air quality in Pakistan and Phytoremediation

Air pollution is a major environmental challenge in Pakistan's cities and costs the national checker trillions of rupees per year (IQAir, 2021). Vehicle emissions, fossil fuel combustion of unleaded petrol, and power plants are the main sources of fine particulate pollution (Asad *et al.*, 2011). Poor air quality in Pakistan is creating problems including the development of aerosols, asthma, and lead to toxicity, and accumulation of greenhouse gas. To save 222 million lives, immediate action is needed.

Title: Top ten countries with poor Air Quality index (AQI) in 2020 by (IQAir, 2021).

S.r.	Country	Population	AVG. US AQI
1	Bangladesh	164'689'383	162
2	Pakistan	220'892'331	153
3	India	1'380'004'385	141
4	Mongolia	3'278'292	128
5	Afghanistan	38'928'341	128
6	Oman	5'106'622	123
7	Qatar	2'881'060	123
8	Kyrgyzstan	6'524'191	121
9	Indonesia	273'523'621	114
10	Bosnia Herzegovina	3'280'815	113

Particulate Matter (PM)

Particulate Matter (PM) is much more visible and intense than the other pollutants mentioned above, but it is less toxic, as it serves as a nucleus for the

deposition of several hazardous chemicals found in any environment. Soot, ash, and chemical byproducts created by combustion or chemical mixing, road construction, and farming are the most common sources

of PM. Pakistan as a whole was polluted with PM_{2.5} in 2018, and the average yearly level in 2019 was 65.81 µg/m³ (IQAir, 2021). The highest average PM_{2.5} mass levels in Karachi were estimated to be 668 µg m⁻³ amongst the 18 biggest cities of the world (Gurjar *et al.*, 2008). Biswas *et al.* (Biswas *et al.*, 2008) found that the average PM_{2.5} mass concentration in Lahore is many times higher than in Hong Kong, Seoul, and New York City. With about 105 thousand deaths per year, Pakistan is one of the most premature death rates in the world due to elevated PM_{2.5} levels (Giannadaki *et al.*, 2016).

The quantity of PM contaminants that plants extract from the air is usually caused by deposition. The quantity of contaminants deposited is dependent on the index of leaf area, deposition, PM concentration, and period of vegetation. The vegetation area of each soil unit surface is used to calculate the index of the leaf area, which may vary considerably between plant species (Janhäll, 2015). Some native plants in Pakistan can reduce PM levels in the air like Hedge or Field maple (*Acer campestre*), *Coniferous corsican*, French hales *Sorbus latifolia*, *Alnus spp.*, sycamore (*Platanus occidentalis*), Poplar (*Populus spp.*), Southern blue gum (*Eucalyptus globulus*), Oak (*Quercus spp.*), Red fir (*Pseudotsuga menziesii*), Woodland Elaecarpus (*E. sylvestris*), English yew (*Taxus baccata*), Scots pine (*Pinus sylvestris*), Laceshrub (*Stephanandra incisa*), Mugo pine (*Pinus mugo*), Anglojap yew (*Taxus media*), Common silver birch (*Betula pendula*), Neem tree (*Azadirachta indica*), Mango (*Mangifera indica*), Banyan (*Ficus bengalensis*), and Hibiscus (*Hibiscus rosasinensis*).

Several studies have looked into the ability of these shrubs and trees to minimize urban PM air pollution (Freer-Smith *et al.*, 2004; Manes *et al.*, 2016; Sæbø *et al.*, 2012; Terzaghi *et al.*, 2013). Freer-Smith *et al.*, 2004, studied the capture efficiency of PM on *Alnus spp.*, *Quercus spp.*, *E. globulus*, and *P. menziesii* using NaCl droplets of 1 mm and wind tunnel. At a 9 m/s wind speed, *P. menziesii* and *Quercus spp* seemed to have the best overall efficiency of 0.671 % and 0.348 %, respectively. Planting trees such as *A. indica*, *F. bengalensis*, *H. rosasinensis* and *M. indica* along urban roadsides. According to Mate and Deshmukh (Mate and Deshmukh, 2015), plants can be a good way to control particulates released by vehicles. The plants that captured the most PM were *P. mugo*, *T. baccata*, *P. sylvestris*, *S. incisa*, *T. media*, and *B. pendula* (Sæbø *et al.*, 2012). Trees such as cedars and oaks planted 25 meters along roadside have been shown to reduce PM_{2.5} and PM₁₀ concentrations by 50%, though tall prairie grass lowers them by 35% (Cowherd *et al.*, 2006).

Sulphur dioxide (SO₂)

Sulphur dioxides were the first air pollutants to cause harm to human health and wildlife. SO₂ has been significantly increased in the air by fossil fuels combustion (Zhang *et al.*, 2013). The expected growth

in economic activity in Pakistan to increase SO₂ emissions by 8.7 between 2005-2030 [4]. The SO₂ (52.5 ppb) was found at Lahore, higher than Karachi and Peshawar (Ghauri *et al.*, 2007). During the study period the amount of SO₂ was below both the National Environmental Quality Standards (NEQS) and United States Environmental Protection Agency (USEPA) standards (Ashraf *et al.*, 2013). SO₂ in Islamabad is within safe limits (Rasheed *et al.*, 2014; Shahid *et al.*, 2019). The average daily SO₂ in Faisalabad was within NEQS limits (Asghar *et al.*, 2018). The plant leaves absorb SO₂ through stomata and are subsequently hydrated and oxidized to sulfite and sulfate, when accumulated to high concentrations, may inhibit photosynthesis and energy metabolism (Wei *et al.*, 2017).

Ground-level Ozone (O₃)

The economic yield of major agricultural crops is impacted by O₃ pollution. Many studies showed that O₃ pollution has a major impact on agricultural productivity, but the health effects of O₃ tend to be less significant in Pakistan than those of other pollutants. The permissible amount of ozone (O₃) in ambient air is 130 g m⁻³ under NEQS (Khawaja and Shams, 2020). For an 8-hour daily average, the World Health Organization has developed a guideline value of 100 g m⁻³ for O₃ levels. According to (Colbeck *et al.*, 2010), O₃ concentrations in Pakistan's major cities were well within WHO air quality guidelines.

Within the leaf structure of the plant, O₃ can be completely detoxified. After entering the stomata, O₃ can be extracted and subsequently reacted within the intercellular zone. O₃ build up in the intercellular space at high O₃ levels and reduce the total O₃ flux (Fares *et al.*, 2010). Plants have been shown to remove O₃ from the atmosphere on an annual basis implying that plants' metabolic pathways will permanently eliminate some O₃ (Mikkelsen *et al.*, 2004; Nowak *et al.*, 2006). In Pakistan, there are some native plants which have the potential to remediate the O₃ level in the air. These are *Larix decidua*, *Picea smithiana*, deciduous conifer, deciduous, and evergreen broadleaved and conifer forests.

So many studies have looked into the effects of various trees on O₃ elimination (Alonso *et al.*, 2011; Manes *et al.*, 2016). The O₃ uptake of the *L. decidua*, the Cembran pine, a deciduous conifer, and the Norway spruce, 1.40, 1.18, and 1.09 nmol/m²s (Wieser *et al.*, 2003). Alonso *et al.*, (2011) looked at how various types of vegetation affected O₃ levels. For evergreen broadleaved, deciduous and conifer forests, 6.64, 6.86, and 3.98 mg/m² were found to be annual absorbed cumulative O₃ fluxes, respectively. Overall, it's important to remember that air pollutant removal effectiveness varies by plant species.

Nitrogen dioxide (NO₂)

Nitrogen dioxide (NO₂), Nitrogen trioxide (N₂O₃), Nitric oxide (NO), and Nitrous oxide (N₂O) are some of the nitrogen (N) oxides found in the atmosphere. Since NO₂ is the most common form of NO_x formed by humans, the USEPA only regulates it. The country's current levels of NO₂ to some extent higher than the WHO air quality permissible value of 40 g/m³, according to annual NO₂ concentrations derived from 48-hour data, with the highest concentrations of 49 ± 28 µg/m³ in Islamabad, 52 ± 21 µg/m³ in Peshawar, 46 ± 15 µg/m³ in Karachi and 49 ± 25 µg/m³ in Lahore (IQAir, 2021). The ability of plants

to absorb some NO₂ through their stomata while also assimilating and metabolizing gaseous nitrogen contaminants appears to allow for varying levels of permanent NO₂ elimination. NO₂ fluxes differ, much like those of other gaseous compounds (Wei *et al.*, 2017). Plants have been shown to minimize regional NO₂ levels in particular (Nowak *et al.*, 2014). The most productive woody plants in Pakistan are *Robinia pseudoacacia*, *Populus nigra*, *Eucalyptus viminalis*, and *Magnolia kobu*, and the most herbaceous plants include *Crassocephalum crepidioides*, *Nicotiana tabacum* and *Erechtites hieracifolia*.

S.r.	Plants	Pollutants	References
1.	<i>Phytolacca Americana</i>	HMs	(Pandey and Bajpai, 2019)
2.	<i>Pinus mugo</i>	PM	(Sæbø <i>et al.</i> , 2012).
3.	<i>Salix schwerinii</i>	HMs	(Mohsin <i>et al.</i> , 2019)
4.	<i>Blumea malcolmii</i>	POP	(Kagalkar <i>et al.</i> , 2011)
5.	<i>Taxus baccata</i>	PMs	(Sæbø <i>et al.</i> , 2012).
6.	<i>Sorghum x drummondii</i>	VOCs	(Dominguez <i>et al.</i> , 2020)
7.	<i>Helianthus annuus</i>	POPs,	(Lee and Yang, 2010)
8.	<i>Picea smithiana</i>	O ₃	(Alonso <i>et al.</i> , 2011)
9.	<i>Mangifera indica</i>	PM	(Mate and Deshmukh, 2015),
10.	<i>Sorghum x drummondii</i>	VOCs	(Dominguez <i>et al.</i> , 2020)
11.	<i>Arundo donax</i>	VOCs,	(Guarino <i>et al.</i> , 2020)
12.	<i>Typha angustifolia</i>	IAP	(Li <i>et al.</i> , 2016)
13.	<i>Phaseolus vulgaris</i>	HMs	(Lee and Yang, 2010)
13.	<i>Juncus effuses</i>	HM	(Najeeb <i>et al.</i> , 2017)
15.	<i>Spirodela polyrhiza</i>	POP	(Kristanti <i>et al.</i> , 2012)
16.	<i>Pinus sylvestris</i> ,	PM	(Sæbø <i>et al.</i> , 2012).
17.	<i>Sorghum x drummondii</i>	HMs	(Dominguez <i>et al.</i> , 2020)
18.	<i>Larix decidua</i>	O ₃	(Wieser <i>et al.</i> , 2003)
19.	<i>Helianthus annuus</i>	PMs	(Lee and Yang, 2010)
20.	<i>Ficus bengalensis</i> ,	PM	(Mate and Deshmukh, 2015),
21.	<i>Carpobrotus aequilaterus</i>	PM	(Terzaghi <i>et al.</i> , 2013)
22.	<i>Juncus effuses</i>	PMs	(Najeeb <i>et al.</i> , 2017)
23.	<i>Stephanandra incisa</i>	PM	(Sæbø <i>et al.</i> , 2012).
24.	<i>Azadirachta indica</i>	PM	(Mate and Deshmukh, 2015),
25.	<i>Sorghum x drummondii</i>	POPS	(Dominguez <i>et al.</i> , 2020)

CONCLUSION

Air pollution in Pakistan has a negative impact on human health and agriculture. Air quality is worsening at enormous pace and the government and many other organizations have identified it as a serious issue. In this respect, however, little work has been done. Due to the complexity of air pollution existence and origins, it is difficult to develop adequate control methods. The plant is therefore an asset to enhance air quality either by metabolism, sequestration or degradation of particular air pollutants. In some plants, toxic pollutants can be assimilated, degraded or modified in air into less toxic pollutants which allow airborne pollutants to be removed using the AP technology. There are several plants in Pakistan that can clean outdoor and indoor air. In roadsides, parks and manmade forests, plants and trees above mentioned should be grown. The phytoremediation of air pollution

is therefore still a developing phenomenon on a commercial scale. The scientists and the general public are well aware of several benefits from tree planting and growing, but there are uncertainty about the capacity and adequacy of individual species for particular pollutants.

REFERENCES

- Ahmad, S.S., Aziz, N. (2013). Spatial and temporal analysis of ground level ozone and nitrogen dioxide concentration across the twin cities of Pakistan. *Environmental monitoring and assessment*, 185(4); 3133–3147.
- Alonso, R., Vivanco, M.G., González-Fernández, I., Bermejo, V., Palomino, I., Garrido, J.L., Elvira, S., Salvador, P., Artíñano, B. (2011). Modelling the influence of peri-urban trees in the air quality of

- Madrid region (Spain). *Environmental pollution*, 159(8–9):2138–2147.
- Asad, F., Haider, K., Shaheen, A. (2018). *Trend Analysis of Ambient Air Quality of Islamabad*. Asghar, Z., Ali, W., Nasir, A., Arshad, A. Atmospheric monitoring for ambient air quality parameters and source apportionment of city Faisalabad, Pakistan. *Earth Sciences Pakistan (ESP)*, 2(1); 1–4.
 - Ashraf, N., Mushtaq, M., Sultana, B., Iqbal, M., Ullah, I., Shahid, S.A. (2013). Preliminary monitoring of tropospheric air quality of Lahore City in Pakistan. *Sustainable Development*, 3(1); 19–28.
 - Biswas, K.F., Ghauri, B.M., Husain, L. (2008). Gaseous and aerosol pollutants during fog and clear episodes in South Asian urban atmosphere. *Atmospheric Environment*, 42(33); 7775–7785.
 - Colbeck, I., Nasir, Z., Ali, Z. (2009). The state of ambient air quality in Pakistan—a review. *Environmental science and pollution research international*, 17; 49–63.
 - Colbeck, I., Nasir, Z.A., Ali, Z. (2010). The state of indoor air quality in Pakistan—a review. *Environmental Science and Pollution Research*, 17(6); 1187–1196.
 - Cowherd, C., Muleski, G., & Gebhart, D. (2006, May). Development of an emission reduction term for near-source depletion. In *Proceedings of 15th International Emission Inventory Conference: “Reinventing Inventories—New Ideas in New Orleans”*, New Orleans, LA.
 - Dominguez, J. J. A., Inoue, C., & Chien, M. F. (2020). Hydroponic approach to assess rhizodegradation by sudangrass (*Sorghum x drummondii*) reveals pH-and plant age-dependent variability in bacterial degradation of polycyclic aromatic hydrocarbons (PAHs). *Journal of hazardous materials*, 387, 121695.
 - Doty, S. L., James, C. A., Moore, A. L., Vajzovic, A., Singleton, G. L., Ma, C., ... & Strand, S. E. (2007). Enhanced phytoremediation of volatile environmental pollutants with transgenic trees. *Proceedings of the National Academy of Sciences*, 104(43), 16816-16821.
 - Fares, S., Goldstein, A., & Loreto, F. (2010). Determinants of ozone fluxes and metrics for ozone risk assessment in plants. *Journal of Experimental Botany*, 61(3), 629-633.
 - Freer-Smith, P. H., El-Khatib, A. A., & Taylor, G. (2004). Capture of particulate pollution by trees: a comparison of species typical of semi-arid areas (*Ficus nitida* and *Eucalyptus globulus*) with European and North American species. *Water, Air, and Soil Pollution*, 155(1), 173-187.
 - Ghauri, B., Lodhi, A., & Mansha, M. (2007). Development of baseline (air quality) data in Pakistan. *Environmental Monitoring and Assessment*, 127(1), 237-252.
 - Giannadaki, D., Lelieveld, J., & Pozzer, A. (2016). Implementing the US air quality standard for PM 2.5 worldwide can prevent millions of premature deaths per year. *Environmental Health*, 15(1), 1-11.
 - Guarino, F., Miranda, A., Castiglione, S., & Cicatelli, A. (2020). Arsenic phytovolatilization and epigenetic modifications in *Arundo donax* L. assisted by a PGPR consortium. *Chemosphere*, 251, 126310.
 - Gurjar, B. R., Butler, T. M., Lawrence, M. G., & Lelieveld, J. (2008). Evaluation of emissions and air quality in megacities. *Atmospheric Environment*, 42(7), 1593-1606.
 - IQAir Air quality in Pakistan, 2021; 1–9.
 - Janhäll, S. (2015). Review on urban vegetation and particle air pollution—Deposition and dispersion. *Atmospheric environment*, 105, 130-137.
 - Kagalkar, A. N., Jadhav, M. U., Bapat, V. A., & Govindwar, S. P. (2011). Phytodegradation of the triphenylmethane dye Malachite Green mediated by cell suspension cultures of *Blumea malcolmii* Hook. *Bioresource technology*, 102(22), 10312-10318.
 - Kelly, F. J., & Fussell, J. C. (2015). Air pollution and public health: emerging hazards and improved understanding of risk. *Environmental geochemistry and health*, 37(4), 631-649.
 - Khwaja, M. A., & Shams, T. (2020). Pakistan National Ambient Air Quality Standards: A comparative Assessment with Selected Asian Countries and World Health Organization (WHO).
 - Kristanti, R. A., Kanbe, M., Hadibarata, T., Toyama, T., Tanaka, Y., & Mori, K. (2012). Isolation and characterization of 3-nitrophenol-degrading bacteria associated with rhizosphere of *Spirodela polyrrhiza*. *Environmental Science and Pollution Research*, 19(5), 1852-1858.
 - Lee, M., & Yang, M. (2010). Rhizofiltration using sunflower (*Helianthus annuus* L.) and bean (*Phaseolus vulgaris* L. var. *vulgaris*) to remediate uranium contaminated groundwater. *Journal of hazardous materials*, 173(1-3), 589-596.
 - Li, Y., Zhang, J., Zhu, G., Liu, Y., Wu, B., Ng, W. J., ... & Tan, S. K. (2016). Phytoextraction, phytotransformation and rhizodegradation of ibuprofen associated with *Typha angustifolia* in a horizontal subsurface flow constructed wetland. *Water research*, 102, 294-304.
 - Macpherson, A. J., Simon, H., Langdon, R., & Misenheimer, D. (2017). A mixed integer programming model for National Ambient Air Quality Standards (NAAQS) attainment strategy analysis. *Environmental Modelling & Software*, 91, 13-27.
 - Manes, F., Marando, F., Capotorti, G., Blasi, C., Salvatori, E., Fusaro, L., ... & Munafò, M. (2016). Regulating ecosystem services of forests in ten Italian metropolitan cities: air quality improvement

- by PM10 and O3 removal. *Ecological indicators*, 67, 425-440.
- Manisalidis, I., Stavropoulou, E., Stavropoulos, A., & Bezirtzoglou, E. (2020). Environmental and health impacts of air pollution: a review. *Frontiers in public health*, 8, 14.
 - Mate, A. R., & Deshmukh, R. R. (2015). To control effects of air pollution using roadside trees. *International Journal of Innovative Research in Science, Engineering and Technology*, 4(11), 11167-11172.
 - Mikkelsen, T. N., Ro-Poulsen, H., Hovmand, M. F., Jensen, N. O., Pilegaard, K., & Egeløv, A. H. (2004). Five-year measurements of ozone fluxes to a Danish Norway spruce canopy. *Atmospheric Environment*, 38(15), 2361-2371.
 - Mohsin, M., Kuittinen, S., Salam, M. M. A., Peräniemi, S., Laine, S., Pulkkinen, P., ... & Pappinen, A. (2019). Chelate-assisted phytoextraction: Growth and ecophysiological responses by *Salix schwerinii* EL Wolf grown in artificially polluted soils. *Journal of Geochemical Exploration*, 205, 106335.
 - Najeeb, U., Ahmad, W., Zia, M. H., Zaffar, M., & Zhou, W. (2017). Enhancing the lead phytostabilization in wetland plant *Juncus effusus* L. through somaclonal manipulation and EDTA enrichment. *Arabian Journal of Chemistry*, 10, S3310-S3317.
 - Nowak, D. J., Crane, D. E., & Stevens, J. C. (2006). Air pollution removal by urban trees and shrubs in the United States. *Urban forestry & urban greening*, 4(3-4), 115-123.
 - Nowak, D. J., Hirabayashi, S., Bodine, A., & Greenfield, E. (2014). Tree and forest effects on air quality and human health in the United States. *Environmental pollution*, 193, 119-129.
 - Pandey, V. C., & Bajpai, O. (2019). Phytoremediation: from theory toward practice. In *Phytomanagement of polluted sites* (pp. 1-49). Elsevier.
 - Peel, J. L., Haeuber, R., Garcia, V., Russell, A. G., & Neas, L. (2013). Impact of nitrogen and climate change interactions on ambient air pollution and human health. *Biogeochemistry*, 114(1), 121-134.
 - Rasheed, A., Aneja, V. P., Ayyer, A., & Rafique, U. (2014). Measurements and analysis of air quality in Islamabad, Pakistan. *Earth's future*, 2(6), 303-314.
 - Raza, H., Qurat-ul-Ain, Asim, M.J., Rehman, A.U., Rasheed, A., Bilal, H., Maqsood, M., Raza, A., Shoukat, M.B. (2021). Bio Remedial Potential for the Treatment of Contaminated Soils. *Current Research in Agriculture and Farming*, 2; 53–58.
 - Sæbø, A., Popek, R., Nawrot, B., Hanslin, H.M., Gawronska, H., Gawronski, S.W. (2012). Plant species differences in particulate matter accumulation on leaf surfaces. *Science of the Total Environment*, 427; 347–354.
 - Shahid, I., Chishtie, F., Bulbul, G., Shahid, M.Z., Shafique, S., Lodhi, A. (2019). State of air quality in twin cities of Pakistan: Islamabad and Rawalpindi. *Atmósfera*, 32(1); 71–84.
 - Singh, S. N., & Verma, A. (2007). Phytoremediation of air pollutants: a review. *Environmental bioremediation technologies*, 293-314.
 - Terzaghi, E., Wild, E., Zacchello, G., Cerabolini, B. E., Jones, K. C., & Di Guardo, A. (2013). Forest filter effect: role of leaves in capturing/releasing air particulate matter and its associated PAHs. *Atmospheric Environment*, 74, 378-384.
 - Wei, X., Lyu, S., Yu, Y., Wang, Z., Liu, H., Pan, D., & Chen, J. (2017). Phylloremediation of air pollutants: exploiting the potential of plant leaves and leaf-associated microbes. *Frontiers in plant science*, 8, 1318.
 - Wei, X., Lyu, S., Yu, Y., Wang, Z., Liu, H., Pan, D., & Chen, J. (2017). Phylloremediation of air pollutants: exploiting the potential of plant leaves and leaf-associated microbes. *Frontiers in plant science*, 8, 1318.
 - WHO Air pollution. 2021; 1–9.
 - Wieser, G., Matyssek, R., Köstner, B., & Oberhuber, W. (2003). Quantifying ozone uptake at the canopy level of spruce, pine and larch trees at the alpine timberline: an approach based on sap flow measurement. *Environmental Pollution*, 126(1), 5-8.
 - Zhang, J. J., Wei, Y., & Fang, Z. (2019). Ozone pollution: a major health hazard worldwide. *Frontiers in immunology*, 10, 2518.
 - Zhang, X., Zhou, P., Zhang, W., Zhang, W., & Wang, Y. (2013). Selection of landscape tree species of tolerant to sulfur dioxide pollution in subtropical China. *Open Journal of Forestry*, 3(04), 104.