

Effects of Black Pepper on Oxidative Stress Markers in Acrylamide-Induced Toxicity in Male Wistar Rats

Onyebuchi Obia^{1*}, Edna Vincent Agorom¹, Edith Reuben¹, Augusta Chiamaka Innocent¹

¹Department of Human Physiology, Faculty of Basic Medical Sciences, College of Medical Sciences, Rivers State University, Nkpolu-Oroworukwo, Port Harcourt, Nigeria

*Corresponding author: Onyebuchi Obia

| Received: 18.03.2025 | Accepted: 24.04.2025 | Published: 26.04.2025 |

Abstract: Background to the study: The common methods of processing carbohydrate based foods require exposure to high temperatures which often results in the formation of a toxicant called acrylamide. The aim of the present study is to determine the effect of black pepper on the oxidative stress markers of wistar rats exposed to acrylamide. **Methodology:** The study involved 20 male wistar rats separated into 4 groups of 5 rats each as follows; Group 1 served as control. Animals in groups 2, 3 and 4 respectively received a daily oral dose of 30mg/kg of Acrylamide (ACR) throughout the experiment. Group 2 remained untreated (ACR only). Animals in groups 3 and 4 received in addition, 50mg/kg/day of black pepper and 150mg/kg/day of vitamin E respectively. All treatments were given orally for 14 days and thereafter animals were sacrificed and blood samples collected to determine the concentrations of oxidative stress markers in the blood using standard methods. **Results:** Our results showed that daily oral administration of ACR caused significant reduction in the plasma concentrations of GSH, GPx and catalase but no significant changes in the SOD and MDA levels. However, concomitant administration of black pepper mitigated the ACR induced depletion of GPx and catalase in a similar fashion with vitamin E but caused no significant changes in the GSH, SOD and MDA. **Conclusion:** Conclusively, black pepper exerts its antioxidant effects on ACR induced toxicity by improving GPx and catalase levels but not GSH, SOD and MDA. Aside from GSH levels, the antioxidant potentials of black pepper in our study showed a similar pattern as vitamin E.

Keywords: Black pepper, Oxidative stress, acrylamide-induced toxicity, wistar rats.

INTRODUCTION

The palatability of most foods consumed in most modern restaurants depends largely on the method of processing including; frying, roasting and baking. These methods mainly involve exposure of the processed food to higher thermal heating (Lingnert *et al.*, 2002; Pan *et al.*, 2020; Rifai & Saleh, 2020; Timmermann *et al.*, 2021; Alefe, 2024) with resultant formation of a biochemical compound known as acrylamide (Zhang *et al.*, 2021; Yan *et al.*, 2023; Adimas *et al.*, 2024). The incidence of acrylamide in foods processed by higher temperatures was first noted by the food administration agency in Sweden (SNFA) in 2002 (Ferrer-Aguirre *et al.*, 2016). Several studies have linked the consumption of acrylamide contaminated food with toxicity to vital organs (Zhang *et al.*, 2021; Zheng & Xiao, 2022; Govindaraju *et al.*, 2024). Acrylamide-induced toxicity is mediated by cellular oxidative stress (Sharafi-Rad *et al.*, 2020; Banc *et al.*, 2022). The optimal health of an organism requires maintenance of a fair balance between

oxidants and antioxidants. Antioxidants play a pivotal role in disease prevention by preventing the formation of free radicals and by implication reduces cellular damage.

Most of the thermally processed foods are usually garnished with pepper such that in any popular Nigerian restaurant, foods like peppered beef, fish, roasted chicken and gizzard, roasted yam, roasted plantain and potatoes abound. This is to buttress the fact that pepper is almost always found in these thermally processed foods especially in a typical Nigerian dish. Black pepper is the commonest variety of pepper spice used in these fried, roasted or baked foods. Some of the nutritional benefits of pepper has been studied including; lipid lowering potential, organo-protective, enhanced intestinal motility and antioxidant effects (Hervert-Hernández *et al.*, 2010; Ludy *et al.*, 2012; Charles *et al.*, 2024; Emmanuel *et al.*, 2025; Obia & Emmanuel, 2025). The present study will determine the impact of black pepper on acrylamide-induced toxicity in wistar rats.

Quick Response Code



Journal homepage:

<https://www.easpublisher.com/>

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

Citation: Onyebuchi Obia, Edna Vincent Agorom, Edith Reuben, Augusta Chiamaka Innocent (2025). Effects of Black Pepper on Oxidative Stress Markers in Acrylamide-Induced Toxicity in Male Wistar Rats. *Cross Current Int J Med Biosci*, 7(2), 28-31.

MATERIALS AND METHODS

This study was carried out in the department of Human Physiology, Faculty of Basic Medical Sciences, Rivers State University with ethical approval number: RSU/FBMS/REC/24/104. The experiment involved 20 male wistar rats separated into 4 groups of 5 rats each which were acclimatized for a period of two weeks being provided with standard animal chow and water *ad libitum*. Acrylamide was procured from a chemical shop (Joechem ventures) while black pepper was purchased from the mile 3 market in Port Harcourt and processed according to the method described by Charles *et al.*, 2024. The groups include; Group 1 which served as control and received distilled water and animal chow throughout the period of the experiment. A daily oral dose of 30mg/kg of Acrylamide was respectively used to induce toxicity (Shler *et al.*, 2015) in Groups 2, 3 and 4. Group 2 remained untreated (Acrylamide only). Group 3

received in addition, 50mg/kg/day of black pepper while Group 4 received in addition, 150mg/kg of vitamin E (Ebuehi *et al.*, 2012). All treatment were given orally for 14 days and thereafter animals were sacrificed and blood samples collected to determine the concentrations of oxidative stress markers in the blood. All the parameters were determined using standard methods and values recorded.

Data were analyzed using SPSS vs 23 and presented in Tables. Continuous variables were expressed as mean \pm SEM. The differences between each group were analyzed using paired sample t-test and ANOVA. Values of $p < 0.05$ were considered significant with a confidence level of 95%.

RESULTS

Table 1: Effect of black pepper on oxidative stress markers in acrylamide-induced toxicity in male wistar rats

| GROUP | GSH (mg/dl) | GPx (U/l) | CAT (U/l) | SOD (U/l) | MDA (mg/dl) |
|--------------------|------------------------------|------------------------------|------------------------------|-----------------|-----------------|
| Control | 2.28 \pm 0.11 | 0.06 \pm 0.00 | 3.90 \pm 0.17 | 0.33 \pm 0.02 | 0.48 \pm 0.02 |
| ACR Only | 1.78 \pm 0.05 ^a | 0.04 \pm 0.00 ^a | 2.06 \pm 0.31 ^a | 0.26 \pm 0.04 | 0.50 \pm 0.02 |
| ACR + Black Pepper | 2.12 \pm 0.17 | 0.06 \pm 0.00 ^b | 3.73 \pm 0.46 ^b | 0.46 \pm 0.22 | 0.50 \pm 0.03 |
| ACR + Vitamin E | 2.18 \pm 0.08 ^b | 0.06 \pm 0.00 ^b | 3.79 \pm 0.20 ^b | 0.30 \pm 0.04 | 0.49 \pm 0.04 |

Values are expressed as Mean \pm SEM (n = 5);

^a Significantly different when compared to group 1 (control).

^b Significantly different when compared to group 2 (acrylamide only).

DISCUSSION

Antioxidants are substances which mop up free radicals or other unstable molecules. These free radicals are capable of causing cellular injury and are implicated in various human diseases (Halliwell, 2007; Sies, 2015). Antioxidant enzymes protect major molecules, such as lipids, proteins, and DNA from oxidative damage by inactivating oxidants. These antioxidant enzymes can act in a coordinated way to protect living tissues from oxidative damage (Khan *et al.*, 2011). A wide range of plant-based foods; fruits, vegetables, nuts and seeds possess abundant antioxidants (Serafini & Peluso, 2016; Obia & Auquo, 2018; Maleki *et al.*, 2019) including vitamin C, vitamin E, and polyphenols such as flavonoids, phenolic acid and carotenoids. Glutathione (GSH) is a non-enzymatic antioxidant that neutralizes reactive oxygen species (ROS) and maintains cellular redox homeostasis. The present study showed that acrylamide (ACR) significantly depleted GSH levels ($P < 0.05$). This is so because acrylamide is known to induce oxidative stress by increasing the utilization of GSH, thus depleting its circulatory levels. ACR also inhibits the enzyme, glutathione synthetase which is essential for the production of GSH (Klaunig *et al.*, 2010). Our findings on the effect of ACR on GSH levels agree with other studies which informed the use of ACR to induce oxidative stress in animal studies (Rifai & Saleh, 2020; Zhang *et al.*, 2023).

Administration of black pepper did not show any significant change in the GSH levels of ACR treated rats. In a previous study, the main component of black pepper (piperine) enhanced glutathione synthetase activity (Vijayakumar *et al.*, 2004) with the expected outcome of increasing GSH levels. In our study, GSH levels improved significantly upon treatment with a known antioxidant (vitamin E). In our study, administration of ACR caused significant reduction in the plasma levels of GSH, glutathione peroxidase (GPx) and catalase compared to the control group. Oral administration of black pepper caused a significant rise in the GPx and catalase levels compared to the ACR only group in a similar manner as vitamin E. The findings suggest that supplementation with black pepper and vitamin E respectively mitigated the ACR-induced GPx and catalase depletion. ACR did not affect the plasma concentrations of both supra-oxide dismutase (SOD) and malondialdehyde (MDA) in our study. Again, administration of black pepper had no significant effect on both SOD and MDA levels.

Conclusively, black pepper exerts its antioxidant effects on ACR induced toxicity by improving the GSH and catalase levels but not GPx, SOD and MDA. Asides GSH levels, the antioxidant potentials of black pepper in our study showed a similar pattern as vitamin E.

REFERENCES

- Adimas, M. A., Abera, B. D., Adimas, Z. T., Woldemariam, H. W., & Delele, M. A. (2024). Traditional food processing and Acrylamide formation: A review. *Heliyon*, 10(9), e30258. <https://doi.org/10.1016/j.heliyon.2024.e30258>
- Alefe, M., Abera, B. D., & Delele, M. A. (2024). The Ethiopian snack food ('Kolo'): Existing knowledge and research directions on processing practices, quality and consumption. *Heliyon*, 10(7).
- Banc, R., Popa, D.S., Cozma-Perut, A., Filip, L., Kiss, B., Farcas, A., Nagy A. Miere, D., Loghin, F. (2022): Protective Effects of Wine Polyphenols on Oxidative Stress and Hepatotoxicity Induced by Acrylamide in Rats. *Antioxidants* 11 (1),1347. <https://doi.org/10.3390/antiox1171347>
- Charles, C., Obia, O., Emmanuel, F. D., Ogba, A., & Ojeka, S. O. (2024). Effects of red, green, and black pepper on intestinal motility and post-prandial bicarbonate concentration in guinea pigs. *EAS Journal of Nutrition and Food Sciences*, 6(4), 125-129
- Ebuehi, O. A. T., Ogedegbe, R. A. and Ebuehi. O. M. (2012). Oral administration of vitamin C and Vitamin E ameliorates lead-induced hepatotoxicity and oxidative stress in rat brain. *Nig Q J Hosp Med*. 22(2), 85-90. PMID: 23175903.
- Emmanuel, F. D., Obia, O., Charles, C., Okari, K. A., Otto, J. B., Reuben, E., & Onyeso, G. (2025). Comparative Assessment of Red, Green and Blackpepper species on Plasma and Fecal Lipid Profile of High-fat diet fed wistar rats. *International Journal of Biochemistry Research & Review*, 34(1), 153-163
- Ferrer-Aguirre, A., Romero-Gonzalez, R., Vidal, J. L. M & French A.G. (2016): simple and Fast Determination of Acrylamide and Metabolites in Potato Chips and Grilled Asparagus by Liquid Chromatography coupled to Mass Spectrometry. *Food Anal. Methods* 9 (1), 1237-1245. doi: 10.1007/s 12161-015-0304-6
- Govindaraju, I., Sana, M., Chakraborty, I., Rahman, M. H., Biswas, R., & Mazumder, N. (2024). Dietary Acrylamide: A Detailed Review on Formation, Detection, Mitigation, and Its Health Impacts. *Foods (Basel, Switzerland)*, 13(4), 556. <https://doi.org/10.3390/foods13040556>
- Halliwell, B. (2007). Oxidative stress and inflammation: A joint statement. *Biochemical Society Transactions*, 35(5), 1144-1146.
- Hervert-Hernández, D., Sáyo-Ayerdi, S. G., & Goñi, I. (2010). Bioactive Compounds of Four Hot Pepper Varieties (*Capsicum annum* L.), Antioxidant Capacity and Intestinal Bioaccessibility. *Journal of Agricultural and Food Chemistry*, 58(6), 3399–3406.
- Khan, M.R.; Afzaal, M.; Saeed, N.; Shabbir, M. (2011). Protective Potential of Methanol Extract of *Digera Muricata* on Acrylamide Induced Hepatotoxicity in Rats. *African J. Biotechnol.* 10 (2), 8456–8464.
- Klaunig, J. E., Kamendulis, L. M., & Hocevar, B. A. (2010). Oxidative stress and oxidative damage in carcinogenesis. *Toxicologic pathology*, 38(1), 96–109. <https://doi.org/10.1177/0192623309356453>
- Lingnert, H., Grivas, S., Jägerstad, M., Skog, K., Törnqvist, M., & Åman, P. (2002). Acrylamide in food: mechanisms of formation and influencing factors during heating of foods. *Scandinavian journal of nutrition*, 46(4), 159-172.
- Ludy, M. J., Moore, G. E., & Mattes, R. D. (2012). The effects of capsaicin and capsiate on energy balance: critical review and meta-analyses of studies in humans. *Chemical senses*, 37(2), 103–121
- Maleki, S. J., Crespo, J. F., & Cabanillas, B. (2019). Anti-inflammatory effects of flavonoids. *Food chemistry*, 299, 125124. <https://doi.org/10.1016/j.foodchem.2019.125124>
- Pan, M., Liu, K., Yang, J., Hong, L., Xie, X., & Wang, S. (2020). Review of Research into the determination of acrylamide in foods. *Foods*, 9(4), 524.
- Rifai, L., & Saleh, F. A. (2020). A review on acrylamide in food: Occurrence, toxicity, and mitigation strategies. *International journal of toxicology*, 39(2), 93-102.
- Serafini, M., & Peluso, I. (2016). Functional Foods for Health: The Interrelated Antioxidant and Anti-Inflammatory Role of Fruits, Vegetables, Herbs, Spices and Cocoa in Humans. *Current pharmaceutical design*, 22(44), 6701–6715. <https://doi.org/10.2174/1381612823666161123094235>
- Sharifi-Rad, M., Anil Kumar, N. V., Zucca, P., Varoni, E. M., Dini, L., Panzarini, E., Rajkovic, J., Tsouh Fokou, P. V., Azzini, E., Peluso, I., Prakash Mishra, A., Nigam, M., El Rayess, Y., Beyrouthy, M. E., Polito, L., Iriti, M., Martins, N., Martorell, M., Docea, A. O., Setzer, W. N., ... Sharifi-Rad, J. (2020). Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Frontiers in physiology*, 11, 694. <https://doi.org/10.3389/fphys.2020.00694>
- Shler, A.F. Mahmood., Kawa, A.M. Amin., and Shilan F.M. Salih (2015). Effect of Acrylamide on Liver and Kidneys in Albino Wistar Rats. *International Journal of Current Microbiology and Applied Sciences* 4(5), 434-44.
- Sies, H., & Jones, D. P., (2020). Oxidative Stress: A Concept in Redox Biology and Medicine, *Redox Biology*, 23, 101-112.
- Timmermann, C. A. G., Mølck, S. S., Kadawathagedara, M., Bjerregaard, A. A., Törnqvist, M., Brantsæter, A. L., & Pedersen, M. (2021). A Review of Dietary Intake of Acrylamide in Humans. *Toxics*, 9(7), 155. <https://doi.org/10.3390/toxics9070155>

- Vijayakumar, R. S., Surya, D., & Nalini, N. (2004). Antioxidant efficacy of black pepper (*Piper nigrum* L.) and piperine in rats with high fat diet induced oxidative stress. *Redox report : communications in free radical research*, 9(2), 105–110. <https://doi.org/10.1179/135100004225004742>
- Yan, F., Wang, L., Zhao, L., Wang, C., Lu, Q., & Liu, R. (2023). Acrylamide in food: Occurrence, metabolism, molecular toxicity mechanism and detoxification by phytochemicals. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*, 175, 113696.
- Zhang, N. N., Zhou, Q., Fan, D. M., Xiao, J., Zhao, Y. L., Cheng, K. W., *et al* (2021). Novel roles of hydrocolloids in foods: Inhibition of toxic maillard reaction products formation and attenuation of their harmful effects. *Trends in Food Science and Technology*. 111:706-715.
- Zhang, L., Yang, L., Luo, Y., Dong, L., & Chen, F. (2023). Acrylamide-Induced Hepatotoxicity Through Oxidative Stress: Mechanisms and Interventions. *Antioxidants & redox signaling*, 38(16-18), 1122–1137. <https://doi.org/10.1089/ars.2022.0055>
- Zheng, J., & Xiao, H. (2022). Editorial: The Effects of Food Processing on Food Components and Their Health Functions. *Frontiers in nutrition*, 9, 837956. <https://doi.org/10.3389/fnut.2022.837956>