

Research Article

Induction of Autophagy in Human Myeloid and Lymphoid Leukaemia Cell Line by Using Polyphenols Alone and Combined with A Stander Chemotherapy

Abdu-Alhameed A. Ali Azzwali¹, Azab Elsayed Azab² and Abdalrauf M.A.B. Alfourt³¹Department of Medical Laboratory, Faculty of Medical Technology, Sabratha University, Surman, Libya²Department of Physiology, Faculty of Medicine, Sabratha University, Sabratha, Libya³Department of Medical Laboratory, Faculty of Medical Technology, Zawia University, Zawia, Libya

*Corresponding Author

Azab Elsayed Azab

Abstract: Background: Autophagy is a major catabolic mechanism that contributes to the degradation of dysfunctional or unnecessary components in cells. The main function of autophagy depends on the variant cellular contexts and inducers. The results of autophagy can stimulate either cell death or cell survival. Autophagy is known as a protective process for cell survival if it provides the cell with the necessary requirements and nutrients, and removes organelles that were damaged. In contrast, anti-cancer drugs are often able to activate autophagy and in this case, autophagy induces apoptotic cell death. The high level of polyphenols found in some fruits and vegetables has been reported to be potentially chemotherapeutic and having a considerable effect on haematological malignancies. **Objectives:** To explore the effect of a stander chemotherapy (Imatinib, Doxorubicin), and/or polyphenols (emodin, rhein, apigenin and cis-stilbene) on inducing autophagy of myeloid and lymphoid leukaemia cells compared to non-tumour cells. **Methods:** one myeloid (K562), one lymphoid (CCRF-CEM), a leukaemia cell line and one non-tumour normal cell line (CD133) were treated with different doses of polyphenols. Using a CYTO-ID® green detection reagent and Hoechst 33342 nuclear stain to detect autophagy by fluoresent microscope. **Results:** Emodin, cis-stilbene, apigenin and rhein showed different levels of effect on inducing autophagy in K562 myeloid cells, CCRF-CEM lymphoid cells and CD133+ normal cells when these types of polyphenols used separately and combined with chemotherapy. **Conclusion:** Generally, the CCRF-CEM lymphoid leukaemia cell line was more sensitive to polyphenol treatments alone and when combined with Doxorubicin compared to the K562 myeloid leukaemia cell line and CD133+ non-tumour cells. These results suggest that polyphenols have different effects according to the type of cell and polyphenol. The variant effect between leukaemia cells and non-tumour cell suggests that polyphenols are a potentially therapeutic agent for leukaemia. Polyphenols can enhance the effect of chemotherapy and reduce the required dose to induce cell death in cancer cells.

Keywords: Induction of Autophagy, Imatinib, Doxorubicin, Polyphenols, Polyphenols combined with chemotherapy., Human Myeloid and Lymphoid Leukaemia Cell Line.

1. INTRODUCTION

Autophagy is a major catabolic mechanism that contributes to the degradation of dysfunctional or unnecessary components in cells. It is also known as macro autophagy, which is a dynamic and an inherently regulated mechanism related to the creation of auto phagosomes and cytoplasmic vesicles, with a double-membrane that engulf cellular constituents (JING, K., & LIM, K. 2012). The formation of auto-phagosomes begins with phagophores, also known as sequestering membranes and is followed by conjugating

microtubule-associated protein (light chain 3) to the phagophores and controlling their elongation (XIE, Z. *et al.*, 2008). As the sequestering membrane expands, it engulfs cytoplasmic components that contain organelles such as the endoplasmic reticulum, aggregated proteins, mitochondria and foreign microorganisms (virus and bacteria). When the auto-phagosome is created, it is fused with lysosome to become autolysosome and degrades the cargo with LC3 and autophagy from the outer membrane. This process acts as a regulatory mechanism for maintaining a homeostatic balance in

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easjpp/>

Article History

Received: 30.03.2019

Accepted: 15.04.2019

Published: 23.05.2019

Copyright © 2019 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

DOI: 10.36349/easjpp.2019.v01i03.001

mammalian cells, by clearing injured organelles and reprocessing autophagy-derived nutrients (Figure 1) (JING, K., & LIM, K. 2012).

However, the main function of autophagy depends on the variant cellular contexts and inducers. The results of autophagy can stimulate either cell death or cell survival. For example, autophagy is known as a protective process for cell survival if it provides the cell with the necessary requirements and nutrients, and

removes organelles that were damaged. In contrast, anti-cancer drugs are often able to activate autophagy and in this case, autophagy induces apoptotic cell death (Amaravadi, R. K. *et al.*, 2007; Song, K. S. *et al.*, 2008). Nonetheless, autophagy damage can delay or stop the development of cell death (Yuk, J. M. *et al.*, 2010), and in other instances, induce cell death by itself. Cell death enhanced by autophagy may be associated with depreparing essential factors for cell survival (Yu, L. *et al.*, 2006).




Figure 1. The different stages of autophagy formation (DANIELLE, G. *et al.*, 2012).

Formation of the phagophore is the first step in autophagy, which conjugates with light chain 3 to control the elongation of the phagophore. The expansion of the isolation membrane (phagophore) leads to formation of the auto-phagosome, which can engulf cytoplasmic components containing organelles and foreign microorganisms (virus and bacteria). When the outer membrane of the auto-phagosome fuses with a lysosome (docking and fusion steps), it forms an autophagolysosome. Finally, the sequestered substance is destroyed inside the autophagolysosome (vesicle breakdown and degradation) and recycled.

The aim of this study was to determine the impact of polyphenols (emodin, apigenin, cis-stilbene and rhein) for inducing autophagy in one myeloid cell line (K562), one lymphoid cell line (CCRF-CEM) and one non-tumour normal cell line (CD133). The polyphenol doses and chemotherapy were then used separately and combined together to investigate if there was an antagonistic or synergistic effect for inducing autophagy.

2. METHODS AND MATERIALS:

2.1. Media preparation:

50 ml of RPMI media was removed into falcon tube. Then, 50 ml of fetal calf serum (FCS) was added, followed by 5 ml (1.5 mmol/L L-glutamine) and 5 ml 1 % penicillin/streptomycin to protect media from contamination by bacteria.

2.2. Cell Culture

Two leukaemia cell lines were obtained from the American Type Culture Collection (ATCC; Middlesex, U.K.). One myeloid cell lines: K562 (chronic myeloid leukaemia) (ATCC: CCL-243, Middlesex, UK) and one lymphoid cell line CCRF-CEM (acute lymphoblastic leukaemia) (ATCC: CCL-119, Middlesex, UK) together with the non-tumour (CD133) positive hematopoietic stem cells. These cell lines were maintained and cultured in RPMI 1640 medium (Invitrogen, Paisley, U.K.) supplemented with 10% (v/v) fetal bovine serum, 1.5 mmol/L L-glutamine, and 1 % penicillin/streptomycin (complete media) and then incubated in an atmosphere of 5% CO₂ at 37°C.

2.3. Cell Feeding

The cells needed to be fed and split every three days, which they were split into 1:4 flasks each Friday and fed with fresh media every 48 to 72 h. This was performed by centrifuging the cells for 5 minutes at 1500 rpm, and then re-suspended in new fresh media. If large quantities of suspension cells are not required, cells were fed by discarding a part of cells and spent media in a clean bleach pot and 20 ml of fresh media was added (Do not overfill > 40ml). These were repeated Monday, Wednesday and on Friday, cell lines were split as described earlier.

2.4. Procedures to Prevent Contamination:

Rings, watches or any decorative ornaments on the arms were removed and a clean lab coat was worn before contacting with cells. Hands were then washed up and using antibacterial detergent to clean the fume hood surfaces including the bottom surface under the plates. Gloves were changed and sprayed after contact with surfaces outside the fume hood or when working with a different cell line. All equipment was sprayed with bactericidal spray before entering the fume hood.

2.5. Counting Cells

Cells were counted using the haemocytometer by calculating the total number of 5 small squares in the central square (4 in corner squares and one in the centre square). After the cells counted, they were then multiplied by 5 and by 1×10^4 to represent cells in a 0.02mm-cubed volume and in 1000 cubic millilitre, respectively (cells number $X \ 5 \ X \ 10^4$). For example, $32 \times 5 \times 10^4 = 160 \times 10^4$, for 96 well plate 2.5×10^3 cells per well was required; therefore, for 100 wells 2.5×10^5 cells were needed which $160 \times 10^4 = 16 \times 10^5$ /ml, and $16 \times 10^5 = 1000 \ \mu\text{l}$ from this $1 \times 10^5 = 1000/16 = 1 \times 10^5 = 62.5 \ \mu\text{l}$. So, $2.5 \times 10^5 = 62.5 \times 2.5 = 2.5 \times 10^5 = 156 \ \mu\text{l}$ from cell suspension was added to 9844 μl media to get 10000 μl for 96 wells, then 100 μl was pipetted to each well and treated with 5 μl polyphenols as described above.

2.6. Polyphenols Treatment Preparation:

Polyphenols (Emodin, Apigenin, Rhein and Cis-stilbene) was prepared to the required concentration (0, 2, 10, 50 and 250 μM) from an original master stock. The polyphenols master stocks for emodin and apigenin were prepared by dissolving and sterilizing 27.024 mg, in 100 μl of 100 % ethanol (Sigma). Then, dissolved in 900 μl serum free media (SFM) (Invitrogen) to make the final concentration of 100mMol/ml with a 10% ethanol (100000 $\mu\text{mol/ml}$). The apigenin has the same molecular weight as emodin, so it prepared by the same procedures. Rhein also was prepared by the same methods just 28.422 mg was weighted from rhein powder instead of 27.024 mg. Cis-Stilbene provided dissolved as a 96% pure solution in a glass bottle.

The polyphenols dosages were prepared according to the well plate was used. For 96 well 5 μl of polyphenol was added to 100 μl of cells suspension per well. This is 1:20 therefore, the stock needed to be X20 more concentrated. For 24 and 12 well plates that were used for apoptosis and autophagy experiments which 10 μl of polyphenols was added to 1ml cell suspension. This is a 1:100 therefore the stock needed to be X100 more concentrated which were prepared as follow:

- To make 0 μM , 1000 μl serum free media was taken = 0 μmol
- To make 2 μM , 2 μl of stock was added to 998 μl (SFM) = 200 μmol .
- To make 10 μM 10 μl of stock was added to 990 μl (SFM) = 1000 μmol .

- To make 50 μM 50 μl of stock was added to 950 μl (SFM). = 5000 μmol .
- To make 250 μM , 250 μl of stock was added to 750 μl (SFM). = 25000 μmol .

For 96 well plate 100 μl of cells was taken and treated with 5 μl of polyphenol, so there was a 1:20 dilution, therefore all stocks were 20 times more concentrated which were prepared by using the above concentration as follow:

- 20 μl of 200 μmol stock was added to 80 μl SFM to make 2 μM stock = 40 μmol .
- 20 μl of 1000 μmol stock was added to 80 μl SFM to make 10 μM stock = 200 μmol .
- 20 μl of 5000 μmol stock was added to 80 μl SFM to make 50 μM stock = 1000 μmol .
- 20 μl of 25000 μmol stock was added to 80 μl SFM to make 250 μM stock = 5000 μmol .

2.7. Preparation of standard chemotherapy agents:

2.7.1. Imatinib:

14.7 mg of imatinib powder was dissolved in 1 ml sterile microbiology filtered distilled water. This made solution with 25 mmol solution. For 96 well plate, 1 μMol concentration was needed out of the well. However, in well it should be 20 times stronger because 5 μl / 100 μl this is a 1/20. This prepared by taking 4 μl from super stock (25 mmol) and added to 4996 μl SFM with 10% ethanol to form 5 ml Imatinib with (1 μMol) concentration.

2.7.2. Doxorubicin:

14.5 mg from doxorubicin powder was dissolved in 1 ml sterile microbiology filtered distilled water. This made solution with 25 mmol solution. For 96 well plate, 1 μMol concentration were needed out of the well. However, in well it should be 20 times strong because 5 μl / 100 μl this is a 1/20. This prepared by taking 5 μl from super stock (25 mmol) and added to 4995 μl SFM with 10% ethanol to form 5 ml Doxorubicin with (1 μMol) concentration.

2.8. Seeding the Cells:

Cells were seeded into 96-well plate (Fisher Scientific, Leicestershire, United Kingdom) at 2.5×10^3 cells per well and then treated with different concentration (0, 2, 10, 50, 250) μM of different types of polyphenols according to cell type. For example, emodin (Sigma, Poole, UK) was used for (CCRF-CEM, K562 and CD133), apigenin and cis-stilbene (Sigma, Poole, UK) were used for (K562 and CD133) and rhein (Sigma, Poole, UK) was used for (CCRF-CEM and CD133). All treatments were done in triplicate, and incubated at 37 $^{\circ}\text{C}$ for 24 hours. Then, 96 well plate and its contents were equilibrated at room temperature (RT) for 30 minutes. After that, 10 μl of reagent was added to each well and mixed well on an orbital shaker at 400 rpm to induce cell lysis and incubated for 10 minutes at RT to stabilize the luminescent signal. Then, 96 well

plat was read by using Wallace 1420 luminescence detector (PerkinElmer, Waltham, USA). Control was prepared with media without cells.

In a white 96 well plate standard curve of the luminescence was designed against cell number between 0 and 50000 cells per well to ensure luminescence was in the linear range. The density of cells that prepared was: 50,000; 25,000; 12,500; 6250; 3125 1562, 780 and 0. Then, IC₅₀ was determined from a linear regression equation of each standard curve for each polyphenol with each cell line which is known as the treatment concentration at which 50% inhibition in cellular proliferation was detected which was performed to determine the effectiveness of treatments. The IC₂₅ and IC₁₀ were also calculated in order to provide treatment ranges for apoptosis detection.

2.9. Combination of Polyphenols with Chemotherapy:

After IC₅₀, IC₂₅ and IC₁₀ were determined, cells were seeded into 96-well plate (Fisher Scientific, Leicestershire, United Kingdom) with 100 µl at 2.5 x10³ cells per well as previously described and then treated with 5 µl of Imatinib for (K562 and CD133) or Doxorubicin for (CCRF-CEM and CD133) alone and combined with 5 µl of IC₅₀, IC₂₅ and IC₁₀ (LSD) of different types of polyphenols according to cell type as above. All treatments were done in triplicate, in three independent experiments and incubated at 37°C for 24 hours. Then, completed as described above. This step was performed to see if there is any synergistic result for polyphenols when combined with chemotherapy.

2.10. Autophagy Assay

CYTO-ID® Autophagy Detection Kit has been optimized for measuring autophagy in live cells by using a novel dye that selectively marks autophagic vacuoles. This was detected by using fluorescence microscope. The best way of observing autophagic activity is by measuring the increased numbers of autophagosomes in cells responding to induction. Monitoring autophagic flux provides a meaningful way the source of auto phagosomes. Rapamycin and starvation are known as inducer of autophagy (positive control), Chloroquine is a lysosome inhibitor. Rapamycin and Chloroquine were used as positive control.

2.10.1. Reagent preparation

2.10.1.1. Rapamycin

Rapamycin was provided lyophilized, so it was re-suspended in 50 µl of DMSO. This formed 500 µM stock solutions.

2.10.1.2. Chloroquine

Chloroquine was also supplied lyophilised (7.5 µmoles) which was centrifuged to gather the material at the bottom of the tube and reconstituted in 125 µl deionized water for 60 mM stock solution.

2.10.1.3. 1x assay buffer

1 ml from 10x assay buffer that was provided was taken and added to 9 ml deionized water.

2.10.1.4. -CYTO-ID® Green detection reagent

This was prepared by taking 1 ml of 1x assay buffer, then 2 µl of cyto-ID® green detection reagent and 1 µl of Hoechst 33342 nuclear stain were added and mixed well.

2.10.2. Procedure

The autophagy assay was performed for cells that showed synergistic effects of polyphenols when combined with chemotherapy.

24 well plates were prepared for each cell line at 1 x 10⁶ cells per well. 1 ml of 1 x 10⁶ cell were pipetted to each well and treated with 10 µl of IC₅₀, IC₂₅, and IC₁₀ of polyphenols, chemotherapy agents and combination of chemotherapies with polyphenols in the cell that showed synergistic results. In addition, the CYTO-ID® Autophagy Detection Kit was pipetted to four well as follow: 10 DMSO, Rapamycin, Chloroquine, and Rapamycin and Chloroquine together in four wells. Then, incubated at 37°C with 5% CO₂ for 18 hours. Then, cells were collected in Eppendorf tubes and centrifuged at 2000 rpm for 3 min. After that, cells were washed twice with 1x Assay Buffer. After the supernatant was removed, about 100 µl of Microscopy Dual Detection Reagent solution was added to each tube and re-suspended by pipette. The samples were after that covered with foil to protect them from the light and incubated at 37°C for 30 min. Another time cells were washed twice with 1x assay Buffer and added 4% paraformaldehyde for 20min to fix cells. Cells were then washed again three times using 1x Assay Buffer. Finally, the supernatant was removed and pellet re-suspended in 100 µl of 1x Assay Buffer. Then, one drop was applied on the microscope slides and a drop of immersion oil was added and coverslip was also applied and sealed with nail varnish.

Slides were analysed by fluorescent microscopy (Olympus, BX60). Cells were analysed by fluorescent microscopy using magnification 400. Pictures were taken by Q capture-pro 8.0 (UVP Bio Imaging Systems) and saved on memory stick.

3. RESULTS

3.1. Effect of combining chemotherapy and polyphenols on the induction of Autophagy in leukaemia cell lines after 24h treatments

Cell lines that showed an additive or synergistic effect on the inhibition of ATP levels when treated with IC₅₀, IC₂₅ and IC₁₀ of polyphenols combined with chemotherapy (Imatinib or Doxorubicin) were also investigated for their effect on the induction of autophagy. After treatments with the selected polyphenols/chemotherapy combination treatments, cells were stained using a CYTO-ID® green detection reagent and Hoechst 33342 nuclear staining, and

examined by florescent microscope, and images were captured. A DMSO negative control and Rapamycin and Chloroquine positive control were also included. By looking to the images that were collected and comparing them to the positive and negative controls, it was concluded that K562 showed autophagic induction when treated with 0.05µM of Imatinib and when

Imatinib was combined with the IC₂₅ of emodin and cis-stilbene (8 µM and 28 µM) respectively. CCRF-CEM cell lines that were treated with 0.05µM of Doxorubicin also showed the induction of autophagy, which is indicated by the arrow. In contrast, CD133 cells did not indicate any autophagic induction following treatment (Figure 2).




Figure 2. Atypical example of autophagic induction in K562 myeloid leukaemia cells, CCRF-CEM lymphoid leukaemia cells and the CD133+ non-tumour cells following treatment combining polyphenols and chemotherapy for 24h and stained using a CYTO-ID® green detection reagent and Hoechst 33342 nuclear staining. DMSO was used as a negative control, while Rapamycin, Rapamycin and Chloroquine were used as positive controls. The results were determined by comparing images of treated cells to the negative and positive controls.

4. DISCUSSION

Nonetheless, the results of this study are consistent with existing evidence on the anticancer properties of polyphenol. A number of research studies have identified the polyphenols apigenin (Zhu, Y. *et al.*,

2013; Shukla, S. *et al.*, 2014), cis-stilbene (Mahbub, A. *et al.*, 2013; Chen, Y. C. *et al.*, 2002), and rhein (Mahbub, A. *et al.*, 2013; Duraipandiyar, V. *et al.*, 2012; Bounda, G. A. *et al.*, 2015) as having anti-proliferative properties.

Polyphenols have equally been shown in several studies to exhibit synergistic and antagonistic effects with chemotherapeutics (Mertens-Talcott, S.U., & Percival, S.S. 2005; Lecumberri, E. *et al.*, 2013).

In this study, results indicated that when 13 μ M (IC₅₀) of emodin and 53 μ M (IC₅₀) of Cis-Stilbene were combined with the chemotherapeutic Imatinib, a 50% increase in cell death of K562 myeloid cells was recorded. Similar results were observed when 150 μ M (IC₅₀) of emodin and 140 μ M (IC₅₀) of rhein were administered alongside the chemotherapeutic Doxorubicin to CCRF-CEM lymphoid cells, which showed a 60% increase in cell death. Likewise, autophagy was observed in K562 myeloid cells and CCRF-CEM lymphoid cells when treated with 0.05 μ M of Imatinib and Doxorubicin respectively and also in K562 when 8 μ M (IC₂₅) of emodin and 28 μ M (IC₂₅) of Cis-Stilbene were combined with the chemotherapeutic Imatinib.

According to some researchers (Arora, A., & Scholar, E. M. 2005; Han, W. *et al.*, 2011), Imatinib inhibits cancer cell growth and proliferation by blocking BCR-ABL kinase, down regulating the MAP kinase and PI3K/Akt pathways and the activation of RSK kinases. It thus promotes emodin and cis-stilbene-induced apoptosis by aggravating cell proliferation in myeloid leukaemia cells. Likewise, the mechanism of action of Doxorubicin involves the up regulation of the production of ROS which triggers autophagy and apoptotic pathways (Thorn, C. F. *et al.*, 2011; Tacar, O. *et al.*, 2013).

A study by Jia *et al.*, (2009) demonstrated that some types of polyphenols are able to induce autophagy in a chronic myeloid leukemia cell line via down regulation of the Bcl-2 protein. These actions were shown when the potent inhibitor of autophagosome-lysosome fusion, bafilomycin A1 and pan-caspase inhibitor zVAD-FMK blocked the cell death. In another study on prostate cancer cells, curcumin was exposed to induce autophagy cell death by down regulation of another important Bcl-2 family member, Bcl-xL. Curcumin did not stimulate the cleavage of procaspase-8, -9, -3, or -7 or PARP however cause appearance of the LC3B-II isoform and led to increase the number of autophagosomes (Teiten, M. H. *et al.*, 2011).

CONCLUSION

It can be deduced from the results of this study that cancer cells display variable sensitivity to the anti-cancer activity of different polyphenols and this depended on the type of leukaemia cell lineage (myeloid vs lymphoid). It has been shown that lymphoid cell line (CCRF-CEM) was more sensitive to polyphenols than myeloid cell line (K562). The anti-cancer activities of polyphenols have been shown to be significantly improved by chemotherapeutic in this in similar studies which also induced autophagy in

leukaemia cells. This suggests that the molecular mechanism of action of polyphenols may differ according to the type of polyphenols and cell lines. It was proved in this study that polyphenols have the capability to synergize the effect of chemotherapeutic drugs and decrease the dosage required to induce autophagy in cancer cells. However, more research is still required to improve current understanding of the action of polyphenols alone and polyphenol-chemotherapeutic interactions to advance anti-proliferative and autophagic activities in cancer cells. Future studies may be required to investigate the exact mechanism of action of polyphenols in induction of autophagy cell death in leukaemia and findings of pharmacokinetic and pharmacodynamics properties of polyphenols.

REFERENCES

1. JING, K., & LIM, K. (2012). Why is autophagy important in human diseases? *Experimental & molecular medicine*, 44 (2), 69-72.
2. XIE, Z., NAIR, U., & KLIONSKY, D. J. (2008). Atg8 controls phagophore expansion during autophagosome formation. *Mol Biol Cell*, 19, 3290-3298.
3. Amaravadi, R. K., Yu, D., Lum, J. J., Bui, T., Christophorou, M. A., Evan, G. I., ... & Thompson, C. B. (2007). Autophagy inhibition enhances therapy-induced apoptosis in a Myc-induced model of lymphoma. *The Journal of clinical investigation*, 117(2), 326-336.
4. Song, K. S., Kim, J. S., Yun, E. J., Kim, Y. R., Seo, K. S., Park, J. H., ... & Lim, K. (2008). Rottlerin induces autophagy and apoptotic cell death through a PKC-delta-independent pathway in HT1080 human fibrosarcoma cells: the protective role of autophagy in apoptosis. *Autophagy*, 4(5), 650-658.
5. Yuk, J. M., Shin, D. M., Song, K. S., Lim, K., Kim, K. H., Lee, S. H., ... & Jo, E. K. (2010). Bacillus calmette-guerin cell wall cytoskeleton enhances colon cancer radiosensitivity through autophagy. *Autophagy*, 6(1), 46-60.
6. Yu, L., Wan, F., Dutta, S., Welsh, S., Liu, Z., Freundt, E., ... & Lenardo, M. (2006). Autophagic programmed cell death by selective catalase degradation. *Proceedings of the National Academy of Sciences*, 103(13), 4952-4957.
7. DANIELLE, G., SANDRA, B., & KAY, F. (2012). Autophagy: cellular and molecular mechanisms. *JPathology*, 221 (1), 3-12.
8. Zhu, Y., Mao, Y., Chen, H., Lin, Y., Hu, Z., Wu, J., & Xie, L. (2013). Apigenin promotes apoptosis, inhibits invasion and induces cell cycle arrest of T24 human bladder cancer cells. *Cancer Cell Int*, 13(1), 54.
9. Mahbub, A., L Le Maitre, C., L Haywood-Small, S., J McDougall, G., A Cross, N., & Jordan-Mahy, N. (2013). Differential effects of polyphenols on proliferation and apoptosis in human myeloid and lymphoid leukemia cell lines. *Anti-Cancer Agents*

- in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents), 13(10), 1601-1613
10. Shukla, S., Fu, P., & Gupta, S. (2014). Apigenin induces apoptosis by targeting inhibitor of apoptosis proteins and Ku70–Bax interaction in prostate cancer. *Apoptosis*, 19(5), 883-894.
 11. Chen, Y. C., Shen, S. C., Lee, W. R., Hsu, F. L., Lin, H. Y., Ko, C. H., & Tseng, S. W. (2002). Emodin induces apoptosis in human promyeloleukemic HL-60 cells accompanied by activation of caspase 3 cascade but independent of reactive oxygen species production. *Biochemical pharmacology*, 64(12), 1713-1724.
 12. Duraipandiyar, V., Baskar, A. A., Ignacimuthu, S., Muthukumar, C., & Al-Harbi, N. A. (2012). Anticancer activity of Rhein isolated from *Cassia fistula* L. flower. *Asian Pacific journal of tropical disease*, 2, S517-S523
 13. Catanzaro, D., Vianello, C., Ragazzi, E., Caparrotta, L., & Montopoli, M. (2014). Cell cycle control by natural phenols in cisplatin-resistant cell lines. *Natural product communications*, 9(10), 1465-1468.
 14. Bounda, G. A., Zhou, W., Wang, D. D., & Yu, F. (2015). Rhein elicits in vitro cytotoxicity in primary human liver HL-7702 cells by inducing apoptosis through mitochondria-mediated pathway. *Evidence-Based Complementary and Alternative Medicine*, Article ID 329831, 19 pages
HYPERLINK
"<http://dx.doi.org/10.1155/2015/329831>"
<http://dx.doi.org/10.1155/2015/329831>
 15. Mertens-Talcott, S.U., & Percival, S.S. (2005). Ellagic acid and quercetin interact synergistically with resveratrol in the induction of apoptosis and cause transient cell cycle arrest in human leukemia cells. *Cancer letters*, 218(2), 141-151.
 16. Adhami, V. M., Malik, A., Zaman, N., Sarfaraz, S., Siddiqui, I. A., Syed, D. N., & Mukhtar, H. (2007). Combined inhibitory effects of green tea polyphenols and selective cyclooxygenase-2 inhibitors on the growth of human prostate cancer cells both in vitro and in vivo. *Clinical Cancer Research*, 13(5), 1611-1619.
 17. Mazumder, M. E. H., Beale, P., Chan, C., Yu, J. Q., & Huq, F. (2012). Epigallocatechin gallate acts synergistically in combination with cisplatin and designed trans-palladiums in ovarian cancer cells. *Anticancer research*, 32(11), 4851-4860.
 18. Lecumberri, E., Dupertuis, Y.M., Miralbell, R., & Pichard, C. (2013). Green tea polyphenol epigallocatechin-3-gallate (EGCG) as adjuvant in cancer therapy. *Clin Nutr*. 32(6), 894–903.
 19. Arora, A., & Scholar, E. M. (2005). Role of tyrosine kinase inhibitors in cancer therapy. *Journal of Pharmacology and Experimental Therapeutics*, 315 (3), 971-979.
 20. Bantscheff, M., Eberhard, D., Abraham, Y., Bastuck, S., Boesche, M., Hobson, S., & Reader, V. (2007). Quantitative chemical proteomics reveals mechanisms of action of clinical ABL kinase inhibitors. *Nature biotechnology*, 25(9), 1035-1044.
 21. Han, W., Pan, H., Chen, Y., Sun, J., Wang, Y., Li, J., & Wang, X. (2011). EGFR tyrosine kinase inhibitors activate autophagy as a cytoprotective response in human lung cancer cells. *PLoS one*, 6(6), e18691.
 22. Thorn, C. F., Oshiro, C., Marsh, S., Hernandez-Boussard, T., McLeod, H., Klein, T. E., & Altman, R. B. (2011). Doxorubicin pathways: pharmacodynamics and adverse effects. *Pharmacogenetics and genomics*, 21(7), 440–446.
 23. Tacar, O., Sriamornsak, P., & Dass, C. R. (2013). Doxorubicin: an update on anticancer molecular action, toxicity and novel drug delivery systems. *Journal of Pharmacy and Pharmacology*, 65(2), 157-170.
 24. Jia, Y. L., Li, J., Qin, Z. H., & Liang, Z. Q. (2009). Autophagic and apoptotic mechanisms of curcumin-induced death in K562 cells. *Journal of Asian natural products research*, 11(11), 918-928.
 25. Teiten, M. H., Gaascht, F., Cronauer, M., Henry, E., Dicato, M., & Diederich, M. (2011). Anti-proliferative potential of curcumin in androgen-dependent prostate cancer cells occurs through modulation of the Wnt signaling pathway. *International journal of oncology*, 38(3), 603-611.