

Research Article

A Retrospective Analysis of Outcome in Dengue Patients Admitted into a Multidisciplinary ICU of a Tertiary Care Teaching Hospital

Bharath Cherukuri*¹, Bala Narasimha Rao Guna², Neeraj Guttikonda³ & Rajendra kumar Medikonda⁴¹Assistant Professor, Department of Critical Care Medicine, NRI Academy Sciences, Andhra Pradesh, India²Senior Resident, Department of Critical Care Medicine, NRI Academy of Sciences, Andhra Pradesh, India³Junior Resident, Department of Critical Care Medicine, NRI Academy of Sciences, Andhra Pradesh, India⁴Junior Resident, Department of Critical Care Medicine, NRI Academy of Sciences, Andhra Pradesh, India**Article History**

Received: 15.04.2020

Accepted: 09.05.2020

Published: 15.05.2020

Journal homepage:<https://www.easpublisher.com/easjacc>**Quick Response Code**

Abstract: Introduction :World wide dengue has become the most common vector bone tropical fever spread by Aedes Egypti. This virus has four circulating serotypes worldwide. In urban and suburban areas of high endemicity this infection is responsible for high mortality and morbidity. WHO estimates incidence of dengue globally ranges between 50 million to 200 million every year. Dengue fever presentations can vary from fatigability and malaise to severe shock and multiorgan dysfunction syndrome. **Methods** All patients had a full work up for fever that included three smears for malarial parasites, serology for dengue, leptospirosis, scrub typhus, enteric fever, blood, urine, sputum or endotracheal cultures and other tests as clinically indicated. A diagnosis of Dengue was made when a patient is tested positive for NS1-Antigen Strip® ELISA (BioRad), IgM and IgG- Capture ELISA Panbio® (Alere™) and other causes of fever excluded. Patients were treated with Ceftriaxone along with enteral doxycycline and intravenous azithromycin as needed. **Results** Of total patients, the vasopressor requirements being Dopamine (12.9%), Noradrenaline (12.9%) & Vasopressin (3.2%). 1.6% received packed red blood cell transfusions whereas 80.6% had platelet transfusions. 4.8% required non-invasive ventilation whereas 9.7% required invasive mechanical ventilation and 3.2% required both non-invasive and invasive mechanical ventilation. 6.5% of patients had received haemodialysis. The mean duration of ICU and hospital stay were 4.25 and 6.54 days respectively. In-hospital mortality was 1.6%. **Conclusion** In our study we hereby conclude the incidence of clinical features is in agreement with other studies whereas the in-hospital mortality was relatively low.

Keywords: Dengue, Shock, ICU, ARDS, AKI.

Copyright @ 2020: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (Noncommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION:

World-wide dengue has become the most common vector bone tropical fever spread by Aedes Egypti (Murray, N. E. A. *et al.*, 2013; Guzman, M.G. & Harris, E. 2015). This virus has four circulating serotypes worldwide. In urban and suburban areas of high endemicity this infection is responsible for high mortality and morbidity (Anand, A. M. *et al.*, 2016). WHO estimates incidence of dengue globally ranges between 50 million to 200 million every year (Murray, N. E. A. *et al.*, 2013). Dengue fever presentations can vary from fatigability and malaise to severe shock and multiorgan dysfunction syndrome. Out of seven identified countries in south east Asia, India is one of them reporting Dengue fever and Dengue hemorrhagic fever regularly. Even though evolution of illness and dengue prodrome are similar in many patients, progress and outcome differ significantly depending upon severity of illness and also treatment given.

METHODS:

All patients admitted to the multidisciplinary ICU of a tertiary care teaching hospital in South India with acute febrile illness over a one year period (2017 – 2018) were analysed retrospectively.

All patients had a full work up for fever that included three smears for malarial parasites, serology for dengue, leptospirosis, scrub typhus, enteric fever, blood, urine, sputum or endotracheal cultures and other tests as clinically indicated. A diagnosis of Dengue was made when a patient is tested positive for NS1-Antigen Strip® ELISA (BioRad), IgM and IgG- Capture ELISA Panbio® (Alere™) and other causes of fever excluded. Patients were treated with Ceftriaxone along with enteral doxycycline and intravenous azithromycin as needed.


Organ support was provided to patients depending upon the organ dysfunction. Respiratory support includes non-invasive ventilation (NIV) or invasive mechanical ventilation (IMV), cardiovascular support includes vasopressor therapy for patients in shock and renal replacement therapy for those with acute kidney injury.


Severity of illness scores (Acute Physiology and Chronic Health Evaluation II, APACHE-II) and Sequential Organ Failure Assessment (SOFA) scores were calculated at admission. The primary outcome of interest was in-hospital mortality. Other outcomes were duration of ICU and hospital stay respectively.

RESULTS

TABLE I. Demographic and clinical profile of patients with dengue

Variable	n (%)	Mean
Total	62	
Age		44.19
Sex		
Male	35 (56.46)	
Female	27 (43.54)	
Symptoms & Signs		
Fever	60 (96.8)	
Bleeding	23 (37.1)	
Rashes	33 (53.2)	
Breathlessness	10 (16.1)	
Myalgias	31 (50.0)	
Splenomegaly	37 (59.7)	
Headache	31 (50.0)	
Vomiting	29 (46.8)	
Diarrhea	30 (48.4)	
Comorbidities		
COPD	12 (19.4)	
Asthma	5 (8.1)	
CKD	9 (14.5)	
Hypertension	11 (17.7)	
Diabetes	10 (16.1)	
Severity illness scores		
APACHE		9.90
SOFA		6.70





Totally, 62 were diagnosed as dengue based on clinical features, a positive IgM ELISA and exclusion of other diagnoses.

Baseline patient characteristics are summarized in table I. Of 62 patients, mean age 44.19 years, 35 (56.46) were male and 27 (43.54) were female. Incidence of symptoms being, fever (96.8%), bleeding (37.1%), rashes

(53.2%), breathlessness (16.1%), myalgias (50%), splenomegaly (59.7%), headache (50%), vomiting (46.8%), diarrhoea (48.4%). The admission APACHE-II and SOFA mean scores were 9.90 & 6.70 respectively.

TABLE II. Laboratory investigations

Variable	Mean
Haemoglobin	14.11 gm/dl
Total leucocyte count	7400.80/mm ³
Platelet count	0.36 lakh/mm ³
Serum Creatinine	1.14 mg/dl
Serum Sodium	134.58 meq/l
Serum Bicarbonate	22.45 meq/l
Serum Lactate	1.90 mmol/l
Total Bilirubin	1.79 mg/dl
INR	1.60
PTT	20.22 sec
SGOT	300.3 IU
SGPT	161.9 IU

Laboratory analysis was summarized in table II. It showed a mean values of TLC-7400.80 cells/mm³, Platelets – 0.36 lakh/mm³, serum creatinine – 1.14 mg/dl, serum sodium-134.58 meq/l, lactate-1.90 mmol/l, Total Bilirubin – 1.79 mg/dl, INR-1.60, PTT-20.22 sec, SGOT & SGPT were 300.3 IU and 161.9 IU respectively.

TABLE III. Interventions and Outcome

Variable	n (%)	Mean (number of days)
Vasopressor requirement		
Dopamine	8 (12.9)	
Noradrenaline	8 (12.9)	
Vasopressin	2 (3.2)	
Transfusion requirements		
Packed cells	1 (1.6)	
Platelets	50 (80.6)	
Other supports		

Haemodialysis	4 (6.5)
NIV	3 (4.8)
IMV	6 (9.7)
Both	2 (3.2)


Outcome

Primary

Mortality	1 (1.6)
-----------	---------

Secondary

LOS in ICU	4.25
LOS in Hospital	6.54



Data on interventions & outcome during hospitalization was summarized in table III. Of total patients, the vasopressor requirements being Dopamine(12.9%), Noradrenaline(12.9%) & Vasopressin(3.2%). 1.6% received packed red blood cell transfusions whereas 80.6% had platelet transfusions . 4.8% required Noninvasive ventilation whereas 9.7% required Invasive mechanical ventilation and 3.2% required both non-invasive and invasive mechanical ventilation. 6.5% of patients had received haemodialysis.

The mean duration of ICU and hospital stay were 4.25 and 6.54 days respectively. In hospital mortality was 1.6%.

DISCUSSION

Juneja *et al.* in their prospective study done over 2.5 yrs has reported mean age group as 39.6 yrs (Juneja, D. *et al.*, 2011). Chun *et al.* in their prospective study done with 4787 cases has reported the incidence to be Fever (78.3%), anorexia(32.9%), abdominal pain(32.2%) respectively (Chen, C. M. *et al.*, 2016). Ong *et al.* in their retrospective study done with 3187 cases has reported incidence in mortality associated

group to be minor bleeding(42.9%), Diarrhea(28.6%), headache(14.3%), Abdominal pain (57.1%), Nausea and vomiting(57.1%) (Ong, A. *et al.*, 2007). In our study, of 62 patients, mean age is 44.19 years , 35(56.46) were male and 27 (43.54) were female. Incidence of symptoms being, fever (96.8%), bleeding(37.1%), rashes(53.2%), breathless(16.1%), myalgias(50%), splenomegaly(59.7%), headache(50%), vomiting(46.8%), diarrhoea(48.4%) .

In a retrospective single centered study done by Mahesha Padyana *et al.*, in 96 patients .Median APACHE II score at 24 hours among survivors was 6.0 and non-survivors 17.5 (p <0.01). SOFA score at 48 hours among survivors and non survivors was 6 and 20, respectively (p<0.01), also demonstrated high AST, ALT among non survivors. (Padyana, M. *et al.*, 2019). In our study, the admission APACHE-II and SOFA mean scores were 9.90 & 6.70 respectively.

In a prospective study done by Nayak *et al.*, in 150 patients abnormal AST was seen in 97.33% cases and abnormal ALT was seen in 84% cases thrombocytopenia was seen in all cases with DHF and DSS. (Nayak, J. A. D. U. M. A. N. I. *et al.*, 2017) In our

study, the laboratory analysis showed a mean values of TLC-7400.80cells/mm³, Platelets – 0.36 lakh/mm³, serum creatinine –1.14mg/dl, serum sodium-134.58meq/l, lactate-1.90mmol/l, Total Bilirubin – 1.79mg/dl, INR-1.60, PTT-20.22sec, SGOT & SGPT were 300.3 IU and 161.9 IU respectively.

Padyana *et al.* has retrospectively shown that AKI and moderate to severe ARDS is significantly associated with mortality along with shock and lactates (Padyana, M. *et al.*, 2019). Thanachartwet *et al.* in their prospective study, has shown shock as a significant predictor of mortality (Thanachartwet, V. *et al.*, 2016). Chen *et al.* has retrospectively shown that moderate ARDS, shock and multiorgan failure has significant correlation with mortality. (10) In our study, the vasopressor requirements being Dopamine(12.9%), Nor adrenaline(12.9%) & Vasopressin(3.2%). 1.6% received packed red blood cell transfusions whereas 80.6% had platelet transfusions. 4.8% required Non invasive ventilation whereas 9.7% required Invasive mechanical ventilation and 3.2% required both non invasive and invasive mechanical ventilation. 6.5% of patients had received haemodialysis.

Pinto *et al.* has retrospectively reported the mortality to be 3.1% (Pinto, R. C. *et al.*, 2016). Ong *et al.* in their retrospective case-control study has reported the case fatality rate to be 5.4%. (6) In our study, mean duration of ICU and hospital stay were 4.25 and 6.54 days respectively. In hospital mortality was 1.6%.

CONCLUSION

In our study we hereby conclude the incidence of clinical features is in agreement with other studies whereas the in hospital mortality was relatively low.

REFERENCES

1. Murray, N. E. A., Quam, M. B., & Wilder-Smith, A. (2013). Epidemiology of dengue: past, present and future prospects. *Clinical epidemiology*, 5, 299.
2. Guzman, M.G. & Harris, E. (2015). Dengue. *The Lancet*. Jan 31; 385(9966):453-65.
3. Anand, A. M., Sistla, S., Dhodapkar, R., Hamide, A., Biswal, N., & Srinivasan, B. (2016). Evaluation of NS1 antigen detection for early diagnosis of dengue in a tertiary hospital in Southern India. *Journal of clinical and diagnostic research: JCDR*, 10(4), DC01.
4. Juneja, D., Nasa, P., Singh, O., Javeri, Y., Uniyal, B., & Dang, R. (2011). Clinical profile, intensive care unit course, and outcome of patients admitted in intensive care unit with dengue. *Journal of critical care*, 26(5), 449-452.
5. Chen, C. M., Chan, K. S., Yu, W. L., Cheng, K. C., Chao, H. C., Yeh, C. Y., & Lai, C. C. (2016). The outcomes of patients with severe dengue admitted to intensive care units. *Medicine*, 95(31).
6. Ong, A., Sandar, M., Chen, M. I., & Sin, L. Y. (2007). Fatal dengue hemorrhagic fever in adults during a dengue epidemic in Singapore. *International journal of infectious diseases*, 11(3), 263-267.
7. Padyana, M., Karanth, S., Vaidya, S., & Gopaldas, J. A. (2019). Clinical Profile and Outcome of Dengue Fever in Multidisciplinary Intensive Care Unit of a Tertiary Level Hospital in India. *Indian Journal of Critical Care Medicine: Peer-reviewed, Official Publication of Indian Society of Critical Care Medicine*, 23(6), 270.
8. Nayak, J. A. D. U. M. A. N. I., Behera, S. U. D. E. S. H. N. A., Swain, S. K., & Panda, S. R. (2017). A study of multiorgan dysfunction in patients with dengue and its clinicohematological correlation with severity. *Asian J Pharm Clin Res*, 10(2), 218-221.
9. Thanachartwet, V., Desakorn, V., Sahassananda, D., Jittmittraphap, A., Oer-areemitr, N., Osothsomboon, S., ... & Wattanatham, A. (2016). Serum procalcitonin and peripheral venous lactate for predicting dengue shock and/or organ failure: a prospective observational study. *PLoS neglected tropical diseases*, 10(8).
10. Chen, C. M., Chan, K. S., Yu, W. L., Cheng, K. C., Chao, H. C., Yeh, C. Y., & Lai, C. C. (2016). The outcomes of patients with severe dengue admitted to intensive care units. *Medicine*, 95(31).
11. Pinto, R. C., de Castro, D. B., de Albuquerque, B. C., de Souza Sampaio, V., dos Passos, R. A., da Costa, C. F., ... & Braga, J. U. (2016). Mortality predictors in patients with severe dengue in the State of Amazonas, Brazil. *PloS one*, 11(8).