

Original Research Article

Nephrotoxicity Induced By Copper Oxide and Zinc Oxide Nanoparticles and Their Mixtures in Male Albino Rats

Rabia A. M. Yahya¹, Azab Elsayed Azab^{2*} and Karema El.M.Shkal¹

¹Pharmacology Department, Faculty of Medicine, Sabratha University, Libya

²Physiology Departments, Faculty of Medicine, Sabratha University, Libya

*Corresponding Author

Azab Elsayed Azab

Abstract: Background: Nanotechnology is a fast growing field that provides for the development of materials that have new dimensions, novel properties, and a broader array of applications. It is now known that the toxic behavior of NPs differ from their bulk counterparts. Even NPs that have the same chemical composition differ in their toxicological properties; the differences in toxicity depend upon size, shape, and surface covering. Hence, before NPs are commercially used it is most important that they be subjected to appropriate toxicity evaluation. **Objectives:** The aim of the present study was to evaluate the nephrotoxicity induced by CuO and/or Zn-O nano- particles in male rats. **Materials and Methods:** Twenty adult male rats were grouped randomly into four groups (n=5 each group). Group I (control): Rats were injected with saline intraperitoneally and at a dose of 1.0 ml/kg b.w. for 28 days. Group II (ZnONPs): Rats were administered orally with ZnONPs (10 mg/kg/day) for 28 days. Group III (CuONPs): Rats were injected with CuONPs (0.5 mg/kg/day, in saline; intraperitoneally) for 28 days. Group IV (ZnONPs + CuONPs): Rats were given orally ZnONPs (10 mg/kg/day) followed by injected with CuONPs (0.5 mg/kg/day, in saline; intraperitoneally), for 28 days. At the end of the experimental period, rats were anesthetized using light ether. Blood and kidney tissue samples were taken and prepared for biochemical and histological measurements. **Results:** Serum urea concentration was elevated due treatment of rats with CuONPs, ZnONPs and their mixture compared to control group. Also, CuONPs increased serum uric acid, while ZnONPs reduced serum uric acid and increased creatinine concentration compared to the control group. Treatments of rats with nanoparticles mixture elevated uric acid and creatinine concentration. histopathological abnormalities were observed in kidney of rats treated with CuO-NPs, ZnO-NPs, and CuO-NPs + ZnO-NPs. CuO-NPs, ZnO-NPs, and CuO-NPs + ZnO-NPs –induced histopathological changes in the kidney. The control group showed normal structure of renal glomeruli, Bowman’s capsule lined by squamous epithelium, distinct urinary space, and normal renal tubules. CuO NPs treated group showed degenerated glomerulus, degenerated, desquamated and necrotic renal tubules, and inflammatory cells in the interstitium. More severe similar changes were observed in the ZnO-NPs, and CuO-NPs + ZnO-NPs–treated groups besides wide, congested blood vessels in ZnO-NPs–treated group. **Conclusion:** It can be concluded that CuONPs, ZnONPs, and their mixture were induced biochemical and morphological changes in renal tissues. Also, these results demonstrate that metal oxide nanoparticles induce a range of biological responses that vary from cytotoxic and can only be properly understood by using a tiered test strategy to study other aspects of nanoparticle toxicity. Toxicological studies must be performed before nano-particles application specially nano-oxide nano-particles. Caution should be taken in nano-particles use in work place, preparations as well as while handling.

Keywords: Copper oxide nanoparticles, Zinc oxide nanoparticles, CuO & ZnO mixture, NPs, Nephrotoxicity Renal Histopathology.

INTRODUCTION

The manufacture and use of metal oxide nanoparticles is continuously expanding. It becomes increasingly important to investigate and identify their possible toxicological effects and to identify which

particles pose the greatest harm to human health (Fahmy, B., & Cormier, S. A. 2009).

Copper oxide nanoparticles (CuO NPs) are of great interest in nanoscience and nanotechnology

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easims/>

Article History

Received: 02.12.2019

Accepted: 16.12.2019

Published: 27.12.2019

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

because of their broad industrial and commercial applications. Therefore, toxicity of CuO NPs needs to be thoroughly understood. Akhtar *et al.*, (2012) investigated the cytotoxicity and oxidative stress induced by CuO NPs in human lung epithelial cells.

The manufacture and use of nanoparticles are increasing, humans are more likely to be exposed occupationally or via consumer products and the environment. Copper oxide nanoparticles (CuO NPs) are of great interest in nanoscience and nanotechnology because of their broad industrial and commercial applications. Therefore, toxicity of CuO NPs needs to be thoroughly understood (Karlsson, H. L. *et al.*, 2008). Sizova *et al.*, (2011) reported that despite an increasing application of copper nanoparticles, there is a serious lack of information concerning their impact on human health and the environment. In this study, Serum biochemical analysis and histopathological examinations of the kidney of all the rats were simultaneously performed. All the results indicated that the effects produced by nano-copper at a dose of 100 or 50 mg/kg/d were less than those induced at a higher dose of 200 mg/kg/d. Nano-copper induced overt nephrotoxicity at 200 mg/kg/d for 5 d, which mainly involved widespread renal proximal tubule necrosis. Valko *et al.*, (2005) (Valko, M. M. H. C. M. *et al.*, 2005) reported that metal-induced toxicity and carcinogenicity, with an emphasis on the generation and role of reactive oxygen and nitrogen species, is reviewed. Metal-mediated formation of free radicals causes various modifications to DNA bases, enhanced lipid peroxidation, and altered calcium and sulfhydryl homeostasis. Oxidative stress may be the cause of the cytotoxicity of CuO NPs in mammalian cells. However, little is known about the genotoxicity of CuO NPs following exposure to human cells. The role of reactive oxygen species, with the subsequent oxidative deterioration of biological macromolecules in the toxicities associated with transition metal ions is reviewed. Studies have shown that metals, including iron, copper, chromium, and vanadium undergo redox cycling, while cadmium, mercury, and nickel, as well as lead, deplete glutathione and protein-bound sulfhydryl groups, resulting in the production of reactive oxygen species as superoxide ion, hydrogen peroxide, and hydroxyl radical. As a consequence, enhanced lipid peroxidation DNA damage (Stohs, S. J., & Bagchi, D. 1995).

Syama *et al.*, (2014) reported that engineered nanoparticles are developed for various applications in industrial, electrical, agricultural, pharmaceutical and medical fields due to their unique properties. Nanoparticles such as ZnO are widely used in cosmetics for UV protection. The toxicological investigations of ZnO NPs are highly recommended because of the increasing use in various industrial and consumer products. The toxic potential of ZnO NPs was assumed to be caused by the release of free Zinc ions in the

medium. Zhang *et al.*, (Zhang, H. *et al.*, 2012) reported that CuO and ZnO generated oxidative stress and acute pulmonary inflammation that is not predicted by E(c) levels, the adverse biological effects of these materials could be explained by their solubility, as demonstrated by ICP-MS analysis. This predictive toxicological paradigm is also of considerable importance for regulatory decision-making about this important class of engineered nanomaterials.

Although nanozinc oxide (nano-ZnO) is applied widely in photocatalysts and gas sensors and in biological fields, it can cause serious oxidative stress and DNA damage to mammalian cells. Along with existing and emerging use of nanoscale materials, growing concerns have arisen about their unintentional health and environmental impact.

OBJECTIVES

Presently, there is only limited knowledge concerning the toxicological effects of NPs. However, it is now known that the toxic behavior of NPs differ from their bulk counterparts. Even NPs that have the same chemical composition differ in their toxicological properties; the differences in toxicity depend upon size, shape, and surface covering. Hence, before NPs are commercially used it is most important that they be subjected to appropriate toxicity evaluation. The aim of the present study was to evaluate the nephrotoxicity induced by nano-CuO and/or Zn-O in male rats.

MATERIAL AND METHODS

Chemicals

Copper oxide and Zinc oxide as nanoparticles with an average size of 6 and 51 nm, respectively, were a gift from Dr. Amina El-Trass. Synthesis, characterization, optical properties and interaction with amino acids of CuO nanoparticles to confirm the negative surface of CuO nanoparticles were performed by El-Trass *et al.*, (2012).

Animals and Housing

Twenty healthy male Wistar Albino rats weighing 150 ± 10 g, were obtained from the Animal Breeding House of the National Research Centre (NRC), Dokki, Cairo, Egypt. The rats were allowed to acclimatize for a week before starting the experiments. Rats were maintained under temperature-controlled conditions (25 °C), and a normal photoperiod of 12 h of darkness and 12 h of light. They were fed with standard food and had free access to water. Animals were randomly divided into 4 groups of five rats each, with one group assigned to be an untreated control. The housing and management of the animals and the experimental protocols were conducted as stipulated in the Guide for Care and Use of Laboratory Animals (NRC. 1996).

Experimental Protocol

Twenty adult male rats were grouped randomly into four groups (n=5 each group). Group I (control): Rats were injected with saline intraperitoneally and at a dose of 1.0 ml/kg b.w. for 28 days. Group II (ZnONPs): Rats were administered orally with ZnONPs (10 mg/kg/day) for 28 days. Group III (CuONPs): Rats were injected with CuONPs (0.5 mg/kg/day, in saline; intraperitoneally) for 28 days (Liao, M., & Liu, H. 2012). Group IV (ZnONPs + CuONPs): Rats were given orally ZnONPs (10 mg/kg/day) followed by CuONPs (0.5 mg/kg/day, in saline; intraperitoneally), for 28 days.

At the end of the experimental period, rats were anesthetized using light ether. Blood samples were taken from the vena cava of rat heart within 1 min after sacrifice. Tubes were used to compile blood drawn from the heart directly; the blood was collected in glass tubes for coagulation and serum formation, blood was allowed to set for 30 min at 4°C to clot, then centrifuged for 5 minutes at 1000 x g. Packed cells were discarded and the supernatant serum samples were decanted and stored into capped sterile poly-ethylene tubes at -20°C until used (within 24 hours). The abdominal cavity of each rat was opened where the liver was excised.

BIOCHEMICAL PARAMETERS IN RAT SERUM AND TISSUES

Determination of Urea

Urea was determined using a commercially available kit (Urea-kit S 180, bioMerieux Vitek, Inc. USA). Concentration of urea in the sample was calculated which is equal = $\Delta A_{\text{sample}} / \Delta A_{\text{standard}} \times \text{concentration of the standard}$ which is known and equal 50 mg/dl (Patton, C.J., & Crouch, S. R. 1977).

Determination of Uric Acid

Uric acid was determined using a commercially available kit (Uric acid-kit, Cat. No. 10690. Human Gesellschaft, Taunusstein, Germany). Concentration of uric acid in the sample was calculated which equal = $\Delta A_{\text{sample}} / \Delta A_{\text{standard}} \times \text{concentration of the standard}$ which is known and equal 8 mg/dl (Guder, W. G. *et al.*, 1996).

Determination of Creatinine

Creatinine was determined using a commercially available kit (CREA-kit MPR3 124192 Boehringer Mannheim). The concentration of creatinine in the sample was calculated which is equal = $\Delta A_{\text{sample}} / \Delta A_{\text{standard}} \times \text{concentration of the standard}$ which is known and equal 2 mg/dl (Henry, R.J. 1974).

Histopathological Analysis

Dissected kidneys were immediately fixed into 10% formaldehyde saline. Tissues were processed by embedding in paraffin. Sections were cut by rotatory microtome and mounted on glass slides. The sections were stained by conventional Hematoxylin & Eosin

(H&E) stain. The sections were examined by light microscope (Drury, R., & Wallington, E. 1980).

Statistical Analysis

Values obtained as mean \pm SEM were subjected to one-way analysis of variance (ANOVA) followed by Tukey test using GraphPad Prism version 4.0 for windows from GraphPad Software, San Diego, California, USA). Values of $P < .05$ were considered significant.

RESULTS AND DISCUSSION

Nanoparticles (NPs) were found to reach the systemic circulation after inhalation, ingestion or intravenous injection. They are known to disseminate to several organs such as liver, spleen, kidneys, brain or heart (Kreuter, J. 2001; Girardin, F. 2006; Oberdörster, G. *et al.*, 2005; Elder, A. *et al.*, 2006). Such translocation depends on the physicochemical properties of NPs, and their migration to distant sites is an important issue with regard to their toxicity.

Effects of Copper Oxide Nanoparticles and Zinc Oxide Nano-Particles on Rats Serum Urea, Uric Acid, and Creatinine

Over a third of the atoms in a nanoparticle are at the surface, and these are extremely reactive systems, which in some cases can generate oxygen radicals; because of the size of nanostructures, it is possible to manipulate the surface interface to allow for interactions with biological systems (Guyton, A.C., & Hall, J.E. 2001). With the correct coating, particles below 50 nm can translocate them into cells relatively easily and are able to interact with channels, enzymes, and other cellular proteins (Zhang, X. D. *et al.*, 2010). This may be the mechanism that CuNPs induced its toxicity to hematological indices. This study aimed to investigate human renal cell responses to manufactured NPs in order to highlight their potential toxicity and/or biological responses. Kidneys play an important role in eliminating xenobiotics from the body and thus, NPs absorbed in the systemic circulation can be excreted by renal clearance (Burns, A. A. *et al.*, 2008; Schipper, M. L. *et al.*, 2009).

An abnormally elevated blood creatinine level is a specific and sensitive indicator of impaired kidney function. The urea, uric acid and blood urea nitrogen values in the CuONPs group were significantly higher compared to control ($P < 0.05$) groups, while creatinine without significant changes. ZnO NPs induced serum urea and creatinine, while reduced uric acid compared to control. On the other hand the mixture of CuO and ZnO NPs induced significant ($P > 0.05$) changes in urea, uric acid, and creatinine (Tables 1 & Figures 1-3). The exposure to ZnO NPs significantly affected cellular viability in a dose-dependent manner and formation of reactive oxygen species (ROS) was found to be the mechanism of cellular toxicity (Syama, S. *et al.*, 2014).

In this study, the biochemical compositions of serum of rats treated with nano-copper and zinc oxide indicating impairment of kidney function. Also, Lei *et al.*, (2008) found that 200 mg/kg/d of nano-copper induced nephrotoxicity reflected with renal proximal tubule necrosis and elevation in creatine levels in

serum. In addition, Liao and Liu (2012) and Sizova *et al.*, (2011) found that nanocopper can induce widespread renal proximal tubule necrosis in rat kidneys with blood urea nitrogen and creatinine increase.

Table (1). Effects of treatment of rats with zinc oxide (ZnONPs), and copper oxide nanoparticles (CuO NPs), and their mixture on serum urea, uric acid, and creatinine

Parameters	Groups			
	Control	CuONP	ZnONP	CuO + ZnONP
Serum urea (mg/dl)	26.80± 4.749 ^{bcd}	43.0± 3.52 ^a	40.6± 5.28 ^a	42.2± 3.82 ^a
Serum uric acid (mg/dl)	2.50 ± 0.08 ^{cd}	3.11 ± 0.28 ^{ac}	2.27 ± 0.15 ^{abd}	3.79 ± 0.69 ^{ac}
Serum creatinine (mg/dl)	0.95± 0.02 ^{bcd}	0.92± 0.06 ^a	1.01± 0.05 ^a	1.03± 0.05 ^a

Significance at $P < 0.05$. ^a Comparison of control and other groups; ^b Comparison of CuONP and other groups;

^c Comparison of ZnONP and other groups; ^d Comparison of CuO+ZnONP and other groups




Figure 1: Serum urea (mg/dl) of rat treated with zinc oxide (ZnO) and copper oxide (CuO) nanoparticles (NPs). Significance at $P > 0.05$. a Comparison of control and other groups; b Comparison of CuO NPs and other groups; c Comparison of ZnO NPs and other groups; d Comparison of CuO+ZnONPs and other groups




Figure 2: Serum uric acid (mg/dl) of rat treated with zinc oxide (ZnO) and copper oxide (CuO) nanoparticles (NPs). Significance at $P > 0.05$. a Comparison of control and other groups; b Comparison of CuO NPs and other groups; c Comparison of ZnO NPs and other groups; d Comparison of CuO+ZnONPs and other groups




Figure 3: Serum creatinine (mg/dl) of rat treated with zinc oxide (ZnO) and copper oxide (CuO) nanoparticles (NPs). Significance at $P > 0.05$. Significance at $P > 0.05$. a Comparison of control and other groups; b Comparison of CuO NPs and other groups; c Comparison of ZnO NPs and other groups; d Comparison of CuO+ZnONPs and other groups

Effects of Copper Oxide Nanoparticles (CuO Nps) and Zinc Oxide Nano-Particles (Zno Nps) On Rats Kidney Histology

Despite an increasing application of copper and zinc nanoparticles, there is a serious lack of information concerning their impact on human health and the environment. Elevation in serum levels of urea could also be attributed to increase in the activities of urea enzymes, ornithine carbomoyl transferase and arginase mostly associated in liver damage in many animal species, since the urea cycle is confined to the liver (James, W. *et al.*, 1998). Damage to the kidney may result in reduced erythropoietin production, resulting in high urea which may in turn be associated with low blood volume. Thereby leading to an elevation in inflammatory cells types, this usually occurs during the inflammatory process. Inflammation exposes the body organs to infections, leading to the release of high amount of white blood cells (James, W. *et al.*, 1998).

Microscopic examination of organs to evaluate the toxic effects of contaminants has been used as a biomarker to evaluate the toxicity of various pollutants (Adams, S. 2002; Mela, M. *et al.*, 2007). Histopathology studies of target organs along with the studies of oxidative stress would give the complete risk assessment and toxic potential of copper and zinc oxide nanoparticles (CuO NPs & ZnO-NPs). In the present study, histopathological abnormalities were observed in kidney of rats treated with CuO-NPs, ZnO-NPs, and CuO-NPs + ZnO-NPs.

Figures (4-7) demonstrate CuO-NPs, ZnO-NPs, and CuO-NPs + ZnO-NPs –induced histopathological changes in the kidney. The control group showed normal structure of renal glomeruli, Bowman’s capsule lined by squamous epithelium, distinct urinary space, and normal renal tubules. CuO NPs treated group showed degenerated glomerulus, degenerated, desquamated and necrotic renal tubules, and inflammatory cells in the interstitium. More severe similar changes were observed in the ZnO-NPs, and

CuO-NPs + ZnO-NPs–treated groups besides wide, congested blood vessels in ZnO-NPs–treated group.

Kidney of the different treated groups showed degeneration of tubular epithelial cells and glomerular degeneration. This may be due to the accumulation of inflammatory cells (Kurtović, B. *et al.*, 2008). Venkatramreddy and Baul, (2010), Aslam *et al.*, (2013), and Patel *et al.*, (2014) reported that lipid peroxidation is the primary source for tubular dilation, and disintegration tubules. Necrosis is a form of cell injury that results in the premature death of cells in living tissue by autolysis (Patel, J. G. *et al.*, 2014). Cells that die due to necrosis exhibit loss of cell membrane integrity and an uncontrolled release of products of cell death into the intracellular space (Proskuryakov, S. Y. *et al.*, 2003). This initiates, in the surrounding tissue, an inflammatory response which prevents nearby phagocytes from locating and eliminating the dead cells by phagocytosis (Kasper, D. L., & Zaleznik, D. F. 2001). The histopathological abnormalities observed in the present study, in kidney of rats treated with CuO-NPs, ZnO-NPs, and CuO-NPs + ZnO-NPs could be explained on basis of the reported increase in lipid peroxidation, and decreased level of reduced glutathione.




Figure 4: Photomicrograph of section of kidney of control group showing normal structure of renal glomeruli (1), Bowman’s capsule lined by squamous epithelium (2), distinct urinary space (3), and normal renal tubules (4). H & E stain x 400.




Figure 5: Photomicrograph of section of kidney of CuO NPs treated group showing degenerated glomerulus (1), degenerated, desquamated and necrotic renal tubules (2), and inflammatory cells in the interstitium (3) (H&E X 400).




Figure 6: Photomicrograph of section of kidney of ZnO NPs- treated group showing degenerated glomerulus (1), degenerated, atrophied, desquamated and necrotic renal tubules (2), inflammatory cells in the interstitium (3), and wide, congested blood vessels (4) (H&E X 400).




Figure 7: Photomicrograph of section of kidney of ZnO + CuO NPs treated group showing degenerated glomerulus (1), degenerated, atrophied, desquamated and necrotic renal tubules (2), and inflammatory cells in the interstitium (3) (H&E X 400).

Furthermore, these histological changes were in line with the reported changes in the kidney function parameters (serum creatinine, urea, and uric acid). The negative surface of CuO nanoparticles reported by El-

Trass *et al.*, (2012) indicates its ability to generate free radicals, and its ability to induce oxidative stress as reported by Karlsson *et al.*, (2009), Song *et al.*, (2013). As a result of this it could play an important role in the peroxidation of unsaturated fatty acids in the cell membranes and development of necrosis as suggested by Lloyd *et al.*, (1997; 1998), and Susa *et al.*, (1996). The redox alterations caused by oxidative agents have been shown to induce apoptosis and necrosis in hepatocytes and other cells (Han, D. *et al.*, 2006).

CONCLUSION

From the previous results, it could be concluded the main following points: The results obtained showed that CuONPs, ZnONPs and their mixture increased serum kidney function tests and produce morphological alterations in the kidney tissues. These results demonstrate that metal oxide nanoparticles induce a range of biological responses that vary from cytotoxic and can only be properly understood by using a tiered test strategy such as we developed for oxidative stress and adapted to study other aspects of nanoparticle toxicity. Toxicological studies must be performed before nano-particles application specially nano-oxide nano-particles. Caution should be taken in nano-particles use in work place, preparations as well as while handling.

REFERENCES

1. Fahmy, B., & Cormier, S. A. (2009). Copper oxide nanoparticles induce oxidative stress and cytotoxicity in airway epithelial cells. *Toxicology In Vitro*, 23(7), 1365-1371.
2. Akhtar, M. J., Ahamed, M., Fareed, M., Alrokayan, S. A., & Kumar, S. (2012). Protective effect of sulphoraphane against oxidative stress mediated toxicity induced by CuO nanoparticles in mouse embryonic fibroblasts BALB 3T3. *The Journal of toxicological sciences*, 37(1), 139-148.
3. Karlsson, H. L., Cronholm, P., Gustafsson, J., & Moller, L. (2008). Copper oxide nanoparticles are highly toxic: a comparison between metal oxide nanoparticles and carbon nanotubes. *Chemical research in toxicology*, 21(9), 1726-1732.
4. Sizova, E., Glushchenko, N., Miroshnikov, S., & Skalny, A. (2011). Influence of Cu10x copper nanoparticles intramuscular injection on mineral composition of rat spleen. *Journal of Trace Elements in Medicine and Biology*, 25, S84-S89.
5. Valko, M. M. H. C. M., Morris, H., & Cronin, M. T. D. (2005). Metals, toxicity and oxidative stress. *Current medicinal chemistry*, 12(10), 1161-1208.
6. Stohs, S. J., & Bagchi, D. (1995). Oxidative mechanisms in the toxicity of metal ions. *Free radical biology and medicine*, 18(2), 321-336.

7. Syama, S., Srekanth, P. J., Varma, H. K., & Mohanan, P. V. (2014). Zinc oxide nanoparticles induced oxidative stress in mouse bone marrow mesenchymal stem cells. *Toxicology mechanisms and methods*, 24(9), 644-653.
8. Zhang, H., Ji, Z., Xia, T., Meng, H., Low-Kam, C., Liu, R., ... & Wang, M. (2012). Use of metal oxide nanoparticle band gap to develop a predictive paradigm for oxidative stress and acute pulmonary inflammation. *ACS nano*, 6(5), 4349-4368.
9. El-Trass, A., ElShamy, H., El-Mehasseb, I., & El-Kemary, M. (2012). CuO nanoparticles: synthesis, characterization, optical properties and interaction with amino acids. *Applied Surface Science*, 258(7), 2997-3001.
10. NRC. (1996). Seventh Revised Edition. Subcommittee on Beef Cattle Nutrition. Committee on Animal Nutrition. *Board on Agriculture. National Research Council*.
11. Liao, M., & Liu, H. (2012). Gene expression profiling of nephrotoxicity from copper nanoparticles in rats after repeated oral administration. *Environmental toxicology and pharmacology*, 34(1), 67-80.
12. Patton, C.J., & Crouch, S. R. (1977). A simple and rapid micro-Kjeldahl method for total nitrogen analysis. *Anal. Chem.*, 49: 464-469.
13. Guder, W. G., Narayanan, S., Wisser, H., & Zawta, B. (1996). List of Analytes; Preanalytical variables. Brochure in: *Samples: From Patient to the Laboratory*.
14. Henry, R.J. (1974). Clinical chemistry. Principles and techniques. 2nd Edition. *Harper and Row, Pp. 525-531*.
15. Drury, R., & Wallington, E. (1980). Carlton, S. Histological Technique, 5 th ed., *Oxford university press, Oxford, New York, Toronto*.
16. Kreuter, J. (2001). Nanoparticulate systems for brain delivery of drugs. *Advanced drug delivery reviews*, 47(1), 65-81.
17. Girardin, F. (2006). Membrane transporter proteins: a challenge for CNS drug development. *Dialogues in clinical neuroscience*, 8(3), 311.
18. Oberdörster, G., Oberdörster, E., & Oberdörster, J. (2005). Nanotoxicology: an emerging discipline evolving from studies of ultrafine particles. *Environmental health perspectives*, 113(7), 823-839.
19. Elder, A., Gelein, R., Silva, V., Feikert, T., Opanashuk, L., Carter, J., ... & Oberdörster, G. (2006). Translocation of inhaled ultrafine manganese oxide particles to the central nervous system. *Environmental health perspectives*, 114(8), 1172-1178.
20. Guyton, A.C., & Hall, J.E. (2001). Textbook of Medical Physiology 10th ed, *New Delhi, pp. 309-310*.
21. Zhang, X. D., Wu, H. Y., Wu, D., Wang, Y. Y., Chang, J. H., Zhai, Z. B., ... & Fan, F. Y. (2010). Toxicologic effects of gold nanoparticles in vivo by different administration routes. *International journal of nanomedicine*, 5, 771.
22. Burns, A. A., Vider, J., Ow, H., Herz, E., Penate-Medina, O., Baumgart, M., ... & Bradbury, M. (2008). Fluorescent silica nanoparticles with efficient urinary excretion for nanomedicine. *Nano letters*, 9(1), 442-448.
23. Schipper, M. L., Iyer, G., Koh, A. L., Cheng, Z., Ebenstein, Y., Aharoni, A., ... & Chen, X. (2009). Particle size, surface coating, and PEGylation influence the biodistribution of quantum dots in living mice. *Small*, 5(1), 126-134.
24. Lei, R., Wu, C., Yang, B., Ma, H., Shi, C., Wang, Q., ... & Liao, M. (2008). Integrated metabolomic analysis of the nano-sized copper particle-induced hepatotoxicity and nephrotoxicity in rats: A rapid in vivo screening method for nanotoxicity. *Toxicology and applied pharmacology*, 232(2), 292-301.
25. Liao, M., & Liu, H. (2012). Gene expression profiling of nephrotoxicity from copper nanoparticles in rats after repeated oral administration. *Environmental toxicology and pharmacology*, 34(1), 67-80.
26. James, W., Lohr, G., Willsky, R., & Margaret, A. (1998). Renal Drug Metabolism. *Am. Soc. Pharmacol. Exp. Therap.*, 50: 108-110.
27. Adams, S. (2002). Biological indicators of aquatic ecosystem stress. *Amer. Fish. Soc.*, 3: 104-112.
28. Mela, M., Randi, M. A. F., Ventura, D. F., Carvalho, C. E. V., Pelletier, E., & Ribeiro, C. O. (2007). Effects of dietary methylmercury on liver and kidney histology in the neotropical fish *Hoplias malabaricus*. *Ecotoxicology and Environmental Safety*, 68(3), 426-435.
29. Kurtović, B., Teskeredžić, E., & Teskeredžić, Z. (2008). Histological comparison of spleen and kidney tissue from farmed and wild European sea bass.
30. Venkatramreddy, V., & Baul, B. (2010). Chromium induced biochemical, genotoxic and histopathologic effects in liver and kidney of Goldfish, *Carassius auratus*. *Mutat. Res.*, 30: 43-51.
31. Aslam, M., Dayal, R., Javed, K., Parray, S. A., Jetley, S., & Samim, M. (2013). Nephroprotective effects of methanolic extract of *Peucedanum grande* against acute renal failure induced by potassium dichromate in rats. *Int J Pharm Sci Drug Res*, 5(2), 45-9.
32. Patel, J. G., Joshi, D. V., Patel, B. J., & Raval, S. H. (2014). Biochemical and pathomorphological study of potassium dichromate-induced nephrotoxicity in wistar rat. *Journal of Advanced Veterinary Research*, 4(1), 12-17.
33. Proskuryakov, S. Y., Konoplyannikov, A. G., & Gabai, V. L. (2003). Necrosis: a specific form of programmed cell death?. *Experimental cell research*, 283(1), 1-16.

34. Kasper, D. L., & Zaleznik, D. F. (2001). Gas gangrene, antibiotic-associated colitis, and other clostridial infections. *HARRISONS PRINCIPLES OF INTERNAL MEDICINE, 1*, 922-926.
35. El-Trass, A., ElShamy, H., El-Mehasseb, I., & El-Kemary, M. (2012). CuO nanoparticles: synthesis, characterization, optical properties and interaction with amino acids. *Applied Surface Science, 258*(7), 2997-3001.
36. Karlsson, H. L., Gustafsson, J., Cronholm, P., & Möller, L. (2009). Size-dependent toxicity of metal oxide particles—a comparison between nano- and micrometer size. *Toxicology letters, 188*(2), 112-118.
37. Song, M. F., Wang, Y. W., Wang, C., & Peng, K. L. (2013). Study of genotoxicity and oxidative stress induced by copper oxide nanoparticles in mice. *Zhonghua lao dong wei sheng zhi ye bing za zhi= Zhonghua laodong weisheng zhiyebing zazhi= Chinese journal of industrial hygiene and occupational diseases, 31*(9), 690-692.
38. Lloyd, D. R., Phillips, D. H., & Carmichael, P. L. (1997). Generation of putative intrastrand cross-links and strand breaks in DNA by transition metal ion-mediated oxygen radical attack. *Chemical research in toxicology, 10*(4), 393-400.
39. Lloyd, D. R., Carmichael, P. L., & Phillips, D. H. (1998). Comparison of the formation of 8-hydroxy-2'-deoxyguanosine and single- and double-strand breaks in DNA mediated by fenton reactions. *Chemical research in toxicology, 11*(5), 420-427.
40. Susa, N., Ueno, S., Furukawa, Y., & Sugiyama, M. (1996). Protective effect of vitamin E on chromium (VI)-induced cytotoxicity and lipid peroxidation in primary cultures of rat hepatocytes. *Archives of toxicology, 71*(1-2), 20-24.
41. Han, D., Hanawa, N., Saberi, B., & Kaplowitz, N. (2006). Mechanisms of liver injury. III. Role of glutathione redox status in liver injury. *American Journal of Physiology-Gastrointestinal and Liver Physiology, 291*(1), G1-G7.