

## Original Research Article

## Estimation of Cholesterol Content in *Synodontis Budgetti* from Amassoma, Tombia, Swali and Otuokpoti Rivers in Bayelsa State

Ezomoh O. O<sup>1</sup>, Agberia S. O<sup>1</sup>, Ekechukwu B<sup>1</sup> and Tamunoene B<sup>1</sup><sup>1</sup>Department of Biochemistry, Faculty of Basic Medical Sciences, Niger Delta University, Wilberforce Island, Amassoma, Bayelsa State, Nigeria

\*Corresponding Author

Ezomoh, O. O

**Abstract:** The investigation was carried out to estimate cholesterol content in *synodontis budgetti* from amassoma, tombia, swali and Otuokpoti Rivers in bayelsa state using the gas chromatography. The result obtained showed the cholesterol content from Otuokpoti River (5.67±0.17), from Amassoma River (5.34±0.02), from Swali River (5.17±0.02), from Tombia River (5.12±0.02). The cholesterol content in *synodontis budgetti* showed a significant decrease ( $p < 0.05$ ) in Okuokpoti, Amassoma, Swali and Tombia Rivers. All cholesterol content showed significant difference ( $p < 0.05$ ) relative to each other. The differences in cholesterol content of fish samples *synodontis budgetti* from the four rivers may be as a result of age, gender, diet, geographical origin and season of the fish.

**Keywords:** investigation, *synodontis budgetti*, Amassoma River.

### INTRODUCTION

According to the Food and Agriculture Organization (FAO), it is estimated that 3 billion people around the world consume fish and other marine organisms as a source of protein (Enrique Murillo, *et al.*, 2013). Lipids are the predominant source of energy for fish. The mechanisms by which fish allocate energy from lipids for metabolism, development, growth and reproduction are critical for understanding key life-history strategies and transitions. Lipids are the major source of nutrition in marine fishes. They are considered an efficient biochemical means of concentrating large amounts of stored energy at small space. The cells of white fat tissue, called adipocytes, are responsible for lipid synthesis, release and storage in the organism. The genus *Synodontis budgetti* commonly known as the up-side down catfish belongs to the family *Mochokidae* which is made up of three genera, *Mochocus*, *Synodontis* and *Chinoglanis*. It is the largest genus of the catfishes of the order *Siluriformes*, and most widely distributed. The specie, *Synodontis budgetti* of the genus *Synodontis*, occur through most of the freshwaters of sub-saharan Africa and the Nile river system, but are restricted to water system within the tropics. They can be found in large rivers, small fast-flowing streams and massive African rift lakes. *Synodontis*

was one of the most abundant genera in the Nile system. *Synodontis* is the most common genus of the *Mochokidae* family and of great commercial importance in the inland waters of Nigeria and West Africa (Akombo *et al.*, 2016). Fish is of high nutritional value due to its high protein content, excellent quality of lipids (omega 3 and 6 series) and low level of total fat, saturated fats and cholesterol (Magobe, *et al.*, 2015). Surveys aimed at establishing the influence of n-3 fatty acids on various health conditions have shown their positive effects, both in disease prevention and health status improvement. The human brain consists of 60% lipids, of which 33% are n-3 fatty acids. Cholesterol is a waxy fat-like substance that is present in all animals but not in plant. It is not soluble in blood and so has to be carried in the bloodstream by molecules called lipoproteins. Lipoproteins are named based on their density. The high density lipoprotein (HDL) is 'good' and transports cholesterol to the liver for further processing. Low density lipoprotein is 'bad' fat and when oxidized gets deposited in the walls of arteries contributing to arteriosclerosis. The former is so named because it is known to contain less cholesterol and more phospholipids and apoproteins than the later. HDL is present in higher amount in fish lipids, whereas the LDL is more abundant in other animal fats and contains more of the saturated fatty acids (SFA).

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easims/>

Article History

Received: 15.04.2019

Accepted: 30.04.2019

Published: 17.05.2019

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

**MATERIAL AND METHODS**

The sample (3.0g) was weighed into the labeled porous thimble; 200ml of the petroleum ether was measured and then added to a dried 250ml capacity flask. The covered porous thimble with the sample was placed condenser of the Soxhlet extractor. Extraction was done for 5hrs.

The porous thimble was removed with care and the petroleum ether in the top container (tube) was collected for reuse. The extraction flask was removed

from the heating mantle arrangement when it was almost free of petroleum ether. The extraction flask with the oil was oven dried at 105°C for the period of one (1) hour. The flask containing the dried oil was cooled in the desiccator and the weight of the cooled flask with the dried oil was measured using gas chromatography. The results (means ± standard deviation) were calculated as both the weight percentage (fatty acid profile) and the concentration (mg/g dry tissues).

**RESULT**

Showing Cholesterol content of fish samples *Synodontis budgetti* from Otuokpoti, Amassoma, Swali and Tombia River.

| RIVERS                      | CHOLESTEROL                                     |
|-----------------------------|---|
| OTUOKPOTI RIVER mg/100g(DW) | <i>Synodontis budgetti</i> Mean±SD<br>5.67±0.17 |
| AMASSOMA RIVER mg/100g (DW) | <i>Synodontis budgetti</i> Mean±SD<br>5.34±0.02 |
| SWALI RIVER mg/100g (DW)    | <i>Synodontis budgetti</i> Mean±SD<br>5.17±0.02 |
| TOMBIA RIVER mg/100g (DW)   | <i>Synodontis budgetti</i> Mean±SD<br>5.12±0.02 |

Values are mean of three determination ± SD. S

The mean (±SD) values of total fat and Cholesterol content of fish samples *Synodontis budgetti* determined from the four different rivers; Otuokpoti (5.67±0.17), Amassoma (5.34±0.02), Swali (5.17±0.02) and Tombia Rivers (5.12±0.02) Bayelsa State are significantly different (p<0.05).

All the cholesterol content in *Synodontis budgetti* showed a significant decrease (p<0.05) in Otuokpoti, Amassoma, Swali, and Tombia Rivers. All cholesterol showed significant difference (p<0.05) relative to each other. The differences in Cholesterol content of fish samples *Synodontis budgetti* from the four rivers may be as a result of age, gender, diet, Geographical origin and season of the fish.

**DISCUSSION**

The result obtained from the Determination showed a significant variation in the lipid profile of *Synodontis Budgetti* found in Amassoma, Tombia, Swali and Otuokpoti River respectively. The fish species *Synodontis Budgetti* studied belong to High fat fish (> 8%) category, going by the classification of Ackman. The lower side of the lipid range (5.12±0.02) is in line (5.12±0.02) with some literature report. Apart from the issues of seasonal variations between dry and rainy seasons which are known to affect fat content of fishes, spawning activities can use-up the fat reserves and give very low fat levels. Other causes of variation in fat content are (as already mentioned) attributed to geographical differences and particularly the feed-type. Of the total cholesterol determined, HDL values were

higher than their corresponding LDL levels and of which was also in consonant with literature report. However it was observed that the presence of trans fatty acids could have a lowering effect on the body's HDL cholesterol - an effect which underscores the need to keep the level of the former low (not to rise above % of total energy intake), if the HDL level should be maintained. The levels of PUFA determined from fishes in OR, AR & SR were similar; but were all higher than that from TR'. The specie *Synodontis Budgetti* from Otuokpoti river was the major contributor of PUFA in each study location, thereby confirming the fact that location differences is one of the major factors in nutrient accumulation. The present study therefore supports the production of aquaculture of C G. Prominent PUFAs in fishes in OR were C20:4, C20:5 and C22:6; while those of AR included, together with the above acids, the C18:2. Fishes from SR and TR featured mostly the C18:3, C20:3; and C18:2, C20:3, C20:4, C22:2, C20:5, C:22:6 acids respectively, and agreed with literature reports. Still among the PUFAs, C18:2 (linoleic) was prominent in fishes from TR, AR & LR. The C18:3(linolenic) was more prominent in SR fishes. Both acids are vital for growth and development and therefore must be provided in the diet since the body cannot synthesize them. With them available, the body can synthesize arachidonic and other eicosanoids. The variations recorded in the concentration of nutritional components in the fish examined could have been as a result of the rate in which these components are available in the water body and the ability of the fish to absorb and convert the essential nutrients from the diet or the water bodies where they live (Yeannes and Almandos, 2003).

In addition, since the chemical composition of synodontis was found to vary with sex, seasons, size, age and locality of catch, therefore, it is essential to determine and evaluate for different species of the fish in relation to these factors.

(Otitologbon, *et al.*, 1997) reported similar findings and attributed this to the dominance of the lipid in the water body where the fish lives.

(Effiong and Mohammed 2008) reported that nutrient content of fish is influenced by several factors such as feeding, processing method, habitat etc.

Lipid content of *Synodontis Budgetti* from Otuokpoti River increased significantly than to Amassoma, Swali and Tombia River as these may be as a result of the population level. The high level of this lipid in *Synodontis Budgetti* may be attributed to the rate in which they are available in the water body and the ability of the fish to absorb it from their diets and the environment where they live.

## CONCLUSION

The fishes studied belong to High Fat Fish category. The fat accumulation was related to species type, nature of feed and hence geographical location. Cholesterol was the prominent fat followed by HDL and LDL in that order. Lower levels were determined from the study areas in Swali and Tombia than in Otuokpoti and Amassoma. The fish, *Synodontis Budgetti* from otuokpoti seemed more nutritionally promising in terms of essential fatty acid (EFA) content. It is therefore recommended for culture production as it contained reasonable levels of EFA and showed higher values of total PUFA when compared to the other fishes from other the habitats studied. The present study has therefore provided good sources of fish oils rich in essential polyunsaturated fatty acids particularly EPA, DHA and the n- 6 PUFA - eicosatetraenoic acid (arachidonic acid). Increased fish production in Otuokpoti and Amassoma areas in Nigeria is recommended so as to provide more EFAs needed for good health.

## REFERENCES

- Achionye-Nzeh, C. G., Adedoyin O. M., Oyebanji, S., & Mohammed, M. O. (2011). Mineral composition of some marine and freshwater fishes agriculture and biology. *Journal of North America Agriculture*, 2(7), 1113-1116.
- Adeyeye, E. I., Oyerekua, M. A., & Adesina, A. J. (2014). Proximate, mineral, amino acid composition and mineral safety index of callinectes latimanus. *International Journal of Development Research*, 1(1), 333-336.
- Adewoye, S. O., Fawole, O. O., & Omotosho, J. S. (2003). Concentrations of selected elements in some freshwater fishes in Nigeria. *Science Focus*, 4(2), 106-108.
- Alas, A., Ozcan, M, M., & Harmankaya, M. (2014). Mineral contents of head, caudal, central fleshy part and spinal columns of some fishes. *Environmental Monitoring and assessment*, 18(6), 889-894.
- Asabga, S. O. (2000). Effect of cadmium on kidney and liver cell membrane integrity and antioxidant enzyme status: implications for Warri River cadmium level. *Tropical Journal of Environment Science Health*, 3(1), 33-39.
- Batra, J., & Seth, P. K. (2002). Effect of cadmium of iron deficiency on developing rat brain. *Indian Journal of Biochemistry*, 17(2), 108-114.
- Beyza, E., & Akif, O. (2009). The effect of cooking methods on mineral and vitamin contents of African catfish. *Food Chemistry*, 11(5), 419-422.
- Beyza, E., & Akif, O. (2009). The effect of cooking methods on mineral and vitamin contents of Africa catfish. *Food Chemistry*, 11(5), 419-422.
- Cerklewski, F. L. (1998). Fluoride: Essential or just beneficial. *Nutritional Review*. 1(4):475-476.
- Elagba, M. H. A., Rabie, A., & Mohamed, H. (2010). Proximate composition, amino acid and mineral contents of five commercial Nile fishes in Sudan. *African Journal of Food Science*. 4(10):650-654.
- Effiong, B. N. and Mohamed, I. (2008). Effect of seasonal variation on the nutrient composition in selected fish species in Lake Kainji, Nigeria. *Nature and Science*, 6(2), 1-5
- Fakwole, O. O., Ogundiran, M. A., Ayandiran, T. A. and Olagunju, O. F. (2007). Mineral composition in some selected fresh water fishes in nigeria. *Journal of Food Safety*, 9(4), 52-55.
- Fakunle, J. O., Effiong, B. N., & Ibanga, U. I. (2009). Nutrient composition of three commercially important fisheries in Laje Kainji Nigeria. *Proceedings of the 24<sup>th</sup> Annual Conference of Fisheries Society of Nigeria, Akure*, 10(1), 46-48.
- Saadettin, G., Barbaros, D., Niger, A., Ahmet, C., & Mehmet, T. (1999). Proximate composition and selected mineral content of commercial fish species from the Black Sea. *Journal of the Science of Food and Agriculture*, 5(5), 110-116.
- Otitologbon, S. A., Agbaji, E. B., Peter, O. A., & Oniye, S. J. (1997). Proximate and mineral composition of three freshwater fishes. *Journal of Food and Agriculture*, 75(3), 312-314.
- Noel, T., Roger, P., Bernard, T., & Fabrice, T. D. (2016). Proximate composition, Fatty Acid and Mineral Contents of Four Freshwater Fish from Maga Lake (Far North Region of Cameroon). *American Journal of Food Science and Technology*, 4(3), 64-69.
- Farhat, J., & Abdul, S. C. (2011). Chemical compositions and fatty acid profiles of three fresh water fish species. *Food chemistry*, 12(5), 991-996.
- Hayam, M. I. (2015). Chemical composition, menirals contents, amino acids bioavailability and

- sensory properties of meat and fish balls containing fish protein isolate. *International Journal of Microbiology and Applied Science*, 4(4), 917-933
19. Magobe, O., Mosepele K., & Masamba, W. R. L. (2015). Essential mineral content of common fish species in Chanoga, Okavango delta, Botswana. *African journal of Food Science*, 2(3), 480-486.
  20. Ravichandran, S., Kumaravel, K., & Florence, P. E. (2011). Nutritive Composition of some Edible Fin Fishes. *International Journal Zoological Research*, 7(1), 341-251.
  21. Sidhu, P., Gorg, M. L., Morgenstern, P., Vogt, J., Butz, T., & Dhawan, D. K. (2004). Role of Zinc in regulating the levels of hepatic elements following nickel toxicity in rats. *Biochegeical Trace Elements Research*, 10(2), 161-172.
  22. Simsek, A., & Aykut, O. (2007). Evaluation of the microelement profile of Turkish hazelnut (*Corylus avellana* L) varieties for human nutrition and health. *International Journal of Food Science*, 5(8), 677-688.
  23. Soetan, K. O., Olaiya, C. O., & Oyewole, O. E. (2010). The importance of mineral elements for humans, domestic animals and plants: A review African. *Journal of Food Science*. 4(5), 200-222.
  24. Tejada-Jimenez M., Galvan A., Fernandez, E., & Lamas A. (2009). Homeostasis of the micronutrients Ni, Mo and Cl with specific biochemical functions. *Journal of Biology*, 1(2), 358-363.
  25. Underwood, E. J. (1971). Trace Elements in Human and Amino Nutrition, 3<sup>rd</sup> Edition, Academic Press, New York.
  26. Wood, R. J. (2000). Assessment of marginal Zinc status in humans. *Journal of Nutrition*. 13(2):1350-1353.
  27. Yeannes, I. M., & Almandos, M. E. (2003). Estimation of fish proximate composition starting from water content. *Journal of Food Composition*, 1(6), 81-92.