

Letter to the Editor

Neurological Disease Triggering Takotsubo Syndrome

Josef Finsterer, MD, PhD¹ and Claudia Stöllberger, MD²¹Krankenanstalt Rudolfstiftung, Messerli Institute, Vienna, Austria²2nd Medical Departments, Krankenanstalt Rudolfstiftung, Vienna, Austria

*Corresponding Author

Josef Finsterer, MD, PhD

Keywords: TTS, ICD9, ICD10 system.

In a recent article, Morris *et al.*, presented a retrospective study of patients with acute neurological disease, who also experienced Takotsubo syndrome (TTS) (Morris, N.A. *et al.*, 2018). We have the following comments and concerns.

TTS has not only been reported in association with epilepsy, subarachnoid bleeding, ischemic, stroke, intracerebral bleeding, migraine, Guillain-Barre syndrome, and traumatic brain injury as mentioned in the article by Morris *et al.*, but also in association with amyotrophic lateral sclerosis (Choi, S.J. *et al.*, 2017), mitochondrial disorder (Finsterer, J. *et al.*, 2007), thrombolysis of ischemic stroke (Kitagawa, T. *et al.*, 2018), non-convulsive status epilepticus (Uemura, J. *et al.*, 2016), zoster virus encephalitis (Bennett, L., & Iqbal, J. A. 2017), HLTV1-associated myelopathy (Yamanaka, S. *et al.*, 2017), myasthenia gravis (Battineni, A. *et al.*, 2017), Miller-Fisher syndrome (Gill, D. *et al.*, 2017), posterior reversible encephalopathy syndrome (PRES) (Grimaldi, S. *et al.*, 2017), syndrome of inadequate SDH secretion (Jha, K.K. *et al.*, 2016), transient global amnesia (TGA) (Sajeev, J. *et al.*, 2017), multiple sclerosis (Peller, M. *et al.*, 2016), baclofen withdrawal (Levy, J. *et al.*, 2016), post-anoxic encephalopathy (Batouche, D.D. *et al.*, 2016), Alzheimer's disease (Zuin, M. *et al.*, 2016), myotonic dystrophy type 1 (Fernández, A. M. *et al.*, 2016), acute disseminated encephalo-myelitis (Venkatraman, A. *et al.*, 2016), eclampsia (Gleich, S. J. *et al.*, 2016), delirium (Joy, P.S., & Kumar, G. 2015), panhypopituitarism (Plácido, R. *et al.*, 2016), entacarpone add-on (Baldacci, F. *et al.*, 2014), and botulism (Tonomura, S. *et al.*, 2017).

A main disadvantage of the study is that only ICD9 was applied. According to ICD10 the identifier for TTS is I51.81. Was this code also considered during the recruitment of patients with acute neurological disease and TTS? Due to ignoring the ICD10 system, a number of patients might have been missed during the search for appropriate patients.

Interestingly, the authors included hypertensive encephalopathy to the list of acute neurological disorders [1]. However, we do not regard hypertensive encephalopathy as “acute”. It is a chronic disease developing due to chronic arterial hypertension. The authors may mean an acute hypertensive crisis, for example due to pheochromocytoma previously reported in association with TTS, but this is not an acute neurological disease.

A further shortcoming of the study is that only diagnoses at dismissal from the hospital were considered. Mentioning TTS with an acute neurological disease on the report does not mean that these diagnoses are causally linked, and does not clarify if there was a timely relation. A patient may have been admitted for ischemic stroke but may have developed TTS four weeks later due to a completely different trigger. Furthermore, TTS may have been the initial event leading to admission and the neurological disease may have developed long after TTS, thus excluding a causal relation. However, both may occur on the report of same hospitalisation.

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easims/>

Article History

Received: 27.06.2019

Accepted: 09.07.2019

Published: 29.07.2019

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

In summary, this interesting study could be more meaningful, if additional neurological disorders would have been considered as triggers of TTS, if the ICD10 codes for TTS would have been additionally used, and if the time relation between the neurological event and TTS would have been clarified.

REFERENCES

- Morris, N.A., Chatterjee, A., Adejumo, O.L., Chen, M., Merkler, A.E., Murthy, S.B., & Kamel, H. (2018). The Risk of Takotsubo Cardiomyopathy in Acute Neurological Disease. *Neurocrit Care*. doi: 10.1007/s12028-018-0591-z.
- Choi, S.J., Hong, Y.H., Shin, J.Y., Yoon, B.N., Sohn, S.Y., Park, C.S., & Sung, J.J. (2017). Takotsubo cardiomyopathy in amyotrophic lateral sclerosis. *J Neurol Sci*, 375, 289-293.
- Finsterer, J., Stöllberger, C., Sehnal, E., Valentin, A., Huber, J., & Schmiedel, J. (2007). Apical ballooning (Takotsubo syndrome) in mitochondrial disorder during mechanical ventilation. *J Cardiovasc Med (Hagerstown)*, 8, 859-63.
- Kitagawa, T., Yamamoto, J., Kureshima, M., Maeda, H., & Nishizawa, S. (2018). Takotsubo Cardiomyopathy and Neurogenic Pulmonary Edema Following Fibrinolytic Therapy for Embolic Stroke: A Case Report. *No Shinkei Geka*, 46, 21-25.
- Uemura, J., Wada, Y., & Yagita, Y. (2016). Non-convulsive status epilepticus with Takotsubo cardiomyopathy: a case report. *Rinsho Shinkeigaku*, 56, 852-856.
- Bennett, L., & Iqbal, J. A. (2017). 68-year-old with cranial nerve neuropathies and a troponin rise. *Clin Med (Lond)*, 17, 575-577.
- Yamanaka, S., Nakayama, K., Tamai, H., Sakamaki, M., & Inokuchi, K. (2017). Adult T-cell leukemia-lymphoma complicated by Takotsubo cardiomyopathy and HTLV-1-associated myelopathy after treatment with the anti-CCR4 antibody mogamulizumab. *Rinsho Ketsueki* 58, 309-314.
- Battineni, A., Mullaguri, N., Thanki, S., Chockalingam, A., & Govindarajan, R. (2017). A Case Report of Recurrent Takotsubo Cardiomyopathy in a Patient during Myasthenia Crisis. *Case Rep Crit Care*, 2017:5702075. doi: 10.1155/2017/5702075.
- Gill, D., & Liu, K. (2017). Takotsubo cardiomyopathy associated with Miller-Fisher syndrome. *Am J Emerg Med* 2017;35:1012. doi: 10.1016/j.ajem.2016.12.050.
- Grimaldi, S., Doche, E., Rey, C., Laksiri, N., Boussen, S., Quilici, J., ... & Pelletier, J. (2017). Association of posterior reversible encephalopathy syndrome and transient apical ballooning syndrome (Takotsubo): first case report of a man and review of the literature. *Case reports in neurology*, 9(2), 173-178.
- Jha, K.K., Kumar, M., Jha, U., & Desai, S. (2016). Takotsubo cardiomyopathy in a patient with SIADH. *Int J Cardiol*, 225, 342-344.
- Sajeev, J., Koshy, A., Rajakariar, K., & Gordon, G. (2017). Takotsubo cardiomyopathy and transient global amnesia: a shared aetiology. *BMJ Case Rep*, 2017. pii: bcr-2017-219472. doi: 10.1136/bcr-2017-219472.
- Peller, M., Balsam, P., Budnik, M., Marchel, M., & Opolski, G. (2016). Reverse Takotsubo syndrome in a patient with diagnosed multiple sclerosis. *Kardiol Pol*, 74,1029.
- Levy, J., De Brier, G., Hugeron, C., Lansaman, T., & Bensmail, D. (2016). Takotsubo cardiomyopathy as a reversible complication of intrathecal baclofen withdrawal. *Ann Phys Rehabil Med*, 59, 340-342.
- Batouche, D.D., Elhalimi, K., & Benatta, N.F. (2016). Cardiac dysfunction secondary to an adrenergic storm or cardiomyopathy of stress among child victims of traumatism. *Ann Cardiol Angeiol (Paris)*, 65,171-4.
- Zuin, M., Dal Santo, P., Picariello, C., Conte, L., Zuliani, G., D'Elia, K., & Roncon, L. (2016). Takotsubo Cardiomyopathy in an Elderly Woman with Alzheimer's Disease: A Rare Association. *Case Report and Mini-Review of the Literature. J Am Geriatr Soc*, 64, 916-7.
- Fernández, A. M., Cejudo, D. C. D. L., Martínez, M. V., & Paule, S. A. (2016). Tako-Tsubo syndrome as first cardiac manifestation in Steinert's disease. *Medicina clinica*, 147(2), e7.
- Venkatraman, A., Bajaj, N. S., Khawaja, A., & Meador, W. (2016). Response to " Glucocorticoids for treating Takotsubo syndrome?" and " Atypical global Takotsubo syndrome in a patient with acute disseminated encephalomyelitis". *Clinical Autonomic Research*, 26(2), 163.
- Gleich, S. J., Barbara, D. W., Arendt, K. W., Rose, C. H., & Blauwet, L. A. (2016). Reverse apical ballooning echocardiographic pattern in eclampsia-related cardiomyopathy. *A & A case reports*, 6(1), 6-9.
- Joy, P.S., & Kumar, G. (2015). Delirium tremens is a risk factor for Takotsubo cardiomyopathy. *Int J Cardiol*, 191,185-6.
- Plácido, R., Martins, A. F., Robalo Martins, S., do Vale, S., Almeida, A. G., Pinto, F., & Martin Martins, J. (2016). Takotsubo Syndrome: A pathway through the pituitary disease. *Case reports in cardiology*, 2016.