Abbreviated Key Title: EAS J Parasitol Infect Dis ISSN 2663-0982 (Print) & 2663-6727 (Online) Open Access Published By East African Scholars Publisher, Kenya



Volume-1 | Issue-3 | May-June-2019 |

Case Report

A Primary Hydatid Cyst of Thigh: A Rare Case Report

Redouane Roukhsi^{1*}, Hassan Doulhousne², Salah ben elhend¹, Nabil H³, Elmehdi A³, Abdelilah Mouhsine³, El fikri A²

¹Radiology service – 3th Military Hospital – Laayoune – Morocco ²Radiology service – 5th Military Hospital – Guelmim – Morocco ³Radiology service – HMA Military Hospital – Marrakech – Morocco

*Corresponding Author Redouane Roukhsi

Abstract: Hydatid disease is a parasitic disease that is endemic in many parts of the world, especially in Morocco. Primary hydatid disease of musculoskletal system is rare. We present an unusual case of a primary hydatid cyst of thigh. **Introduction:** Primary hydatid disease of musculoskletal system is rare. We present an unusual case of a primary hydatid cyst of thigh. **Introduction:** Primary hydatid disease of musculoskletal system is rare. We present an unusual case of a primary hydatid cyst of thigh.

Keywords: Hydatid disease – Musculoskletal - Thigh - MRI.

Case Report:

A 53 year old man presented with larg swelling in medial aspect of the left thigh. The pain is severe on walking, and associated with a limp. On local examination, swelling was approximately 20×08 cm. On examination, he was febrile with a temperature of

37.8°C. Blood pressure was 130/80 mmHg and pulse rate was 100/min. Ultrasonography revealed to be a cystic swelling. Magnetic resonance imaging (MRI) (Figure 1 /a, b, c, d, and e) further reinforced the diagnosis of hydatid disease. A MRI scan of the hip shows classic features of the hydatid cyst of thigh.



Quick Response Code



Journal homepage: http://www.easpublisher.com/easjpid/ Article History Received: 10.06.2019 Accepted: 25.06.2019

Published: 13.07.2019

Copyright @ 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited. Redouane Roukhsi et al., EAS J Parasitol Infect Dis; Vol-1, Iss-3 (May-June, 2019): 61-62



Figure 1: Longitudnal (a, b, c) and transverse (d, e) MRI section through the lesion.

DISCUSSION:

Hydatid disease is a parasitic infestation of and herbivorous animals, humans caused hv echinococcus granulosus. Hydatid disease can affect virtually any organ system in body and should be kept as differential diagnosis of cystic lesion. The imaging methods used for diagnosis and evaluation of the extent of Hydatid disease are ultrasonography, computed tomography, magnetic resonance imaging, and less commonly radiography and urography (Mehta, P. et al., 2016). MRI was advised to confirm the diagnosis and for clear identification of involved structures and for surgical planning (Bansiwal, R. K. et al., 2011). Typical signs of hydatidosis are multivesicular lesions with or without hypointense peripheral ring (Garcia-Diez, A. I. et al., 2000) (rim sign). En bloc resection alone is curative for intramuscular hydatid disease (García-Alvarez, F. et al., 2002; Arazi, M. et al., 2005).

CONCLUSION:

Hydatid disease can affect virtually any organ system in body. Commonly, it demonstrates characteristic imaging findings that help to differentiate hydatid disease from other entities.

REFERENCES:

- 1. Mehta, P., Prakash, M., & Khandelwal, N. (2016). Radiological manifestations of hydatid disease and its complications. *Tropical parasitology*, 6(2), 103.
- Bansiwal, R. K., Sharma, R., & Attri, A. K. (2011). A large primary hydatid cyst of thigh: a case report. *Indian Journal of Surgery*, 73(2), 158-160.
- Garcia-Diez, A. I., Mendoza, L. R., Villacampa, V. M., Cozar, M., & Fuertes, M. I. (2000). MRI evaluation of soft tissue hydatid disease. *European* radiology, 10(3), 462-466.
- García-Alvarez, F., Torcal, J., Salinas, J. C., Navarro, A., García-Alvarez, I., Navarro-Zorraquino, M., ... & Lozano, R. (2002). Musculoskeletal hydatid disease: a report of 13 cases. *Acta Orthopaedica Scandinavica*, 73(2), 227-231.
- Arazi, M., Erikoglu, M., Odev, K., Memik, R., & Ozdemir, M. (2005). Primary echinococcus infestation of the bone and muscles. *Clinical Orthopaedics and Related Research*[®], 432, 234-241.