

## Letter to the Editor

## Kawasaki Disease in a *LDB3* Mutation Carrier with Subclinical Myopathy and Noncompaction

Josef Finsterer, MD, PhD

Krankenanstalt Rudolfstiftung, Messerli Institute, Vienna, Austria

\*Corresponding Author  
Josef Finsterer, MD, PhD

**Keywords:** myopathy, non-compaction, hypertrabeculation, mutation, gene, congenital muscular dystrophy.

In a recent article, Hachiya *et al.*, reported about a 6 months-old male with Kawasaki disease (KD) who also carried a *LDB3* mutation and presented with noncompaction(LVHT) (Hachiya, A. *et al.*, 2016). We have the following comments and concerns.

*LDB3* is one of the six cardinal genes (*DES*, *CRYAB*, *LDB3/ZASP*, *MYOT*, *FLNC*, *BAG3*), of which mutations cause myofibrillar myopathy(MM), characterised by focal abnormal accumulation of intra-sarcoplasmic proteins, the presence of vacuoles, and disorganisation of the inter-myofibrillar network beginning at the Z-disk (Béhin, A. *et al.*, 2015). In addition, structural and functional abnormalities of mitochondria have been described (Jackson, S. *et al.* 2015). Recently, it has been shown that mutations in the *DNAJB6*, *FHL1*, *PLEC*, or *TTN* gene, encoding for Z-disc proteins as well, may also cause MM (Jackson, S. *et al.*, 2015). Did the index case or any of his relatives present with clinical features of MM, such as slowly progressive diffuse weakness of the limb muscles, weakness of the distal muscles, or respiratory failure due to affection of the respiratory muscles? Did any of the mutation-carriers report muscle stiffness, aching, or cramps (Selcen, D. *et al.*, 2005 Jan 28). Were there indications for peripheral neuropathy occurring in 20% of the cases (Selcen, D. *et al.*, 2005 Jan 28). Which were the results of nerve-conduction studies, muscle MRI, and muscle biopsy?

We should be informed about those first-degree relatives who also carried the *LDB3* mutation. In which relatives was the *LDB3* mutation detected? Which of those who carried the mutation were

symptomatic and which were asymptomatic? Which subjects had developed only cardiac manifestations and which presented with muscle disease and cardiac involvement? Did transmission of the mutation follow an autosomal dominant or autosomal recessive trait?

Mutations in *LDB3/ZASP/cypher* may not only be associated with LVHT but also with arrhythmogenic right ventricular dysplasia. Did the patient ever develop phenotypic features such as hepatomegaly, palpitations, syncope, dyspnoea, exercise-intolerance, or coma? Did endomyocardial biopsy show right ventricular fibrofatty replacement, predisposing for premature ventricular contractions, sustained or non-sustained re-entrant ventricular tachycardia, supra-ventricular ectopic beats, supra-ventricular tachycardia, atrial flutter, atrial fibrillation, or sick sinus syndrome?

LVHT is frequently complicated by heart failure, ventricular arrhythmias, including sudden cardiac death, or cardiac embolism. Was heart failure and systolic dysfunction attributed to KD rather than LVHT or was it a complication of another type of cardiomyopathy, occurring in 15-30% of the cases (Selcen, D. *et al.*, 2005 Jan 28)?

Since cardiac disease in MM may go along with severe ventricular arrhythmias and conduction disturbances (Celestino-Soper, P. B. *et al.*, 2015), we should be informed about the results of long-term ECG recordings and if the individual or family history was positive for palpitations or syncopes. Did any of the first-degree relatives require a pacemaker or an ICD?

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easms/>

Article History

Received: 12.09.2019

Accepted: 28.09.2019

Published: 10.10.2019

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

We do not agree with the notion that mutations in *LDB3* cause LVHT [1]. Though LVHT is associated with mutations in >40 different genes and a number of chromosomal defects, a causal relation has never been proven.

Overall, this interesting case would profit from a more clearcut delineation between manifestations of KD and those of the *LDB3* mutation. Relatives need to be investigated for the mutation and LVHT. Muscle manifestations of the *LDB3* mutation, including hyperCKemia, need to be encountered.

#### REFERENCES

1. Hachiya, A., Motoki, N., Akazawa, Y., Matsuzaki, S., Hirono, K., Hata, Y., ... & Koike, K. (2016). Left ventricular non-compaction revealed by aortic regurgitation due to Kawasaki disease in a boy with *LDB3* mutation. *Pediatrics International*, 58(8), 797-800.
2. Béhin, A., Salort-Campana, E., Wahbi, K., Richard, P., Carlier, R. Y., Carlier, P., ... & Franques, J. (2015). Myofibrillar myopathies: state of the art, present and future challenges. *Revue neurologique*, 171(10), 715-729.
3. Jackson, S., Schaefer, J., Meinhardt, M., & Reichmann, H. (2015). Mitochondrial abnormalities in the myofibrillar myopathies. *European journal of neurology*, 22(11), 1429-1435.
4. Selcen, D., Engel, A.G., Myofibrillar, M. (2005 Jan 28). [updated 2012 Oct 29]. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD, Fong CT, Mefford HC, Smith RJH, Stephens K, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2016. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1499/>
5. Celestino-Soper, P. B., Doytchinova, A., Steiner, H. A., Uradu, A., Lynnes, T. C., Groh, W. J., ... & Liu, Y. (2015). Evaluation of the genetic basis of familial aggregation of pacemaker implantation by a large next generation sequencing panel. *PLoS one*, 10(12), e0143588.