

Original Research Article

“Intra-Operative & Post-Operative Complications in Pediatric Anesthesia- A Prospective Observation Study”

Md. Anwarul Islam^{1*}, Abu Zahed Md. Firoz²¹Assistant Professor, Department of Anaesthesiology, Shaheed Ziaur Rahman Medical College, Bogura, Bangladesh²Assistant Professor, Department of Anaesthesiology, Shaheed Ziaur Rahman Medical College, Bogura, Bangladesh**Article History**

Received: 05.06.2021

Accepted: 01.07.2021

Published: 12.07.2021

Journal homepage:<https://www.easpublisher.com>**Quick Response Code**

Abstract: Introduction: Typical complications in pediatric anesthesia are respiratory problems, medication errors, difficulties with the intravenous puncture and pulmonary aspiration. Pediatric patients differ in their drug requirements because of their smaller body size, differences in body composition and handling capacity of drugs. Usually dosages are based on body weight, because it correlates so intimately with body water compartments. **Objective:** To find out the Intra-Operative & Post-Operative Complications in Pediatric Anesthesia. **Materials & Methods:** The present study was conducted in the Department of Anaesthesiology, Shaheed Ziaur Rahman Medical College Hospital, Bogura, Bangladesh during from April 2020 to March 2021. It included 83 pediatric patients of both gender. Parents were informed regarding the study and written consent was obtained. Ethical clearance was taken from institutional ethical committee. General information such as name, age, sex, ASA status, technique of anesthesia and complications were recorded. **Results:** The prospective study was conducted to identify anesthesia-related complications in pediatric patients. The present study was conducted on 83 pediatric patients of both genders. Factors such as conscious state, colour, respiration, pulse and blood pressure were also assessed. Age group 1 month- 1 year had 12 males and 18 females. 1-5 years had 10 males and 15 females, 6-10 years had 11 males and 15 females and 11-18 years had 01 males and 01 females. The difference was non-significant (P-0.5). Patients had ASA I (72), ASA 2 (17), ASA 3 (5), ASA 4 (1) and ASA 5 (1) Status. The difference was significant (p-0.05). Type of anesthesia was general anesthesia (50), general anesthesia with caudal block (14), subarachnoid block (1) and general anesthesia with local anesthesia (18). The Difference was significant (p-0.01). Common intra-operative complications were bradycardia (3), tachycardia (2), hypotension (2), cardiac arrest (2), bronchospasm (1), hypoventilation (2), pain (3) and prolonged unconsciousness (1). Shows that post-operative complications were shivering (3), pain (4), hemorrhage (4), hypoventilation (2), respiratory arrest (1) and laryngospasm (2). The difference was non-significant (P-0.1). **Conclusion:** In order to avoid or minimize these complications occurring at induction to the barest minimum, a consultant anesthetist or an experienced anesthetist must always be in attendance to handle difficult cases.

Keywords: General Anesthesia, Laryngospasm, Pediatric.

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

I. INTRODUCTION

Typical complications in pediatric anesthesia are respiratory problems, medication errors, difficulties with the intravenous puncture and pulmonary aspiration. In the postoperative setting, nausea and vomiting, pain, and emergence delirium can be mentioned as typical complications. Major differences in anatomy and physiology in the small infant have important consequences in many aspects of anesthesia. The physical disparity between the adult and child diminishes at 10-12 years of age although major psychological differences continue till adolescence. In

order to reduce the morbidity and mortality associated with pediatric anesthesia, the anesthetist should have a sound knowledge of the physiologic, anatomic and pharmacologic peculiarities of each developmental stage of childhood. Laryngospasm is a common occurrence in pediatric anesthesia and is usually easy to manage. It is however a significant cause of severe adverse outcome including cardiac arrest and its sequelae [1]. Pediatric patients differ in their drug requirements because of their smaller body size, differences in body composition and handling capacity of drugs. Usually dosages are based on body weight,

because it correlates so intimately with body water compartments. Pediatric anesthesia morbidity and mortality in the perioperative period has been studied by Cohen et al [2], an incidence of 35% was reported by the same author. In another multicenter study, complications related to anesthesia in infants and children were also reported. A study carried out in Nigeria revealed an incidence of 10% adverse events in pediatric surgical emergencies. Advances in monitoring and the declining use of halothane have led to a change in the types of risk. Risk is defined as the chance or possibility of danger, loss or injury. It is therefore an ever-present part of life. What constitutes a serious or worrying risk is dependent upon perspective. This is readily illustrated in the health care industry. Risk is viewed in quite different lights by patients, doctors, hospital managers, insurance companies and policy-makers. While patients worry about the prospect of failed treatment and unpleasant side effects, anesthetists will be more concerned about the immediate risks of major disability and death associated with surgery. Similarly, while individual hospital managers and insurance companies are keen to minimize their exposure to financial liability, health policy-makers aim to ensure patients' access to the most appropriate centers for their treatment [3]. The present study was conducted to identify anesthesia related complications in pediatric patients.

II. MATERIALS & METHODS

The prospective observation study was conducted in the Department of Anaesthesiology, Shaheed Ziaur Rahman Medical College Hospital, Bogura, Bangladesh during from April 2020 to March 2021. It included 83 pediatric patients of both gender. Parents were informed regarding the study and written consent was obtained. Ethical clearance was taken from institutional ethical committee. General information such as name, age, sex, ASA status, technique of anesthesia and complications were recorded. Factors such as conscious state, colour, respiration, pulse and blood pressure were also assessed. Results were subjected to statistical analysis using chi-square test. P-value 0.05 was considered significant. All data analysis windows "SPSS" version 20.0.

III. RESULTS

The present study was conducted on 83 pediatric patients of both genders. General information such as name, age, sex, ASA status, technique of anesthesia, and complications were recorded. Factors such as conscious state, colour, respiration, pulse and blood pressure were also assessed. Age group 1 month-1 year had 12 males and 18 females. 1-5 years had 10 males and 15 females, 6-10 years had 11 males and 15 females and 11-18 years had 01 males and 01 females. The difference was non-significant (p=0.5) (Table-1).

Table-1: Age distribution of patients (N=83)

Age	Males	Females	p value
1-12 months	12	18	0.5
1-5 years	10	15	
6-10 years	11	15	
11-18 years	01	01	
Total	34	49	




Fig-1: ASA status in patients was significant (P<0.05).

Shows that patients had ASA I (72), ASA 2 (17), ASA 3 (5), ASA 4 (1) and ASA 5 (1) Status. The difference was significant (P<0.05) (Fig 1).




Fig-2: Choice of Anesthesia

Shows that type of anesthesia was general anesthesia (50), general anesthesia with caudal block (14), subarachnoid block (1) and general anesthesia with local anesthesia (18). The Difference was significant (p<0.01) (Fig 2).




Fig-3: Intra-operative complications

Shows that common intra-operative complications were bradycardia (3), tachycardia (2), hypotension (2), cardiac arrest (2), brochospasm (1),

hypoventilation (2), pain (3) and prolonged unconsciousness (1). The difference was significant ($P=0.01$) (Fig 3).




Fig-4: Post-operative complications

Shows that post-operative complications were shivering (3), pain (4), hemorrhage (4), hypoventilation (2), respiratory arrest (1) and laryngospasm (2). The difference was non-significant ($P=0.1$) (Fig 4).

IV. DISCUSSION

Standard monitoring is essential in pediatric anesthesia care particularly for early detection and prevention of dangerous complications such as bradycardia, hypotension, hypoxia and adverse temperature changes. The hallmark of pediatric anesthesia has always been the precordial stethoscope because it allows constant monitoring of breath sounds, along with the heart sounds, rate and rhythm. Changes in clinical status happen quickly in the newborn with only subtle signs heralding serious alterations in the patient's clinical status [4]. Difficulty of intravenous cannulation is sometimes encountered especially in the preterm neonates, overweight babies and when most peripheral veins had been ruined from withdrawal of blood sample for laboratory investigations and intravenous therapy. This problem was sometimes solved with the help of pediatricians or rarely the surgeons who perform venous cut downs [5]. The present study was conducted to identify anesthesia-related complications in pediatric patients. In this study, maximum patients were in age group 1-12 months. The type of anesthesia was general anesthesia, general anesthesia with caudal block, subarachnoid block and general anesthesia with local anesthesia. This is similar to Murray et al., [6] complications arising at induction of anesthesia in neonates and small infants in our study were mainly respiratory and cardiovascular. They may manifest as apnea, bronchospasm, laryngospasm, cyanosis, bradycardia and cardiac arrest. These are likely to be due to stimulation of the larynx during laryngoscopy and tracheal intubation at light planes of anesthesia. Neonates and infants have very active airway reflexes and therefore require adequate depth of anesthesia for laryngoscopy and tracheal intubation. The preterm infants are more prone to developing postoperative respiratory complications as evidenced by our study. In neonates, if bradycardia is not promptly

treated with 100% oxygen, they readily develop hypoxic cardiac arrest [7]. We found that common intra-operative complications were bradycardia, tachycardia, hypotension, hypertension, cardiac arrest, bronchospasm, hypoventilation, pain and prolonged unconsciousness. Post-operative complications were shivering, pain, hemorrhage, hypoventilation, respiratory arrest, laryngospasm and hypoventilation. This is in agreement with Jimenez N [8]. It has been recognized that infants and children require adequate pain management and if not received may actually adversely affect patient outcome. Age appropriate pain assessment is essential and both subjective and objective assessment tools may be utilized depending on patient age and clinical status. Many pain assessment scales have been developed to assess pain in children. The most reliable form of pain assessment is self-report. Infants less than 1 month old have the highest rate of adverse events both intraoperatively and in the recovery room [9]. Anesthesia, because of its depression on metabolism and dilatation of cutaneous vessels, tends to cause a progressive fall in body temperature [10, 11]. In the recovery room, pain was treated with opioid analgesics after excluding other causes of crying and discomfort. Our study showed a similar finding. The presence of pain implies inadequate analgesia following surgery. There have been a number of myths and misunderstandings related to pain in children which has led to historically inadequate treatment of pediatric pain [12].

V. CONCLUSION

In order to avoid or minimize these complications occurring at induction to the barest minimum, a consultant anesthetist or an experienced anesthetist must always be in attendance to handle difficult cases.

REFERENCES

1. Murat, I., Constant, I., & Maud'huy, H. (2004). Perioperative anaesthetic morbidity in children: a database of 24 165 anaesthetics over a 30-month period. *Pediatric Anesthesia*, 14(2), 158-166.
2. Tired, L., Nivoche, Y., Hatton, F., Desmonts, J. M., & Vourc'h, G. (1988). Complications related to anaesthesia in infants and children: a prospective survey of 40240 anaesthetics. *British journal of anaesthesia*, 61(3), 263-269.
3. Cohen, M. M., Cameron, C. B., & Duncan, P. G. (1990). Pediatric anesthesia morbidity and mortality in the perioperative period. *Anesthesia & Analgesia*, 70(2), 160-167.
4. Tay, C. L. M., Tan, G. M., & Ng, S. B. A. (2001). Critical incidents in paediatric anaesthesia: an audit of 10 000 anaesthetics in Singapore. *Pediatric Anesthesia*, 11(6), 711-718.
5. Flick, R. P., Sprung, J., Harrison, T. E., Gleich, S. J., Schroeder, D. R., Hanson, A. C., ... & Warner, D. O. (2007). Perioperative cardiac arrests in children between 1988 and 2005 at a tertiary

- referral center: a study of 92,881 patients. *The Journal of the American Society of Anesthesiologists*, 106(2), 226-237.
6. Morray, J. P., Geiduschek, J. M., Caplan, R. A., Posner, K. L., Gild, W. M., & Cheney, F. W. (1993). A comparison of pediatric and adult anesthesia closed malpractice claims. *Anesthesiology*, 78(3), 461-467.
 7. Gobbo Braz, L., Braz, J. R. C., Módolo, N. S. P., Nascimento, P. D., Brushi, B. A. M., & Raquel De Carvalho, L. (2006). Perioperative cardiac arrest and its mortality in children. A 9- year survey in a Brazilian tertiary teaching hospital. *Pediatric Anesthesia*, 16(8), 860-866.
 8. Jimenez, N., Posner, K. L., Cheney, F. W., Caplan, R. A., Lee, L. A., & Domino, K. B. (2007). An update on pediatric anesthesia liability: a closed claims analysis. *Anesthesia & Analgesia*, 104(1), 147-153.
 9. Coté, C. J., Rolf, N., Liu, L. M., Goudsouzian, N. G., Ryan, J. F., Zaslavsky, A., ... & Alifimoff, J. K. (1991). A single-blind study of combined pulse oximetry and capnography in children. *The Journal of the American Society of Anesthesiologists*, 74(6), 980-987.
 10. Feldman, S. (1978). Neurological disease in relation to Anaesthesia – *A Practice of Anaesthesia* 4th Edition 937.
 11. Roe, C.F. (1970). Temperature Regulation in Anaesthesia. *Physiological and Behavioural Temperature regulation*.
 12. Cote, C.J. (1990). *Paediatric Anaesthesia*: In *Anesthesia* Miller R.D. 1990, Churchill Livingstone. 3rd Edition, 2; 1897-1926.

Cite this article: Anwarul Islam & Abu Zahed Md. Firoz (2021). "Intra-Operative & Post-Operative Complications in Pediatric Anesthesia-A Prospective Observation Study". *EAS J Anesthesiol Crit Care*, 3(4), 47-50.