

Letter to the Editor

Neurogenic Pulmonary Edema – The Takotsubo Syndrome of the Lungs

Josef Finsterer, MD, PhD*

Krankenanstalt Rudolfstiftung, Messrli Institute, Vienna, Austria

*Corresponding Author
Josef Finsterer, MD, PhD**Keywords:** lung edema, stress reaction, Takotsubo, epilepsy, brain-lung disease.

In a recent article Cho *et al.*, reported about a 31 years old male who experienced an inhalation injury and CO-poisoning after a fire in his apartment (Cho, K. *et al.*, 2018). During hyperbaric oxygen therapy (HBOT), applied for the CO-poisoning, he experienced a single generalised tonic-clonic seizure followed by lung edema (Cho, K. *et al.*, 2018). The case report has a number of shortcomings and raises concerns.

We do not agree with the conclusion that lung edema resulted from HBOT (Cho, K. *et al.*, 2018). Acute lung edema after a generalised seizure may be the direct consequence of the seizure, as has been previously reported (July, M. *et al.*, 2014). Acute lung edema immediately after an acute, severe neurological injury, such as acute meningitis, subarachnoid bleeding, seizures, ischemic stroke, intracerebral bleeding, epidural bleeding, traumatic brain injury, multiple sclerosis exacerbation, or vertebral artery dissection, is termed neurogenic pulmonary edema (NPE) (Al-Dhahir, M.A., & Sharma, S. 2018).

An argument in favour of NPE as the consequence of a stress-reaction to a serious cerebral event is the simultaneous occurrence of NPE and TTS in a patients with meningioma (Goyal, A., & Bharadwaj, S. 2019), with ischemic stroke and thrombolysis (Kitagawa, T. *et al.*, 2018), in patients with brainstem strokes (Probasco, J. C. *et al.*, 2014), in a patient with epidural hemorrhage (Nakamori, Y. *et al.*, 2018), in patients with subarachnoid bleeding (Manto, A. *et al.*, 2014), and in patients with brainstem hemangioblastoma (Gekka, M. *et al.*, 2014).

A further shortcoming of the study is that no MRI of the brain was carried out. CO-poisoning is well known to cause hyperintensities on diffusion weighted imaging (DWI). In a retrospective study of 387 patients experiencing CO-poisoning, 26.9% had DWI hyperintensities on cerebral MRI (Jeon, S. B. *et al.*, 2018). DWI lesions could easily explain why the patient developed a seizure. DWI lesions may not show up on cerebral CT scans why carrying out an MRI is obligatory. Since CO-poisoning goes along with cerebral lesions, they are more plausible to explain the development of seizures than excessive oxygen consumption.

The main therapeutic options for treating NPE are supportive measures and lowering of the intracranial pressure (Sarı, M. Y. *et al.*, 2015). Thus we should be informed if the index case had developed any clinical features or features on imaging of increased intracranial pressure. In case the patient had experienced an increase in intracranial pressure we should be informed which kind treatment was applied.

In a recent study it has been shown that endothelin-1 may be elevated in the cerebrospinal fluid (CSF) of patients with NPE due to infection with enterovirus-71 (Tu, Y. F. *et al.*, 2015). We should be informed if endothelin-1 was elevated in the CSF of the index case. Diagnosing NPE is still challenging but transpulmonary thermodilution studies can be helpful to establish the diagnosis. We are interested to know if this diagnostic approach was applied and how NPE was diagnosed in the index case. Missing in the report is a description of the EEG.

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easjms/>

Article History

Received: 02.12.2019

Accepted: 15.12.2019

Published: 26.12.2019

Copyright © 2019 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

Overall, this interesting case report could be more meaningful if an MRI of the brain had been carried out in the acute stage of CO-poisoning, if NPE triggered by the tonic-clonic seizure would have been more extensively discussed, and if treatment of NPE would have been more extensively described.

REFERENCES

1. Cho, K., Minami, T., Okuno, Y., Kakuda, Y., Tsutsumi, T., Kogame, T., ... & Koike, K. (2018). Convulsive seizure and pulmonary edema during hyperbaric oxygen therapy: A case report. *The Journal of Medical Investigation*, 65(3.4), 286-288.
2. July, M., Williamson, J. E., & Lolo, D. (2014). Neurogenic pulmonary edema following non-status epileptic seizure. *The American journal of medicine*, 127(3), e9-e10.
3. Al-Dhahir, M.A., & Sharma, S. (2018). Pulmonary Edema, Neurogenic. Oct 30. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2018 Jan-. Available from <http://www.ncbi.nlm.nih.gov/books/NBK532984/>
4. Goyal, A., & Bharadwaj, S. (2019). Neurogenic Pulmonary Edema and Stunned Myocardium in a Patient With Meningioma: A Heart-Brain Cross Talk. *Journal of neurosurgical anesthesiology*, 31(2), 264-265.
5. Kitagawa, T., Yamamoto, J., Kureshima, M., Maeda, H., & Nishizawa, S. (2018). Takotsubo cardiomyopathy and neurogenic pulmonary edema following fibrinolytic therapy for embolic stroke: a case report. *No shinkei geka. Neurological surgery*, 46(1), 21-25.
6. Probasco, J. C., Chang, T., Victor, D., & Nyquist, P. (2014). Isolated pulmonary edema without myocardial stunning in brainstem strokes. *Journal of neurology & translational neuroscience*, 2(1), 1040.
7. Nakamori, Y., Miyazawa, N., Yoshitani, K., & Yamamoto, S. (2018). A Pediatric Case With Takotsubo Cardiomyopathy and Neurogenic Pulmonary Edema Due to an Epidural Hemorrhage. *Journal of neurosurgical anesthesiology*, 30(3), 279-280.
8. Manto, A., De Gennaro, A., Manzo, G., Serino, A., Quaranta, G., & Cancellata, C. (2014). Early endovascular treatment of aneurysmal subarachnoid hemorrhage complicated by neurogenic pulmonary edema and Takotsubo-like cardiomyopathy. *The neuroradiology journal*, 27(3), 356-360.
9. Gekka, M., Yamaguchi, S., Kazumata, K., Kobayashi, H., Motegi, H., Terasaka, S., & Houkin, K. (2014). Hemorrhagic onset of hemangioblastoma located in the dorsal medulla oblongata presenting with tako-tsubo cardiomyopathy and neurogenic pulmonary edema: A case report. *Case reports in neurology*, 6(1), 68-73.
10. Jeon, S. B., Sohn, C. H., Seo, D. W., Oh, B. J., Lim, K. S., Kang, D. W., & Kim, W. Y. (2018). Acute brain lesions on magnetic resonance imaging and delayed neurological sequelae in carbon monoxide poisoning. *JAMA neurology*, 75(4), 436-443.
11. Sarı, M. Y., Yıldızdaş, R. D., Yükselmiş, U., & Horoz, Ö. Ö. (2015). Our patients followed up with a diagnosis of neurogenic pulmonary edema. *Turkish Archives of Pediatrics/Türk Pediatri Arşivi*, 50(4), 241.
12. Tu, Y. F., Lin, C. H., Lee, H. T., Yan, J. J., Sze, C. I., Chou, Y. P., ... & Huang, C. C. (2015). Elevated cerebrospinal fluid endothelin 1 associated with neurogenic pulmonary edema in children with enterovirus 71 encephalitis. *International Journal of Infectious Diseases*, 34, 105-111.