

Review Article

A Review on Ethnobotanical, Phytochemistry, Bioactivities and Medicinal Mysteries of *Fumaria officinalis* (Common Fumitory)

Rintu Dutta^{1*}, Mukesh Kumar Sharma¹ and Megha Jha²¹Maharaj Vinayak Global University, Jaipur, Rajasthan 302015, India²Pinnacle Biomedical Research Institute (PBRI), Near, Bharat Scout and Guides Campus, Shanti Marg, Shyamla Hills Road, Bhopal, Madhya Pradesh 462003, India

*Corresponding Author

Rintu Dutta

Abstract: Plant preparations are said to be medicinal or herbal when they are used to promote health beyond basic nutrition. It is indeed quite interesting to observe that our ancients were duly equipped with the vast, in depth and elaborated knowledge of drugs from the vegetable origin but unfortunately they possessed a scanty knowledge with regard to the presence of chemically pure compounds in most of them. *Fumaria officinalis* (*F. officinalis*, Fumariaceae) is a well-known traditional herb rich in biochemically active components. *F. officinalis* (common fumitory or earth smoke) is the most common species of the genus *Fumaria* in Western and Central Europe. Fumitory was officially recognized in 1986 by the French Health authorities as an herbal medicine which was traditionally used in renal and digestive elimination functions. By accurately looking to the name of fumitory, it is said to be derived either from the fact that its whitish, blue-green colour gives it the appearance of smoke rising from the ground, or according to Pliny and Olivier de Serres (XIV century) because the juice of the plant brings on such a flow of tears that the sight becomes dim as with smoke and hence its reputed use in affections of the eye. This herb has been known since antiquity and was described in herbals from the Middle-Ages. It was mainly the Mediterranean genus which was once used as medicine and wound healing. In a meanwhile traditionally fumitory has been used as digestive and diuretic. This review highlights the traditional, ethnobotanical, phytochemical, pharmacological information available on *F. officinalis*, which might be helpful for scientists and researchers to find out new chemical entities responsible for its claimed traditional uses.

Keywords: *Fumaria officinalis*, Fumariaceae, Fumitory, Mediterranean genus, Ethnobotanical, Phytochemical, Pharmacological.

INTRODUCTION

The genus *Fumaria* (Fumariaceae) consists of 46 species in the world and *Fumaria* species are known as fumitory, earth smoke, beggary, fumus, vapor, fumittery or wax dolls in English (Orhan, I. *et al.*, 2010). As per data available over three-quarters of the world population relies mainly on plants and plant extracts for their health care needs. More than 30% of the entire plant species, at one time or other was used for medicinal purposes. Treatment with medicinal plants is considered very safe as there is no or minimal side effects. These remedies are in sync with nature, which is the biggest advantage. The ancient scholars only believed that herbs are only solutions to cure a number of health related problems and diseases (Rajagopal, P.L. *et al.*, 2018). Fumitory is on the United

Kingdom General Sales List (GSL) and is approved by the German Commission E Monograph. It has been used in Europe and universal, as a traditional medicine for more than 30 years without safety problems. Phytochemical studies revealed the attendance of several alkaloids such as adlumidicine, copticine, fumariline, perfumine, protopine (Popova, M.E. *et al.*, 1982), fumaranine, fumaritine, paprafumicin and paprarine (Rahman, A.R. *et al.*, 1992). Fumarity has also been evaluated for pharmacologically effects, therapeutic effects and shown to possess antihelmantic (Hördegen, P. *et al.*, 2003), antipyretic (Khattak, S.G. *et al.*, 1985) and hypoglycemic properties (Akhtar, M.S. *et al.*, 1984). It was usually thought to be good and causes healing effects for the eyes and to remove skin blemishes. Nowadays herbalists use it to treat skin

Quick Response Code



Journal homepage:

<http://www.easpublisher.com/easjpp/>

Article History

Received: 15.07.2019

Accepted: 29.07.2019

Published: 24.08.2019

Copyright © 2019 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

DOI: 10.36349/easjpp.2019.v01i04.004

diseases and conjunctivitis; as well as to cleanse and pure the kidneys. This plant has been extremely valued since at least Roman times for its tonic and blood cleansing effect upon the body the smoky or fummy origin of its name comes from the translucent colour of its flowers giving them the appearance of smoke or of hanging in smoke, and the slightly grey-blue haze colour of its foliage, also resembling smoke coming from the ground, particularly after morning dew (Hördegen, P. *et al.*, 2003). In this article we have gathered a briefly and targeted pack of information about *F. officinalis* which hope to be useful in near future for scientist society. Diverse medicinal uses of the plant stand in confirmation to its pharmacological activity profile revealed in the recent past. Consolidation of its pharmacological activities and its correlation with its traditional uses would open new areas of research for discovery of drugs and various formulations (Gowher, G. 2017).

Taxonomy

Kingdom: Plantae
 Subkingdom: Tracheobionta
 Superdivision: Spermatophyta
 Division: Magnoliopsida
 Class: Magnoliopsida
 Subclass Magnoliidae
 Order: Papaveraceae
 Family: Fumariaceae – Fumitory family
 Genus: *Fumaria* L.
 Species: *Fumaria officinalis* L. ssp. *officinalis*

Common Name

Arab- Buklat-ul-malik, Shahtara
 Sanskrit-Khsetra parpati, Yavanaparpati
 Tamil-Tura
 Telugu -Chata-rashi
 Hin-Pit-parapara
 Kannada-D
 Urdu-Shahterah
 English- Common Fumitory, Earth-Smoke, Wax doll,
 Hedge fumitory

Botanical Description

It is an herbaceous annual plant that grows weakly erect and scrambling, with stalks about 10-50 cm (3.9-19.7 in) long. It has slender green leaves (Reader's Digest Field Guide to the Wild Flowers of Britain. 1981). Its pink 7-9 mm (0.28-0.35 in) flowers appear from April to October in the northern hemisphere¹⁰, or May to September in the UK (Reader's Digest Field Guide to the Wild Flowers of Britain. 1981). They are two lipped and spurred, with sepals running a quarter the length of the petals (Richard, F. *et al.*, 1974). The plant commonly has more than 20 and up to 60 flowers per spike (Popova, M.E. *et al.*, 1982). The fruit is an achene containing one seed. It is approximately globular, slightly wider than high and with an apical notch (Murphy, R.J. 2009).

Geographical Distribution

The *Fumaria* is a genus of herbs distributed in Asia, Europe and Africa. It is native to temperate regions of North Africa, Europe and parts of Western Asia. It is found in North Africa, within Macaronesia, Canary Islands, Algeria, Egypt, Libya, Morocco and Tunisia. Within Western Asia it is found in the Caucasus, Cyprus, Iraq, Israel, Lebanon, Siberia, Syria and Turkey. In Eastern Europe, it is found within Belarus, Estonia, Latvia, Lithuania and Ukraine. In middle Europe, it is in Austria, Belgium, Germany, Hungary, Netherlands, Poland, Slovakia and Switzerland. In northern Europe, in Denmark, Ireland, Norway, Sweden and United Kingdom. In southeastern Europe, within Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Greece, Italy, Macedonia, Montenegro, Romania, Serbia and Slovenia. Also in southwestern Europe, it is found in France, Portugal and Spain (Taxon: *Fumaria officinalis* L. 2017).

Alkaloids

The main alkaloid of *F. officinalis* is protopine. It is the major secondary metabolite extracted from *F. officinalis* and purified by column chromatography. Urine samples were composed from horses and a human volunteer that had been administered with either *F. officinalis* or protopine free base. In other hand, urine and plant samples were acetylated and analysed by GC-MS after solid-phase extraction. It was recognised that the urinary metabolites of protopine were identified as 4,6,7,13-tetrahydro-9,10-dihydroxy-5-methyl-benzo[e]-1,3-benzodioxolo[4,5-1] [2]benzazecin-12(5H)-one, 4,6,7,13-tetrahydro-10-hydroxy-9-methoxy-5-methyl-benzo[e]-1,3-benzodioxolo[4,5-1][2]benzazecin-12(5H)-one and 4,6,7,13-tetrahydro-9-hydroxy-10-methoxy-5-methyl-benzo[e]-1, 3-benzodioxolo[4,5-1][2]benzazecin-12(5H)-one, chelanthifolia, isochelanthifoline and 2-O-desmethylchelanthifoline¹³. Some important alkaloids which have discovered and identified by gas chromatography mass spectrophotometry method from the *F. agraria*, *F. bastardii*, *F. capreolata*, *F. sepium*, *F. densiflora*, *F. faurei*, *F. officinalis* subsp. *officinalis*, *F. parviflora*, *F. petteri* subsp. *calcarata* and *F. macrosepala*, are the isoquinoline alkaloids such as protopine, cryptopine, sinactine, stylophine, bicuculline, adlumine, parfumine, fumariline, fumarophycine, fumaritine, dihydrofumariline, parfumidine and dihydrosanguinarine (Sharma, U.R. *et al.*, 2012). The isoquinoline alkaloids in *F. officinalis* mostly have shown the biological activity (Hentschel, C. *et al.*, 1995; Gilani, H.A. *et al.*, 2005). Traditionally, the juices of *F. officinalis*, after undergoing evaporation process, could be used to treat chronic eczema, dermatological problems and cutaneous eruptions (Dermadosian, A., & Beutler, J.A. 2005; Duke, J.A. 2002).

Etymology

The "smoky" or "fummy" origin of its name comes from the translucent color of its flowers, giving them the appearance of smoke or of hanging in smoke, and the slightly gray-blue haze color of its foliage, also resembling smoke coming from the ground, especially after morning dew. The plant was already called *fūmus terrae* (smoke of the earth) in the early 13th century, and two thousand years ago, Dioscorides wrote in *De Materia Medica* (Περὶ ὕλης ἰατρικῆς) and Pliny the Elder in *Naturalis Historia* that rubbing the eyes with the sap or latex of the plant causes tears, like acrid smoke (*fūmus*) does to the eyes. Its Greek name is *kapnos* (καπνός, for *smoke*) and the name *fumewort*

now applies mostly to the genus *Corydalis*, especially the similar looking *Corydalis solida* (formerly *Fumaria bulbosa*), which was thought to belong to the same genus as fumitory (The Names of Plants, 4TH Edition Gledhill, D. 1985–2008).

Herbalism

It was traditionally thought to be good for the eyes, and to remove skin blemishes. In modern times herbalists use it to treat skin diseases, and conjunctivitis; as well as to cleanse the kidneys. However, Howard (1987) warns that fumitory is poisonous and should only be used "under the direction of a medical herbalist (Howard, M. 1987).

Pharmacological Activity

Table 1 Pharmacological activity of *F. officinalis*

System	Effect	Description	Active compounds	References	
All	Anti-malaria		Drank	Lombardi Satriani	(1951)
			different types		
			of decoctions		
		Bactericidal activity against the Gram-positive organisms		Preininger (1975)	
	Anti-bacterial	<i>Bacillus anthracis</i>			
		and <i>Staphylococcus</i> has been reported			
		Antibacterial activities of aqueous extract of <i>F. officinalis</i>	Extract		Dulger <i>et al.</i> , (2004)
					Boucard <i>et al.</i> ,
Digestive	Amphocholeretic activity				Boucard, M., & Laubenheimer, B. (1966)
					Reynier, M. <i>et al.</i> , (1977)
	Fumitory extract inhibited the formation of gall bladder calculi in animals		Extract		Lagrange, E. (1973)
	Management of disorders of hepatobiliary tract, spastic discomfort in the area of the gallbladder bile ducts as well as gastrointestinal tract				Gruenwald <i>et al.</i> , 2007
			Extract		Gruenwald <i>et al.</i> , (2007); Duke 2002
		<i>Plantago major</i> with <i>F. officinalis</i> ethanolic extracts together	Extract		Denden S <i>et al.</i> ,
	Antiallergic and choleric	(Ethanolic extract)			Denden, S. (2010); Ali Nazarizadeh <i>et al.</i> , 2013

	Ameliorate bile duct blockage in animals			Zacharewicz <i>et al.</i> , 1979
	and assist in the management of similar disorders in humans. Biliary dyskinesia		Extract	Boucard <i>et al.</i> , 1966
	Antibiliary, the alkaloid in it increases			
	the secretion of bile, and also at times		Alkaloid	Bisset <i>et al.</i> , (2001)
	of increased pathologic bile, reduces its secretion			
	Antispasmodic			Ivancheva <i>et al.</i> , 1999
	Colicky pain affecting the gallbladder and biliary system, together with the gastrointestinal tract			Hentschol <i>et al.</i> , 1995
	Extracts inhibited formation of gallbladder calculi in animals		Extracts	Lagrange <i>et al.</i> , 1973
Liver	Hepatoprotective activity	Effect from each side of his body	Ethanollic extract	Uday Raj Sharma <i>et al.</i> , (2012)
	Cytoprotective effect		Alkaloids	Taborska <i>et al.</i> , (1996)
Cardiovascular	Cardiovascular activity		Alkaloid fractions	Gorbunov <i>et al.</i> , (1980)
	In dogs reduced ischaemia caused by experimental ligation of the circumflex artery			Rao <i>et al.</i> , (1998)
System	Effect	Description	Active compounds	References
	Hypotensive, bradycardic and sedative activities in small doses in animals		Protopine	Goetz <i>et al.</i> , (2009)
	Larger doses cause excitation and convulsions		Protopine	Preininger <i>et al.</i> , 1975
Cell	Cytotoxicity effects		Protopine	Saglam <i>et al.</i> , (2003) British

				Herbal	
Eye	The use as an eye lotion in conjunctivitis			Pharmacopoeia (BHP)	
Respiratory	Has been used in Afghanistan for the treatment of asthma			Delaveau (1980)	
Muscle	Antispasmodic activity on smooth muscle has been reported			Reynier <i>et al.</i> , (1977)	
Skin	Valuable agent in cutaneous eruptions such as eczema and psoriasis, in scabies and also in syphilis.		Syrup	Mir Heidari (1993)	
	In Iranian folk medicine in skin diseases.			Amin (1991)	
	Anti-scabies, anti-scorbite, anti-bronchite				
	Leprosy, scabs, tatters, and itches, and such like breakings-out of the skin		Juice or syrup or seed	Baker <i>et al.</i> , (1993)	
Immune system	The immunocompromised condition of chronic alcoholics		Chronic alcoholics	Baker <i>et al.</i> , (1993); Blank <i>et al.</i> , (1993)	
	Significant antioxidant activity			Memnune <i>et al.</i> , (2009); Howard (1987)	
	Reduced glutathione, a free radical scavenger, plays a key role in the activation of T cells and macrophages			Reynier <i>et al.</i> , (1977)	

Clinical Trials

One clinical study which cholelithiasis, hepatopathy and post operation cholecystectomy syndrome were dimensioned and shown that *F. officinalis* water extract could have therapeutic aspects on them⁴⁷. Another test showed very positive amphocholeretic effects on biliary syndrome by *F. officinalis* extract⁴⁸. The effect of water extract of fumitory on choleric activity has been measured by a clinical trial on 1969 by Heully *et al.*, (1969).

CONCLUSIONS

F. officinalis plant has been explored exhaustively for their phytochemical and pharmacological activities. From the foregoing

accounts, it is evident that *F. officinalis* plant has been used ethno-medicinally as a valuable therapeutic agent for a variety of diseases, as we have illustrated in this article. Moreover, numerous research works have proven its uses beyond the ethno-medicinal ones in experimental animals. Various compounds which were separated from this plant may be responsible for its pharmacological activities.

REFERENCES

- Orhan, I., Sener, B., & Musharraf, S.G. (2010). Antioxidant and hepatoprotective activity appraisal of four selected *Fumaria* species and their total phenol and flavonoid quantities. *Exp Toxicol Pathol*, 64(3), 205-209.

2. Rajagopal, P.L., Jasmin, R., Sreejith, K.R., Premaletha, K., & Aneeshia, S. (2018). *Fumaria* species-An elaborative pharmacological review. *Br J Med Health Res*, 1(01), 22-41.
3. Popova, M.E., Simánek, V., Dolejs, L., Smysl, B., & Preininger, V. (1982). Alkaloids from *F. parviflora* and *F. kralikii*. *Planta Med*, 45:120-2.
4. Rahman, A.R., Bhatti, M.K., & Choudhary, M.I. (1992). Chemical constituent of *F. indica*. *Fitoterapia*, 13:129.
5. Hördegen, P., Hertzberg, H., Heilmann, J., Langhans, W., & Maurer, V. (2003). The anthelmintic efficacy of five plant products against gastrointestinal trichostrongylids in artificially infected lambs. *Vet Parasitol*, 17, 51-60.
6. Khattak, S.G., Giliani, S.N., & Ikram, M. (1985). Antipyretic studies on some indigenous Pakistani medicinal plants. *J Ethnopharmacol*, 14, 45-51.
7. Akhtar, M.S., Khan, Q.M., & Khaliq, T. (1984). Effect of *Euphorbia prostrata* and *F. parviflora* in normoglycaemic and alloxan-treated hyperglycemic rabbits. *Planta Med*, 50, 138-42.
8. Gowher, G. (2017). Pharmacological activity of *Fumaria indica* -A review. *J Phytopharmacol*, 6(6), 352-355.
9. Reader's Digest Field Guide to the Wild Flowers of Britain. (1981). Reader's Digest, p. 37.
10. Richard, F., Alastair, F., & Marjorie, B. (1974). *The Wild Flowers of Britain and Northern Europe*. London: Collins, p. 78.
11. Murphy, R.J. (2009). *Fumitories of Britain and Ireland*. BSBI Handbbok No. 12. London: Botanical Society of Britain and Ireland..
12. Taxon: *Fumaria officinalis* L." (<https://npgsweb.ars-grin.gov/gringlobal/taxonomydetail.aspx?id=70905>) ars-grin.gov. Retrieved 5 November 2017.
13. Wynne, P.M., Vine, J.H., & Amiet, R.G. (2004). Protopine alkaloids in horse urine. *J Chromatogr B Analyt Technol Biomed Life Sci*, 811, 85-91.
14. Sharma, U.R., Prakash, T., Surendra, V., Roopakarki, N., & Goli, D. (2012). Hepatoprotective activity of *Fumaria officinalis* against CCl₄-induced liver damage in rats. *Pharmacologia*, 3:20.
15. Hentschel, C., Dressler, S., & Hahn, E.G. (1995). [*Fumaria officinalis* (fumitory)-clinical applications]. *Fortschr Med*, 113, 291-2.
16. Gilani, H.A., Bashir, S., Janbaz, K.H., & Khan, A. (2005). Pharmacological basis for the use of *Fumaria indica* in constipation and diarrhea. *J Ethnopharmacol*, 96:585-9.
17. Dermaderosian, A., & Beutler, J.A. (2005). *The Review of Natural Products*, 4th ed. Lippincott Williams & Wilkins, Wolters Kluwer Health Inc, pp. 458-60.
18. Duke, J.A. (2002). *Handbook of Medicinal Herbs*. 2nd ed. Boca Raton: CRC Press.
19. *The Names of Plants*, 4TH Edition Gledhill, D. (1985-2008). *The Names of Plants*. United Kingdom: Cambridge University Press. p. 443.
20. Howard, M. (1987). *Traditional Folk Remedies* (century, 1987). pp.142-3.
21. Lombardi Satriani, R. (1951). *Credenze popolari calabresi*. Napoli: Edizioni De Simone.
22. Preininger, V. (1975). The pharmacology and toxicology of the papavera- ceae alkaloids. In: RHF Manske (ed): *The Alkaloids XV*. London: Academic Press. pp. 207-61.
23. Dulger, B., & Gonuz, A. (2004). Antimicrobial activity of certain plants used in Turkish traditional medicine. *Asian J Plant Sci*, 3, 104-7.
24. Boucard, M., & Laubenheimer, B. (1966). Action du nébulisat de fumeterre sur le débit biliaire du rat. *Therapie*, 21:903-11.
25. Reynier, M., Lagrange, E., & Godard, F. (1977). Action du nebulisat de Fumeterre officinal sur la musculature lisse. [*Fumaria officinalis* spray action on smooth muscles.] *Travaux de la Société de Pharmacie de Montpellier*, 37(2), 85-102.
26. Lagrange, E., & Aourousseau, M. (1973). Effect of spray-dried product of *Fumaria officinalis* on experimental gall bladder lithiasis in mice. *Ann Pharm Fr*, 31, 357-62.
27. Gruenwald, J., Brendler, T., Jaenicke, C., LaGow, B., editors. (2007). *PDR for Herbal Medicines*, 4th ed. Montvale: Thomson PDR, p. 340.
28. Duke, J.A. (2002). *Handbook of Medicinal Herbs*. 2nd ed. Boca Raton: CRC Press.
29. Denden, S., Braham, W., Gorcii Aloui, S., Lakhdar, R., Kahloun, H., Mahdouani, K., Knani, J., Haj Khelil, A. (2010). Analyse biologique et phytothérapie Clinique de l'asthme dans une population de la Tunisie centrale. *Acta Horticulturae*, 853, 391-6.
30. Nazarizadeh, A., Mikaili, P., Moloudizargari, M., Aghajanshakeri, S.H., & Javaherypour, S. (2013). Therapeutic uses and pharmacological properties of *Plantago major* L. and its active constituents. *J Basic Appl Sci Res*, 3(9), 212-21.
31. Zacharewicz, M., Chorazy, W., Mossor, S., & Zacharewicz, J. M. (1979). Treatment of cholecystopathies by *Fumaria* nebulisate. *Wiener medizinische Wochenschrift (1946)*, 129(8), 221-224.
32. Bisset, N.G., & Wichtl, M. (2001). *Herbal Drugs and phytopharmaceuticals*, 2nd ed. London: CRC Press, pp. 214-6.
33. Ivancheva, S., & Stantcheva, B. (2000). Ethnobotanical inventory of medicinal plants in Bulgaria. *J Ethnopharmacol*, 69(2), 165-72.
34. Hentschol, C., Dressler, S., & Hahn, E.G. (1995). *Fumaria officinalis* (fumitory)-clinical applications. *Fortschr Med*, 113, 291-2.
35. Sharma, U.R., Prakash, T., Surendra, V., Roopakarki, N., Goli, D. (2012). Hepatoprotective activity of *Fumaria officinalis* against CCl₄-induced liver damage in rats. *Pharmacologia*, 3:20.

36. Taborska, E., Bochorakova, H., Sousek, J., Sedmera, P., Vavreckova, C., Simanek, V. (1996). Alkaloids of *Fumaria densiflora* DC. Coll Czech Chem Commun, 61, 1064-72.
37. Gorbunov, N.P., Sukhanov, A.A., & Bolotova, M.F. (1980). Pharmacological correction of myocardial ischemia and arrhythmias in reversible coronary blood flow disorders and experimental myocardial infarct in dogs. Kardiologiia, 20: 84-7.
38. Rao, K.S., & Mishra, S.H. (1998). Antihepatotoxic activity of monomethyl fumarate isolated from *Fumaria indica*. J Ethnopharmacol, 60, 207-13.
39. Goetz, P., Ghedira, K., & Le Jeune, R. (2009). *Fumaria officinalis* L. (Fumariaceae). Phytotherapie, 7: 221-5.
40. Saglam, H., & Arar, G. (2003). Cytotoxic activity and quality control determinations on *Chelidonium majus*. Fitoterapia, 74: 127-9.
41. Bradley, P.R. editor. (1992). British Herbal Compendium. Fumitory-*Fumariae herba*. Bournemouth: British Herbal Medicine Association, pp. 102-4.
42. Delaveau, P. (1980). Fumeterre. Les actualites pharmaceutiques, 172, 33-4.
43. Mir Heidari, H. (1993). Encyclopedia of medicinal plants of Iran. Islamic Culture Press. Tehran (in Persian).
44. Amin, G.H. (1991). Popular medical plants of Iran. Pharmacologyonline, 3, 385-93.
45. Baker, R.C., & Jerrells, T.R. (1993). Recent developments in alcoholism: immunological aspects. Recent Dev Alcohol, 11: 249-71.
46. Blank, S.E., Pfister, L.J., Gallucci, R.M., & Meadows, G.G. (1993). Ethanol-induced changes in peripheral blood and splenic natural killer cells. Alcohol Clin Exp Res, 17, 561-65.
47. Fiegel, G. (1971). Die amphocholoretische Wirkung der *Fumaria officinalis*. Z Allgemeinmed Landarzt, 34, 1819-20.
48. Roux, M. (1977). Le nebulisat de fumeterre dans la régulation des troubles de intestinaux chroniques d'origine biliaire. Gazette Médicale de France, 84(34), 3794-8.
49. Heully, F., Drouin, P., Laurant, J., Bas, M., & Mauuary, G. (1969). Appreciation Des propriétés cholérétiques du nebulisat de *Fumaria officinalis* par le tubage duodenal minute. Ann Med Nancy, 8: 295-9.