

Original Research Article

“To Assess the Quality of Life among Cancer Patients in Relation To Type of Treatment (Chemo Therapy vs. Radio Therapy)”

H. N. Ashikur Rahaman^{1*}¹Registrar, Dept. of Clinical Oncology, Enam Medical College & Hospital, Savar, Dhaka, Bangladesh**Article History**

Received: 14.04.2021

Accepted: 19.05.2021

Published: 26.06.2021

Journal homepage:<https://www.easpublisher.com>**Quick Response Code**

Abstract: Background: With the advent of treatment like chemotherapy and radiotherapy, survival rates of many cancers have increased, but patient may experience side effects from the disease itself and/or from treatment for the disease which can eventually hinder patient's quality of life. **Aim and Objective:** To assess the quality of life among cancer patients in relation to type of treatment (chemo therapy vs. radio therapy). **Material and methods:** It was a cross sectional, descriptive and hospital based study. Total duration of study was 6 months (January 2020-Jun 2020), conducted in Dept. of Clinical Oncology, Enam Medical College & Hospital, Savar, Dhaka, Bangladesh. A total of 113 cancer patients undergoing treatment were selected as study subjects out of which 64 were undergoing chemotherapy and 49 were undergoing radiotherapy. Participants were interviewed by a validated questionnaire. **Results:** A total of 113 cancer patients were included in the study, of which 67 (59.3%) were females and 46 (40.7%) were males. None of the participants had above average or significantly high quality of life. 22.1%, 54%, 21.9% had average, below average and significantly poor quality of life respectively. Patients undergoing radiotherapy had a comparatively higher quality of life than patients undergoing chemotherapy (p value <0.05). Among patients undergoing chemotherapy, those who have undergone 3 or more cycle had better quality of life than those with less than 3 cycle (p value <0.05). **Conclusion:** Our study showed that majority of cancer patients undergoing treatment had poor quality of life and among them patients undergoing chemotherapy had lower quality of life compared to patients undergoing radio therapy.

Keywords: Cancer, Chemotherapy and Radiotherapy, Quality Of Life.

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

I. INTRODUCTION

Cancers are the leading causes of morbidity and mortality worldwide with approximately 14.1 million new cases and 8.2 million cancer related deaths and five-year prevalence of 32.6 million cancers [1]. IARC world cancer report 2014 estimates indicate a substantive increase to 19.3 million new cancer cases by 2025 [2]. Globally most common cancers diagnosed in 2012 were lung, breast and colorectum and most common causes of cancer death were cancers lung, liver and stomach [3]. Bangladesh, at 142 million people, is the 9th most populous country in the world. There are 13 to 15 lakh cancer patients in Bangladesh, with about 2 lakh patients newly diagnosed with cancer each year [4, 5]. As an overview, lung cancer and mouth and oropharynx cancer rank as the top 2 prevalent cancers in males. Other types of cancers commonly noted include esophagus cancer, stomach cancer, lymphomas, and multiple myeloma. In women, cervix uteri cancer and breast cancer are most prevalent. Cancers of oral cavity, lungs, & stomach were common cancers in men and breast, cervix & colorectum in women [4]. In

Bangladesh, cancers account for 6% of the NCD related DALYs [5]. The diagnosis of cancer and its treatment have a major impact on every aspects of patient's quality of life. Quality of life is an individual's perception of their position in life in the context of the culture and value system in which they live and in relation to their goals, expectations, standards and concerns [6]. It is a broad concept incorporating an individual's physical health, psychological state, level of independence, social relationships, personal beliefs and her/his relationships to salient features of the environment [7]. Although chemotherapy and radiotherapy have enhanced better prognosis in many cancers, they are not without side effects. The development of treatment related side effects are the most significant disadvantage of chemotherapy. It is a concentrated and repeated treatment drug regimen and has many adverse reactions including gastro intestinal, musculoskeletal and constitutional symptoms like nausea, vomiting, loss of appetite, constipation, diarrhea, fever and fatigue. The most apparent and emotionally challenging side effect is complete falling

of hair. Besides, it requires extended periods of treatment and repeated admission to hospital. Side effects with radiotherapy are mainly anemia and fatigue due to bone marrow suppression, skin reaction, gastro intestinal symptoms and sleep problems. Most side effects generally go away within few days of finishing treatment. However, some side effects may continue after treatment is over because it takes time for healthy cells to recover from the effects of radiation therapy. Whatever be the treatment methods, the intensity of treatment and their adverse reactions, can eventually affect the quality of life of cancer patients. Hence quality of life assessment should be a part of cancer treatment which helps physicians to properly document the individual impact of such treatments and decide which treatment modality would be better for individual patients. In our study we assessed the difference of quality of life among cancer patients undergoing chemotherapy and radiotherapy and also the impact of number of chemotherapy cycles on patients' quality of life.

II. MATERIAL AND METHODS

It was a cross sectional study done at Dept. of Clinical Oncology, Enam Medical College & Hospital, Savar, and Dhaka, Bangladesh conducted between January 2020 to Jun 2020. All patients more than 18 yrs undergoing chemotherapy and radiotherapy were included in the study. Patients who were undergoing both chemotherapy and radiotherapy and those who were critically ill and didn't give consent were excluded from the study. Ethical clearance was taken from institutional ethical committee of RIMS. 64 patients who were undergoing chemotherapy and 49 patients who were undergoing radiotherapy were interviewed. The quality of life of patients was assessed using a QOL questionnaire designed under EORTC guidelines and validated in Indian scenario by Vidhubala E, et al. [8]. The questionnaire consisted of questions relating to 10 factors.

1. Factor 1 - physical well-being
2. Factor 2 - psychological well-being
3. Factor 3- self adequacy.
4. Factor 4- confidence in self ability
5. Factor 5- external support attained by the patient.
6. Factor 6- extent of pain experienced
7. Factor 7- mobility of the patients
8. Factor 8 -optimism and belief
9. Factor 9 -interpersonal relationship
10. Factor 10- self-sufficiency and independence

Likert-type four-point rating scale was used to elicit responses from the respondents ranged from a minimum score of 1 and maximum score of 4 for each question. The total scores were summed up and QOL was interpreted as follows: 88 and below=significantly poor QOL, 89-108=below average QOL, 109-132=average QOL, 133-144=above average QOL, above 144=significantly high QOL

STATISTICAL ANALYSIS

Data were entered in MS Excel and analysis was done with SPSS statistical software. Chi-square test was performed to assess the difference in QOL of the patients. $p < 0.05$ was considered significant.

III. RESULTS

A total of 113 patients participated in the study. Majority were females (59.2%) belonging to Muslim religion (69.0%). Most (49.6%) patients were in the age group 40-60 years, mean age being 48.41 ± 11.35 years. 29.2% were aged less than 40 years, and 21.2% were more than 60 yrs. 69% of participants were married. Majority (51.6%) of the patients had primary education, and 23.6 % were illiterate. only 6.2% were postgraduate. Most of the participants were housewives (46%) and 40.7% were daily wagers. Majority were from low socioeconomic class (38.1% from class 4 and 20.4% from class 5).

Table-1: Socio-demographic profile

variable	category	frequency	Percentage
Age	<40	33	29.2
	40-60	56	49.6
	>60	24	21.2
gender	Male	46	40.8
	female	67	59.2
religion	Muslim	78	69.0
	Hindu	25	22.1
	Christian	10	8.9
education	Illiterate	28	23.6
	primary secondary/	57	51.6
	higher secondary	21	18.6
	Graduates/post graduates	7	6.2
occupation	House wife	52	46
	Daily wagers	46	40.7
	Business	9	8
	Student	2	1.8

	service	4	3.5
Marital status	Married	78	69
	Un married	12	10.6
	Divorced/separated	2	1.8
	Widow/widower	11	9.7
Socio-economic status(modified BG Prasad)	Class 1	11	9.6
	Class 2	15	13.3
	Class 3	21	18.6
	Class 4	43	38.1
	Class 5	23	20.4

Table-2: Treatment * Quality of life (QOL), n=113

Treatment * Quality of life (QOL), n=113						
		QOL			Total	Pearson chi-square test
		Average QOL	Below average QOL	Significantly poor QOL		P value
Treatment	Radiotherapy	14(28.6%)	29(59.2%)	6(12.2%)	49(43.4%)	Chi-square=6.97. df= 2 P =.031
	Chemotherapy	11(17.2%)	32(50%)	21(32.8%)	64(56.6%)	
Total		25(22.1%)	61(54%)	27(23.9%)	113(100%)	

In the present study we didn't get any participant with significantly high or above average quality of life. Majority of the patients, i.e. 61(54%) were leading below average quality of life and 27 patients (23.9%) were having significantly poor quality of life. only 25 patients (22.1%) had above average quality of life. Out of 113(100%) participants, 49(43.4%) were undergoing radiotherapy and 64 (56.6%) were undergoing chemotherapy. Among

patients undergoing radiotherapy, 14%, 59.2% and 12.2% were leading average, below average, and significantly poor quality of life respectively. Among chemotherapy group, majority (54%) had significantly poor quality of life. 50% were leading below average quality of life and only 17.2% had average quality of life. Patients undergoing radiotherapy had a significantly better quality of life than those undergoing chemotherapy (p=.031).

Table-3: Different domains of QOL and theirs scores.

Factors	Treatment	Score				Pearson Chi - Square Test
		4	3	2	1	
FACTOR 1	RT	10(20.4%)	21(42.9%)	12(24.5%)	6(12.2%)	Chi square= 8.330 P=.04
	CT	7(10.9%)	16(25%)	27(42.2%)	14(21.9%)	
FACTOR 2	RT	5(10.2%)	12(24.5%)	17(34.7%)	15(30.6%)	Chi square=.567 P=.904
	CT	7(10.9%)	17(26.6%)	18(28.2%)	22(34.3%)	
FACTOR 3	RT	20(40.8%)	15(30.6%)	8(16.3%)	6(12.3%)	Chi square=11.633 P=.009
	CT	12(18.8%)	14(21.9%)	18(28.1%)	20(31.2%)	
FACTOR 4	RT	8(16.3%)	16(32.7%)	14(28.6%)	11(22.4%)	Chi square=1.374 P=.712
	CT	14(21.9%)	19(29.7%)	21(32.8%)	10(15.6%)	
FACTOR 5	RT	15(30.6%)	18(36.7%)	12(24.5%)	4(8.2%)	Chi square=9.564 P=.023
	CT	9(14%)	16(25%)	27(42.2%)	12(18.8%)	
FACTOR 6	RT	8(16.3%)	13(26.5%)	19(38.8%)	9(18.4%)	Chi square=.126 P=.989
	CT	12(18.7)	17(26.6%)	24(37.5%)	11(17.2%)	
FACTOR 7	RT	22(44.9%)	18(36.7%)	6(12.3%)	3(6.1%)	Chi square=14.811 P=.002
	CT	14(21.9%)	17(26.6%)	28(43.7%)	5(7.8%)	
FACTOR 8	RT	20(40.8%)	17(34.7%)	8(16.3%)	4(8.2%)	Chi square=15.521 P=.001
	CT	10(15.6%)	16(25%)	21(32.8%)	17(26.6%)	
FACTOR 9	RT	3(6.1%)	7(14.3%)	18(36.7%)	21(42.9%)	Chi square=6.973. P=.042
	CT	11(17.2%)	13(20.3%)	27(42.2%)	13(20.3%)	
FACTOR 10	RT	21(42.8%)	16(32.7%)	9(18.4%)	3(6.1%)	Chi square=17.082 P=.001
	CT	11(17.2%)	14(21.9%)	21(32.8%)	18(28.1%)	

Patients undergoing radiotherapy showed better score in most of the domains of quality of life.

Physical well-being, social well-being, mobility, self-sufficiency and self-adequacy (factors 1,3,5,7,8,9 and

10) were better in radiotherapy group compared to chemotherapy group($p < 0.05$). However psychological

well-being, confidence and pain experienced showed no significant difference (factors 2, 4, and 6).

Table-4: Chemotherapy cycle * QOL, n=64

Chemotherapy cycle * QOL, n=64		QOL			Total	Pearson chi-square test
		Average QOL	Below average QOL	Significantly poor QOL		P value
Chemotherapy cycle	<3 cycles	3(4.7%)	9(14.1%)	13(20.3%)	25(39.1%)	Chi square=6.854 df=2 p=.032
	>_3 cycles	8(12.5%)	23(35.9%)	8(12.5%)	39(60.9%)	
Total		11(17.2%)	32(50%)	21(32.8%)	64(100%)	

Out of 64(100%) patients undergoing chemotherapy, 25(39.1%) had undergone less than 3 cycles and 39(60.9%) had undergone 3 or more cycles. Those patients who had taken 3 or more cycles had significantly better quality of life than those who had taken less than 3 cycles($p < 0.05$)

IV. DISCUSSION

In cancer care, “global well-being” including physical, emotional, mental, social, and behavioral components is the main surrogate objective apart from concluding cure [9]. Over the last decade, clinicians have accepted that while survival and disease-free interval are critical factors for cancer patients, overall quality-of life is fundamental to understand the impact of cancer upon the patient especially when the aims of treatments are palliative rather than curative [10]. In our study, we observed that none of our participants had above average or significantly higher quality of life. Most of them had a below average and poor quality of life. Similar results were found in a study conducted in tertiary care hospital, Dhaka, Bangladesh [11]. Beside these, some other major cancer programs have taken place for early detection of breast, cervical and oral cancer by Bangladesh Government and NGOs such as ICDDR’B, BRAC, Ahsania Mission Cancer Hospital, BSMMU, Bangladesh Cancer Society, Ashic Foundation, Amader Gram, AK Khan Healthcare Trust, CANSUP, Oncology club etc. Piloting of cervical cancer vaccination has recently been completed. Improving the cancer scenario overnight is not an easy task but policy makers may become interested and push this agenda forward, if the huge health impact and economic loss caused by cancer become evident to them. Besides, Bangladesh has accepted reduction of cancer morbidity and mortality targets set by United Nations and World Health Organization as a part of global non-communicable disease prevention agreement. Our study was conducted in an oncology clinic among those cancer patients who were undergoing chemotherapy or radiotherapy and we found that those patients who were undergoing chemotherapy had a significantly lower quality of life compared to those undergoing radiotherapy. A similar study by Awring M. Raof, et al. [12] showed the same results.

By analyzing each domain of quality of life separately in both groups, we found that patients undergoing radiotherapy showed better score in most domains. Radiotherapy had no significant effect on their overall physical well-being and working capacity. On the other hand physical well-being of patients undergoing chemotherapy was unsatisfactory. It may be because the toxicity and intensity of chemotherapy treatment regime and their adverse effects like nausea, vomiting, tiredness, alopecia were interfering with their daily activities, mobility and self-sufficiency. In radio therapy patients such side effects are comparatively less and are not severe enough to hamper their routine activities and mobility. Psychological well-being was similar in both the groups. Many patients from both the groups were having sadness and depression and were worried about their reduced economic status. A study by Chaturvedi S [13] also showed that psychological domain had no correlation with mode of treatment of cancer. Majority of patients undergoing chemotherapy were not satisfied with the external and family support they were getting but social well-being in radiotherapy patients was satisfactory. Pain was experienced by both the groups and chemotherapy patients experienced more pain but the difference was not statistically significant. We also observed that patients undergoing radiotherapy were more optimistic compared to patients undergoing chemotherapy. Present study also showed that the patients who had undergone more than 3 cycles of chemotherapy were having a better quality of life. Ali Dehkordi [14] also found in a study among cancer patients that chemotherapy cycles improved patients „quality of life. This could be because, the symptoms related to cancer gets better with successive cycles and also patients get adjusted to the treatment and their side effects. In another study done in Sweden [15] also, it was seen that chemotherapy improved quality of life in biliary and pancreatic cancer patients. Contradictory to our findings, a study conducted in US [16] among cancer patients undergoing chemotherapy showed that the quality of life of cancer patient did not improve with chemotherapy. It could be because the sample population selected was end stage cancer patients who were near death. In our study participants were from different ethnic, educational and religious background

from both rural and urban areas. The findings of our study showed that there was no correlation between quality of life and socio-demographic characteristics like age, gender, marital status socioeconomic class, education, occupation etc. Studies conducted by dehkordi [14], Vedat I [17] and Nematollahi [18] also showed similar results.

V. CONCLUSION

The most commonly used treatment method chemotherapy is more troublesome for patients compared to radiotherapy due to its side effects especially in the beginning of treatment as patients take time to get adjusted with the treatment and its side effects. Though cure and survival are the main aims of cancer treatment physicians should also focus more on patient's quality of life to offer a better productive life to them.

REFERENCES

1. World cancer fact sheets -2014 by IARC/WHO.
2. GLOBOCAN. (2012). Estimated Cancer Incidence, Mortality and Prevalence Worldwide in 2012.
3. Ferlay. (2015). Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN; 2012. *Int.J.Cancer*, 136:60-64.
4. National Cancer Control Strategy and Plan of Action 2009-2015. Directorate General of Health Services. Ministry of Health and Family Welfare, Bangladesh. [Last accessed on 2013 May 02]. Available from: http://www.ban.searo.who.int/LinkFiles/Publication_Cancer_Strategy.pdf.
5. Islam, A., Eden, T. (2013). Brief report on pediatric oncology in Bangladesh. *South Asian J Cancer*. 2:105-6. [PMC free article] [PubMed] [Google Scholar].
6. WHOQOL. (1997). Measuring quality of life, 4:1-3
7. The WHOQOL Group. (1995). World Health Organization. Quality of Life Assessment (the-WHOQOL): position paper from the World Health Organization. *SocSci Med*.
8. Vidhubala. (2005). Validation of quality of life questionnaire for patients with cancer. *Indian Journal of Cancer*, 42(3).
9. Harminder, S. (2014). Quality of Life in Cancer Patients Undergoing Chemotherapy in a Tertiary Care Center in Malwa Region of Punjab. *Indian journal of palliative care*, 20(2):116-118
10. Billingham, L.J., Abrams, K.R., Jones, D.R. (1999). Methods for the analysis of quality of life and survival data in health technology assessment. *Health technol Assess*, 3:1-152
11. Hussain, S.M.(2013). Comprehensive update on cancer scenario of Bangladesh. *South Asian J Cancer*. 2013;2(4):279-284.
12. Awring, M., Raoof. (2015). Quality of life among cancer patients treated with chemotherapy or radiotherapy in Erbil city. *Cancer and clinical oncology*, 4:19-27
13. Chaturvedi, S. (2003). What's important for quality of life to Indians in relation to cancer? *Indian J Palliat Care*, 9; 62- 70.
14. Ali, Dehkordi. (2009). Quality of life in cancer patients undergoing chemotherapy. *OMJ*, 24:204-207
15. Glimelius, B. (1996). Chemotherapy improves survival and quality of life in advanced pancreatic and biliary cancer. *Ann Oncol*, 7(6): 593-600
16. Holly. G., Pigersonl. (2015). Chemotherapy use, performance status, and quality of life at the end of life. *JAMA Oncol*, 6; 778-784
17. Vedat, I., Perihan, G., Seref, K. (2001). Improving quality of life in patient's with non small lung cancer: research experience with gemetabine. *Eur J Cancer*; 33:8-13
18. Nematollahi, A. (2006). Quality of life in Iranian women with breast cancer. PhD Thesis, Tehran University of Medical Sciences.

Cite This Article: H. N. Ashikur Rahaman (2021). "To Assess the Quality of Life among Cancer Patients In Relation To Type of Treatment (Chemo Therapy vs. Radio Therapy)". *East African Scholars J Med Sci*, 4(5), 124-128.