

Original Research Article

“Fresh versus Frozen Embryo Transfers-A Randomised Controlled Trial Study”

Ahmed B^{1*}, Angel², Sharma S³¹Fellow, International Fertility Center, Delhi, India²Senior Embryologist, International Fertility Center, Delhi, India³Fellow, International Fertility Center, Delhi, India**Article History**

Received: 19.05.2021

Accepted: 24.06.2021

Published: 29.06.2021

Journal homepage:<https://www.easpublisher.com>**Quick Response Code**

Abstract: Introduction: The freeze-all strategy was inferior to the conventional strategy in terms of cumulative ongoing pregnancy rate per woman. IVF without transfer of fresh embryos, thus with frozen-thawed embryo transfer only (freeze-all strategy), is increasingly being used in clinical practice because of a presumed benefit. It is still unknown whether this new IVF strategy increases IVF efficacy. **Materials and Methods:** A single-centre, open label, two arm, parallel group, randomised controlled superiority trial was conducted. The trial was conducted between January 2019 and March 2021 in the India. The intervention was one IVF cycle with frozen-thawed blastocyst transfer(s) versus one IVF cycle with fresh and frozen-thawed blastocyst transfer(s). The primary outcome was cumulative ongoing pregnancy resulting from one IVF cycle within 12 months after randomisation. Couples were allocated in a 1:1 ratio to the freeze-all strategy or the conventional strategy with an online randomisation programme just before the start of down-regulation. Participants were subfertile couples with any indication for IVF undergoing their first IVF cycle, with a female age between 18 and 43 years. Differences in cumulative ongoing pregnancy rates were expressed as relative risks (RR) with 95% CI. All outcomes were analysed following the intention-to-treat principle. **Results:** Two-hundred-and-five couples were randomly assigned to the freeze-all strategy (n ¼ 102) or to the conventional strategy (n ¼ 102). The cumulative ongoing pregnancy rate per woman was significantly lower in women allocated to the freeze-all strategy (19/102 (19%)) compared to women allocated to the conventional strategy (32/102 (31%); RR 0.59; 95% CI 0.36–0.98). **Conclusion:** Our results indicate that there might be no benefit of a freeze-all strategy in terms of cumulative ongoing pregnancy rates. The efficacy of the freeze-all strategy in subgroups of patients, different stages of embryo development, and different freezing protocols needs to be further established and balanced against potential benefits and harms for mothers and children.

Keywords: Endometrial receptivity, Endometrial receptivity analysis (ERA), Fresh embryo transfer (ET), Frozen embryo transfer (FET).

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

I. INTRODUCTION

IVF without fresh embryo transfer, thus with frozen-thawed embryo transfer only (freeze-all strategy), is increasingly being used in clinic practice in women undergoing IVF in an attempt to increase their chances of a pregnancy [1]. The underlying rationale is to avoid possible side effects of ovarian stimulation on endometrial receptivity during the initial treatment cycle by postponing embryo transfer to a subsequent cycle without ovarian stimulation [2-5]. In women at risk for ovarian hyperstimulation syndrome (OHSS), the freeze-all strategy has been applied for years, but with the aim to prevent OHSS [6-8]. Thus far, six randomised controlled trials (RCTs) have compared

cumulative live birth rates between a freeze-all strategy and a conventional strategy in women with a high risk of OHSS [9], in good prognosis women based on the number of follicles [10,11], in women with polycystic ovary syndrome (PCOS) [12], and in young women without PCOS [13,14]. A recent systematic review and meta-analysis of these studies showed that there was no difference between the strategies in cumulative live birth rate, i.e. the proportion of women achieving a live birth following the transfer of fresh or frozen embryos from one single cycle with ovarian stimulation [15]. The risk of OHSS for these groups of women was significantly lower in the freeze-all strategy [15]. Before the [15], systematic review and the largest study

included in this review [12], were available, we initiated a single-centre RCT comparing cumulative ongoing pregnancy rates per woman in a freeze-all strategy with frozen-thawed blastocyst transfer to a conventional strategy with fresh and frozen-thawed blastocyst transfer, as evidence of a benefit was lacking. In contrast to the other RCTs conducted thus far, we included sub fertile couples with. Any IVF indication, independent of the number of follicles or available embryos.

II. MATERIALS AND METHODS

We conducted a single-centre, open label, two arm, parallel group, randomised controlled superiority trial between January 2019 and March 2021. Women between 18 and 43 years of age who were scheduled for their first IVF cycle and who had no previous failed IVF cycles, in either International Fertility Center, Delhi, India were eligible for inclusion. All laboratory procedures were carried out in the couples treated in received ovarian stimulation and oocyte retrieval in embryo transfer International Fertility Center, Delhi, India. The use of this so-called transport-IVF is routine practice in the India. The protocol involves shipment of the follicular fluids immediately after oocyte retrieval in a controlled environment at 37°C to the IVF laboratory in the collaborating hospital. Data over the years show no difference in terms of pregnancy rates between transport-IVF centres and centres in which oocytes are not transported. Couples undergoing a preim-plantation genetic diagnosis cycle or undergoing a modified natural cycle were not included, as were couples with an HIV, hepatitis B virus or hepatitis C virus infection since all these cycles' required modified IVF protocols. The study protocol was approved by the Central Committee on Research Involving Human Subjects in the India. The institutional review boards of International Fertility Center, Delhi, India provided local approval. All couples in this trial provided written informed consent.

Randomisation and Masking

We randomised women with an online randomisation programme during their visit to plan the first IVF cycle, just before the start of down-regulation, using block randomisation with a maximum block size of six, stratified for age (18 years through 35 and 35 through 43 years), and study centre (International Fertility Center, Delhi, India). Couples were allocated in a 1:1 ratio to the freeze-all strategy or the conventional strategy. The randomisation programme generated a unique study number with allocation code after entry of the patient's date of birth and randomisation date. The gynaecologists, embryologists and the researchers who analysed the data could not access the randomisation sequence. Blinding of couples, clinicians and embryologists was not possible due to the nature of the comparison under study.

IVF protocol

Pituitary down-regulation was achieved with a long GnRH agonist protocol with or without oral contraceptive pill (OCP) pre-treatment. Ovarian stimulation was conducted with hMG india or recombinant FSH in women with PCOS starting from the seventh day without OCP. The starting dose depended on the antral follicle count (AFC) [16]. Ovulation was triggered with 5000 or 10 000 IU hCG india. We adhered to a single embryo transfer policy for women below 38 years of age and a double embryo transfer policy for women of 38 years of age and above, if two or more embryos were available.

Conventional strategy

We transferred embryos at Day 5 of culture. The morphologically best embryo(s) was transferred first. All surplus embryos were cryopreserved on Day 6 of culture. Women with a fresh transfer had luteal support with 600 mg vaginal micronized utrogestan (Utrogestan Besins Healthcare, Healthcare, and Utrecht in India) and continued this until the pregnancy test, 14days after the fresh transfer. If not pregnant, frozen embryo transfer was scheduled in artificial cycles with oral estrogen India and vaginal micronized progesterone supplementation (Utrogestan Besins Healthcare, Utrecht the India). If the endometrium had reached 8 mm on vaginal ultrasound, women started vaginal micronized progesterone (Utrogestan Besins Healthcare, Utrecht, and the India) and thawing and transfer were scheduled. Women continued estrogen and progesterone supplementation until 11½weeks gestation. If not pregnant, a subsequent artificial cycle was started.

Freeze-all strategy

Embryos were cryopreserved on Day 6 of culture. After oocyte retrieval, women waited for their menstruation and started an artificial cycle with oral estrogen and vaginal micronized progesterone supplementation, similar to the frozen embryo transfer in the conventional strategy. If not pregnant, a subsequent artificial cycle was started. See Supplementary methods for details of the IVF protocol, including embryo culture, embryo grading and details on the artificial frozen embryo transfer cycle.

End of study

End of study was the achievement of an ongoing pregnancy, transfer of all embryos derived from the first oocyte retrieval within 12months, or being 12months after randomisation regardless of any remaining supernumerary cryopreserved embryos. A pregnancy test was performed on serum 2weeks after blastocyst transfer and we considered a serum hCG >2IU/L as positive. We confirmed clinical and ongoing pregnancies by ultrasonography at 7 and 12weeks of gestation, respectively. If the ultrasound at 12weeks of gestation was not performed in the participating centres,

and if data on live births were missing, we contacted the couples by phone or postal mail to obtain data.

Outcomes

The primary outcome was cumulative ongoing pregnancy per woman after one cycle [17]. An ongoing pregnancy was defined as visible foetal cardiac activity at ultrasound from 12 weeks of gestation onwards, or a pregnancy that resulted in a live birth. Secondary outcomes were time to pregnancy defined as the time to ongoing pregnancy from the date of randomisation to the date of embryo transfer that led to an ongoing pregnancy, live birth defined as the delivery of a live foetus 20 weeks of gestation, clinical pregnancy defined as the presence of at least one intrauterine gestational sac at 7 weeks of gestation, and biochemical pregnancy defined as serum hCG >2 IU/l. Safety outcomes were OHSS, multiple pregnancy, premature birth and congenital abnormalities. We also report on miscarriage rate, ectopic pregnancy rate and birth weight of the children born.

Sample size

The cumulative ongoing pregnancy rate per started cycle among women in the two centres after one cycle of IVF was ~20% at the time of designing the study [18]. The null hypothesis assumed no difference while the alternative hypothesis assumed that pregnancy rates after the freeze-all strategy would be higher than after the conventional strategy. We expected a cumulative ongoing pregnancy rate after one cycle of 40% in the freeze-all strategy and of 20% in the conventional group. In a superiority design with a 5% significance level, 80% power and a two-sided test, we needed 164 evaluable couples. To account for a 15% loss to follow-up, we planned to include 193 couples.

STATISTICAL ANALYSIS

Analyses followed the intention-to-treat principle. We assessed the primary outcome cumulative ongoing pregnancy per randomised woman. Natural conceptions that occurred after randomisation but before IVF treatment or between frozen-thawed embryo cycles were included in the analysis of the primary outcome. If women had an on-going pregnancy from an embryo derived from the first cycle but transferred beyond 12 months after randomisation, these pregnancies were registered but not included in the analysis as primary outcome. We estimated differences in the binary outcomes as relative risks (RR) with 95% CI. The Chi-square test was used for categorical data, as appropriate. For continuous outcomes, we calculated mean and SD or median with range, and we evaluated differences using ANOVA or Mann-Whitney U tests, where appropriate. We constructed Kaplan-Meier curves to estimate the cumulative probability of an ongoing pregnancy over time and we analysed differences between the Kaplan-Meier curves with the log-rank test for significance. All statistical analyses were performed using SPSS (version 21.0, IBM,

Armonk, NY, USA). We considered P-values <0.05 statistically significant.

III. RESULTS

Between January 2019 and March 2021, 205 couples were randomly assigned: 103 couples allocated to the freeze-all strategy and 102 couples to the conventional strategy (Fig. 1). The reasons for not completing one cycle were similar between strategies and were primarily owing to personal reasons of the couple. One woman was randomised although she fulfilled one of the exclusion criteria. We excluded this woman from the analyses. In total, 204 couples were included in the analysis. Follow-up ended in Jun 2021. The baseline characteristics were similar between the two strategies (Table I).

Outcomes

Cumulative ongoing pregnancy was achieved in 19 couples (19%) in the freeze-all strategy and in 32 couples (31%) in the conventional strategy leading to an RR of 0.59 (95% CI 0.36–0.98) (Table II). The cumulative live birth rate (RR 0.62; 95% CI 0.34–1.04) between the two strategies was not significantly different. The cumulative clinical pregnancy rate (RR 0.58; 95% CI 0.36–0.95) and the cumulative biochemical pregnancy rate (RR 0.6; 95% CI 0.38–0.87) between the strategies were significantly lower in the freeze-all strategy. After the first embryo transfer, ongoing pregnancy (RR 0.32; 95% CI 0.15–0.68), live birth (RR 0.32; 95% CI 0.14–0.71), clinical pregnancy (RR 0.35; 95% CI 0.17–0.71) and biochemical pregnancy rates (RR 0.43; 95% CI 0.25–0.74) were significantly lower in the freeze-all strategy. In subsequent embryo transfers, the RR for the freeze-all strategy versus the conventional strategy was 2.00 (95% CI 0.71–5.65) for ongoing pregnancy, 2.00 (95% CI 0.71–5.65) for live birth, 2.00 (95% CI 0.71–5.65) for clinical pregnancy and 1.11 (95% CI 0.47–2.62) for biochemical pregnancy. The time to ongoing pregnancy was significantly higher in the freeze-all strategy (46.9 weeks versus 39.6 weeks, log-rank $P=0.02$; Fig. 2). Additional secondary outcomes are shown in Table III. No woman developed OHSS in the freeze-all strategy and three women developed OHSS in the conventional strategy. There was no significant difference in miscarriage rate between the freeze-all and conventional strategy. None of the women had an ectopic pregnancy in the freeze-all strategy and one woman had an ectopic pregnancy, i.e. cervical pregnancy, in the conventional strategy. None of the women had a multiple pregnancy in the freeze-all strategy and three women had a multiple pregnancy in the conventional strategy. There was no difference between the two strategies in birth weights after the first embryo transfer ($P=0.29$) or subsequent embryo transfers ($P=0.91$). In addition, there were no differences in the total birth weights lower than 2500 g and in preterm births before a gestational age of 37

weeks between the strategies. Congenital abnormalities did not occur in either group.

Treatment and embryological characteristics

The treatment characteristics were similar between the strategies, with exception of the endometrium thickness, which was significantly thinner in the freeze-all strategy than in the conventional

strategy (9.1 ± 2.1 versus 10.9 ± 2.7 ; $P < 0.0001$; Supplementary Table SI). The cumulative embryo implantation rate per woman (resulting in a gestational sac) was significantly lower in the freeze-all strategy than in the conventional strategy (RR 0.49; 95% CI 0.31–0.77). For the first embryo transfer, the implantation rate was significantly lower in the freeze-all strategy than in the

Table-I: Baseline characteristics of the included couples in a randomised controlled trial of fresh or frozen embryo transfer

		Freeze all	Conventional
Women randomised (n)		102	102
International Fertility Treatment centre		102	102
Maternal age (year)		35.2 ± 4.7	35.1 ± 4.5
Paternal age (year)		37.5 ± 6.2	38.6 ± 7.5
Duration of subfertility (year)		3.0 ± 2.1	3.2 ± 2.4
Primary diagnosis of subfertility			
	Tuba factor	10 (10)	11(11)
	Anovulation	2 (2)	3(3)
	Endometriosis	1 (1)	1(1)
	Cervix factor	1 (1)	0(0)
	Male subfertility	56 (55)	54 (53)
	Unexplained	28 (28)	25 (25)
	Mixed female and male factor	4 (4)	8 (8)
Sperm donor			
	Yes	10 (10)	5 (5)
	No	92 (90)	96 (95)
Parity		0.22 ± 0.6	0.27 ± 0.6
Smoking			
	Yes	16 (16)	16 (16)
	No	81 (79)	79(77)
Education			
	Primary school	2 (2)	1 (1)
	Intermediate vocational education	23 (23)	33 (32)
	Higher general education	3 (3)	6 (6)
	Higher education/university	56 (55)	42 (41)
	Not reported	18 (18)	20 (20)
Mean BMI (kg/m2)		23.6 ± 3.6	25.1 ± 5.0
Antral follicles		9.4 ± 6.8	11.6 ± 8.9

AMC Academic Medical Centre; international fertility center, delhi, india, Onze Lieve Vrouwe Gathuis. Data presented as means ± SD or number (%) conventional strategy (RR 0.26; 95% CI 0.13–0.51). For the subsequent embryo transfers, there was no

significant difference in the implantation rate between the freeze-all strategy and the conventional strategy (RR 1.83; 95% CI 0.71–4.77) (Supplementary Table SII).

Table-II: Effectiveness of the embryo transfer strategies: pregnancy outcomes

		Freeze all (n 5102)	Conventional (n 5102)	Relative risk	95% CI		P-value
Ongoing pregnancy ^(a)	Cumulative	19(19)	32 (31)	0.59	0.36	0.98	0.036*
	First ET	8 (8)	25 (25)	0.32	0.15	0.68	0.001*
	Subsequent ETs	10 (10)	5(5)	2.00	0.71	5.65	0.180
Live birth ^(b)	Cumulative	18 (18)	29 (28)	0.62	0.37	1.04	0.067
	First ET	7 (7)	22 (22)	0.32	0.14	0.71	0.003*
	Subsequent ETs	10 (10)	5 (5)	2.00	0.71	5.65	0.180
Clinical pregnancy	Cumulative	19 (19)	33 (33)	0.58	0.36	0.95	0.027*
	First ET	9 (9)	26 (25)	0.35	0.17	0.71	0.002*
	Subsequent ETs	10 (10)	5 (5)	2.00	0.71	5.65	0.180
Biochemical pregnancy	Cumulative	24 (24)	42 (41)	0.57	0.38	0.87	0.007*
	First ET	15 (15)	35 (34)	0.43	0.25	0.74	0.001*
	Subsequent ETs	10 (10)	9 (9)	1.11	0.47	2.62	0.810

Data presented as number (%). In the freeze-all strategy, one woman underwent elective termination of a pregnancy at 22weeks of gestation because of trisomy 21. In the conventional strategy, one woman terminated the gestation because of a cervical pregnancy and two

women had spontaneous miscarriages at 14 and 16weeks of gestation. (a)Cumulative pregnancy rates include pregnancies without IVF. (b)Not all couples who achieved ongoing pregnancy ended in a live birth. *P-value < 0.05. ET, embryo transfer.

Table-III: Secondary outcomes of the randomised controlled trial

		Freeze all (n 5102)	Conventional (n 5102)	Relative risk	95% CI		P-value
OHSS (a)		0 (0)	3 (3)				
Miscarriage (b)	Cumulative	14 (14)	20 (20)				
	First ET	10 (10)	16 (16)	0.63	0.30	1.31	0.21
	Subsequent ETs	4 (4)	6 (6)	0.67	0.19	2.29	0.52
Ectopic pregnancy (c)		0	1 (1)				
Multiple pregnancy	Cumulative	0	3 (3)				
	First ET	0	2 (2)				
	Subsequent ETs	0	1 (1)				
Birthweight (d)	Cumulative	3528 (518)	3283 (704)				0.20
	First ET	3561 (257)	3249 (743)				0.29
	Subsequent ETs	3479 (672)	3440 (579)				0.91
Birthweight	<2500 g	1 (1)	5 (5)				
Preterm birth	<37 week	2 (2)	5 (5)				

Data presented as mean±SD or number (%). (a)Ovarian hyper stimulation syndrome (OHSS) that required hospitalisation. (b)All miscarriages occurred before 20weeks of gestation. (c)One woman had a cervical pregnancy. (d)Data of the first ET include two twin live borns, data in the subsequent ETs include one twin live born, both in the conventional group, ET, embryo transfer.

IV. DISCUSSION

Our study suggests that a freeze-all strategy with transfer of frozen thawed blastocysts significantly reduces cumulative ongoing pregnancy rate per woman after one cycle compared to a conventional strategy with transfer of fresh and frozen-thawed blastocysts in a general IVF population where all patients undergoing IVF, including those with a poor prognosis, were included. There was also evidence of reduced cumulative live birth, and significantly reduced cumulative clinical pregnancy, and cumulative biochemical pregnancy. When comparing our results to other studies, we cannot compare our findings with RCTs that only reported ongoing pregnancy or live birth after the first transfer [19, 20]. However, in comparison to other RCTs that reported cumulative results, the success rate in the freeze-all strategy with frozen-thawed blastocyst transfer seems lower in the present study [21, 13]. These other RCTs have shown comparable cumulative live birth and ongoing pregnancy rates in both arms (Cochrane systematic review of Wong et al. [15]. In the freeze-all arm, our cumulative ongoing pregnancy rate was 19% compared to 50–70% in the other RCTs. The cumulative ongoing pregnancy rate of the conventional arm was also lower than found in other studies but the difference was less.

There could be several explanations for these differences. The RCTs performed by [21], and Shapiro et al. [22] n¼122 included high responders, defined as women at risk for OHSS and as women with more than 15 antral follicles at baseline ultrasound, respectively. The other RCT performed by [22], n¼137 included normal responders, defined as women that had 8–15 antral follicles at baseline ultrasound that were expected to have a good response to ovarian stimulation. The study by Chen et al. [12], n¼1508 included women with PCOS in their first IVF cycle with four or more oocytes available after oocyte retrieval. Two studies included a general, but younger, IVF population (mean age 32 and 28 years, respectively) where the study by Vuong had to have a minimum of one available high-grade embryo, and the study by Wei a minimum of at least four high-grade quality embryos available before a women would be randomly assigned to one of the two groups [13]. Our study consisted of an unselected population undergoing IVF, including (expected) poor responders. The selection criteria of normal and high responders or of young women used in previous studies may represent women with a relatively better prognosis and are therefore more likely to have one or more (good quality) blastocysts for transfer resulting in a pregnancy. An exploratory post-hoc analysis from our data in women with more than two top-quality embryos on Day 3 of culture, indeed shows that the cumulative ongoing pregnancy rates per woman are similar between the strategies in our study (freeze all 39% (14/36) versus conventional 42% (15/36), RR 0.93, 95% CI 0.53– 1.63). This suggests that the results of our study are in line with the outcomes of previous RCTs when analysing a more comparable patient population [15]. It also made clear that the majority of

the included patients in our study (65% 132/204) did not have at least three top-quality embryos on Day 3 of culture, which could possibly be explained by the patient characteristics of an IVF population where expectant management for at least 12 months is promoted before IVF treatment is started. Apart from differences in study population, several other factors were different between our trial and those previously conducted that could possibly explain the differences in results. First, the timing of randomization differed; either after retrieval [21, 13] or before retrieval (our study). Second, the timing of embryo cryopreservation; either at the two pronuclei stage [21], the cleavage stage [12], or the blastocyst stage and our study on Day 6, compared to our fresh blastocyst transfer on Day 5 [14]. Third, the method of cryopreservation; either slow freezing or vitrification [21, 12]. And fourth, the embryo developmental stage at transfer; cleavage stage [21], morula or blastocyst stage (our study) or blastocyst stage only. Although the single Centre design did not allow for analyses considering differences in study protocols and clinical performance, one could speculate that our vitrification-thawing programme might have underperformed. However, based on the high survival rates of embryos after thawing in this study (96%, Supplementary Table SII), we conclude that we performed our vitrification-thawing correctly and that the frozen-thawed embryos were viable and were not affected in their morphological development by freezing. However, we cannot exclude possible damage in frozen-thawed embryos as a result of the vitrification procedure that cannot be determined by morphological development. Since we used fixed criteria, i.e. morulae and blastocysts, for embryo transfer and cryopreservation regardless of the developmental stage of embryos, this could have led to transfer and cryopreservation of embryos of suboptimal quality. We intentionally lowered the morphological cut-off criteria for cryopreservation by freezing embryos with identical morphologic criteria on Day 6 in the freeze-all strategy, and on Day 5 in the conventional strategy. We opted for this fixed transfer protocol, so embryos in the freeze-all strategy were not discarded that otherwise would have been transferred in the conventional strategy outside the RCT setting. Cryopreserving and later transferring slower developing embryos, i.e. morulae or early blastocysts, on Day 6 may be unfavourable for the freeze-all strategy when compared to the same embryo developmental stage, i.e. morulae or early blastocysts, on Day 5 in the conventional strategy. This could have caused a skewed comparison. In both study groups, multiple embryos were still cryopreserved at the end of the study (95 embryos in the freeze-all arm and 103 embryos in the conventional arm). We choose to set a limit on the time frame during which each included patient could remain in the study, in order to limit the duration of the study. The primary endpoint was achievement of an ongoing pregnancy, transfer of all embryos derived from the first oocyte retrieval within 12months, or being 12months

after randomisation regardless of any remaining supernumerary cryopreserved embryos. The vast majority of the embryos that were still cryopreserved at the end of the study were from couples that achieved an ongoing pregnancy during the study, which seems to suggest that the chosen period did not affect the cumulative pregnancy rate in our study to a large extent as these women have already been counted as being pregnant. These embryos remain available to the couples should they request an additional child. We used a fixed protocol for endometrium preparation regardless of the developmental stage of embryos and thus a possible mismatch in embryo development and endometrium development may have occurred. In general, the quality of the endometrium is difficult to determine, but seems an unlikely cause of the lower pregnancy rate in the freeze-all strategy since the endometrium was never thinner than 7mm, a cut-off that has been related to a lower chance of pregnancy. In addition, we used an accepted common protocol for endometrium preparation for transfer of frozen-thawed embryos [22]. The decision to offer the freeze-all strategy should be balanced against potential benefit and harm for mother and child. It is known that the freeze-all strategy is associated with an increased risk of hypertensive disorders in pregnancy and large for gestational age as well as a higher birth weight in singleton babies. In conclusion, our findings suggest that a freeze-all strategy of blastocysts does not result in a better cumulative ongoing pregnancy rate in unselected couples undergoing IVF compared to a fresh transfer strategy.

V. CONCLUSION

Our results indicate that there might be no benefit of a freeze-all strategy in terms of cumulative ongoing pregnancy rates. The efficacy of the freeze-all strategy in subgroups of patients, different stages of embryo development, and different freezing protocols needs to be further established and balanced against potential benefits and harms for mothers and children.

REFERENCES

1. Barnhart, K. T. (2014). Introduction: are we ready to eliminate the transfer of fresh embryos in in vitro fertilization?. *Fertility and sterility*, 102(1), 1-2.
2. Mastenbroek, S., Van Der Veen, F., Aflatoonian, A., Shapiro, B., Bossuyt, P., & Repping, S. (2011). Embryo selection in IVF. *Human Reproduction*, 26(5), 964-966.
3. Maheshwari, A., & Bhattacharya, S. (2013). Elective frozen replacement cycles for all: ready for prime time?. *Human reproduction*, 28(1), 6-9.
4. Wong, K. M., Mastenbroek, S., & Repping, S. (2014). Cryopreservation of human embryos and its contribution to in vitro fertilization success rates. *Fertility and sterility*, 102(1), 19-26.
5. Chida, S., & Mettler, L. (1989). Elective preembryo cryopreservation in ovarian hyperstimulation syndrome. *Journal of in vitro*

- fertilization and embryo transfer, 6(5).
6. Pattinson, H. A., Hignett, M., Dunphy, B. C., & Fleetham, J. A. (1994). Outcome of thaw embryo transfer after cryopreservation of all embryos in patients at risk of ovarian hyperstimulation syndrome. *Fertility and sterility*, 62(6), 1192-1196.
 7. Shaker, A. G., Zosmer, A., Dean, N., Bekir, J. S., Jacobs, H. S., & Tan, S. L. (1996). Comparison of intravenous albumin and transfer of fresh embryos with cryopreservation of all embryos for subsequent transfer in prevention of ovarian hyperstimulation syndrome. *Fertility and Sterility*, 65(5), 992-996.
 8. Devroey, P., Polyzos, N. P., & Blockeel, C. (2011). An OHSS-Free Clinic by segmentation of IVF treatment. *Human reproduction*, 26(10), 2593-2597.
 9. Ferraretti, A. P., Gianaroli, L., Magli, C., Fortini, D., Selman, H. A., & Feliciani, E. (1999). Elective cryopreservation of all pronucleate embryos in women at risk of ovarian hyperstimulation syndrome: efficiency and safety. *Human Reproduction*, 14(6), 1457-1460.
 10. Shapiro, B. S., Daneshmand, S. T., Garner, F. C., Aguirre, M., Hudson, C., & Thomas, S. (2011). Evidence of impaired endometrial receptivity after ovarian stimulation for in vitro fertilization: a prospective randomized trial comparing fresh and frozen-thawed embryo transfer in normal responders. *Fertility and sterility*, 96(2), 344-348.
 11. Shapiro, B. S., Daneshmand, S. T., Garner, F. C., Aguirre, M., Hudson, C., & Thomas, S. (2011). Evidence of impaired endometrial receptivity after ovarian stimulation for in vitro fertilization: a prospective randomized trial comparing fresh and frozen-thawed embryo transfer in normal responders. *Fertility and sterility*, 96(2), 344-348.
 12. Chen, Z. J., Shi, Y., Sun, Y., Zhang, B., Liang, X., Cao, Y., & Legro, R. S. (2016). Fresh versus frozen embryos for infertility in the polycystic ovary syndrome. *N engl j med*, 375, 523-533.
 13. Vuong, L. N., Dang, V. Q., Ho, T. M., Huynh, B. G., Ha, D. T., Pham, T. D., & Mol, B. W. (2018). IVF transfer of fresh or frozen embryos in women without polycystic ovaries. *New England Journal of Medicine*, 378(2), 137-147.
 14. Wei, D., Liu, J. Y., Sun, Y., Shi, Y., Zhang, B., Liu, J. Q., & Chen, Z. J. (2019). Frozen versus fresh single blastocyst transfer in ovulatory women: a multicentre, randomised controlled trial. *The Lancet*, 393(10178), 1310-1318.
 15. Wong, K. M., van Wely, M., Mol, F., Repping, S., & Mastenbroek, S. (2017). Fresh versus frozen embryo transfers in assisted reproduction. *Cochrane Database of Systematic Reviews*, (3).
 16. van Tilborg, T. C., Eijkemans, M. J., Laven, J. S., Koks, C. A., de Bruin, J. P., Scheffer, G. J., ... & Broekmans, F. J. (2012). The OPTIMIST study: optimisation of cost effectiveness through individualised FSH stimulation dosages for IVF treatment. A randomised controlled trial. *BMC women's health*, 12(1), 1-7.
 17. Braakhekke, M., Kamphuis, E. I., Dancet, E. A., Mol, F., Van Der Veen, F., & Mol, B. W. (2014). Ongoing pregnancy qualifies best as the primary outcome measure of choice in trials in reproductive medicine: an opinion paper. *Fertility and sterility*, 101(5), 1203-1204.
 18. NVOG. (2011). http://www.nvog.nl/Sites/Files/0000003049_IVF_cijfers_2011_per_kliniek.pdf
 19. Aghahosseini, M., Aleyasin, A., Sarfjoo, F. S., Mahdavi, A., Yaraghi, M., & Saeeadabadi, H. (2017). In vitro fertilization outcome in frozen versus fresh embryo transfer in women with elevated progesterone level on the day of HCG injection: an RCT. *International Journal of Reproductive BioMedicine*, 15(12), 757.
 20. Coates, A., Kung, A., Mounts, E., Hesla, J., Bankowski, B., Barbieri, E., & Munné, S. (2017). Optimal euploid embryo transfer strategy, fresh versus frozen, after preimplantation genetic screening with next generation sequencing: a randomized controlled trial. *Fertility and sterility*, 107(3), 723-730.
 21. Ferraretti, A. P., Gianaroli, L., Magli, C., Fortini, D., Selman, H. A., & Feliciani, E. (1999). Elective cryopreservation of all pronucleate embryos in women at risk of ovarian hyperstimulation syndrome: efficiency and safety. *Human Reproduction*, 14(6), 1457-1460.
 22. Shapiro, B. S., Daneshmand, S. T., Garner, F. C., Aguirre, M., Hudson, C., & Thomas, S. (2011). Evidence of impaired endometrial receptivity after ovarian stimulation for in vitro fertilization: a prospective randomized trial comparing fresh and frozen-thawed embryo transfer in normal responders. *Fertility and sterility*, 96(2), 344-348.

Cite This Article: Ahmed B *et al* (2021). "Fresh versus Frozen Embryo Transfers-A Randomised Controlled Trial Study". *East African Scholars J Med Sci*, 4(5), 133-139.